

RESEARCH ARTICLE

Open Access

Is the high-risk strategy to prevent cardiovascular disease equitable? A pharmacoepidemiological cohort study

Helle Wallach-Kildemoes^{1*}, Finn Diderichsen², Allan Krasnik³, Theis Lange⁴ and Morten Andersen^{5,6}

Abstract

Background: Statins are increasingly prescribed to prevent cardiovascular disease (CVD) in asymptomatic individuals. Yet, it is unknown whether those at higher CVD risk – i.e. individuals in lower socio-economic position (SEP) – are adequately reached by this high-risk strategy. We aimed to examine whether the Danish implementation of the strategy to prevent cardiovascular disease (CVD) by initiating statin (HMG-CoA reductase inhibitor) therapy in high-risk individuals is equitable across socioeconomic groups.

Methods:

Design: Cohort study.

Setting and participants: Applying individual-level nationwide register information on socio-demographics, dispensed prescription drugs and hospital discharges, all Danish citizens aged 20+ without previous register-markers of CVD, diabetes or statin therapy were followed during 2002–2006 for first occurrence of myocardial infarction (MI) and a dispensed statin prescription (N = 3.3 mill).

Main outcome measures: Stratified by gender, 5-year age-groups and socioeconomic position (SEP), incidence of MI was applied as a proxy for statin need. *Need-standardized statin incidence rates* were calculated, applying MI incidence rate ratios (IRR) as need-weights to adjust for unequal needs across SEP. Horizontal equity in initiating statin therapy was tested by means of Poisson regression analysis. Applying the need-standardized statin parameters and the lowest SEP-group as reference, a need-standardized statin IRR > 1 translates into horizontal inequity favouring the higher SEP-groups.

Results: MI incidence decreased with increasing SEP without a parallel trend in incidence of statin therapy. According to the regression analyses, the need-standardized statin incidence increased in men aged 40–64 by 17%, IRR 1.17 (95% CI: 1.14–1.19) with each increase in income quintile. In women the proportion was 23%, IRR 1.23 (1.16–1.29). An analogous pattern was seen applying education as SEP indicator and among subjects aged 65–84.

Conclusion: The high-risk strategy to prevent CVD by initiating statin therapy seems to be inequitable, reaching primarily high-risk subjects in lower risk SEP-groups.

Keywords: Statins to prevent cardiovascular disease, The high-risk strategy, Socioeconomic gradient, Horizontal equity

* Correspondence: hewk@sund.ku.dk

¹Centre for Healthy Aging, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, Copenhagen 1014, Denmark
Full list of author information is available at the end of the article

Background

A steep inverse relationship between socioeconomic position (SEP) and incidence of cardiovascular disease (CVD) has consistently been shown across high-income Western countries [1-3]. The social gradient has widened over the last decades [4,5] and is to a large extent mediated by the conventional risk factors (i.e., smoking, high blood pressure and serum cholesterol) – when evaluated in absolute terms [1,6-8]. This holds also for the most important CVD component, myocardial infarction (MI) [9].

As CVD is one of the leading causes of premature death in the Western world, preventive strategies are on political agendas, all focusing on the conventional risk-factors, either through their socio-cultural determinants (population (structural) strategies) or through individual behaviour/risk-factors, such as the high-risk strategy to prevent CVD in general practice. In the high-risk strategy (often termed ‘primary prevention’), asymptomatic individuals are screened to determine the need for preventive interventions, such as antihypertensives or lipid lowering drugs (almost exclusively statins). In the present study, we focus on statins (HMG-CoA reductase inhibitors), introduced in 1994 [10] to, reduce post-MI mortality in middle-aged men with hypercholesterolemia. Following subsequent randomised clinical trials (RCTs), recommendations for statins have broadened, including now also asymptomatic individuals (i.e., without established CVD or diabetes) irrespective of lipid-levels age and gender. “The question of ‘at what lipid level to initiate treatment’ has to be replaced by ‘at what cardiovascular risk should statins be started’” [11].

The high-risk strategy has been implemented in Denmark as an opportunistic screening strategy i.e. clients who show up in the general practitioner’s (GP) office may be screened for high CVD risk for possible prescription of preventive drugs. In line with the European guidelines and the European Systematic Coronary Risk Evaluation [12], Danish GPs are recommended to use a matrix of serum lipid and blood pressure levels (stratified by age, gender and smoking status) for identifying high-risk individuals, applying an estimated 10-year risk of fatal atherosclerotic events above 5% as high risk threshold. While risk thresholds and CVD end-points vary slightly according to country, all risk-score charts are based on the same risk-factor matrix, providing risk estimates based on data and risk-equations from historic cohort studies and RCTs [12-14].

The Danish health care system is mainly tax-financed and based on the egalitarian principle, with equal health care (access/treatment/outcome) for equal (legitimate) health care needs [15] – also called horizontal equity [16]. While authorized GP services are free, prescription drugs require patient co-payment. Based on decisions by

an authority under the Ministry of Health, the actual amount of reimbursement depends on whether a particular drug is reimbursable and the actual reimbursement schedule for reimbursable drugs [17]. The current ‘need-dependent’ reimbursement schedule (introduced March 2000) has a number of reimbursement levels, the reimbursed percentage increasing stepwise with the individual’s annual drug expenditures. Reimbursement is based on the cheapest generic drug [17]. Despite near universal health care coverage in many European countries, income-related inequalities in the use of physician services have been observed [18-20]. In Denmark this holds true especially in regards to elective procedures [21] and services with co-payments, such as prescription drugs [22-24]. Yet, European health-care systems are under pressure due to increasing health care expenditures and the challenges of an ageing population [25,26], which includes shortage of GP’s partly due to the retirement of the baby-boom generation.

There is an ongoing debate about the high-risk strategy [14,27-29], encompassing allocation of scarce health care resources [28] and “The strategy of preventive medicine”, by Geoffrey Rose [30], i.e., the high-risk strategy versus the population strategy [29,31,32]. As reduction of social inequalities in health is a central goal in WHO and EU programmes [33,34], it is also being debated whether or not these strategies will reduce inequalities in CVD [35,36]. A range of studies have explored inequalities in utilisation of CVD drugs, but without explicitly taking need-determined measures into account, some focusing on regional or socioeconomic inequalities [37-40], others restricting analyses to individuals with the same medical condition [23,41]. In a study of equity in statin prescribing by GPs in the UK, the authors explore to what extent prescribing variations in different primary care trusts are associated with the frequency of CVD admissions and socio-demographic characteristics [38]. Assuming implicitly equal needs across these groups, the results of the UK study could indicate inequitable statin prescribing. Yet, inequality in health care delivery can only be interpreted as inequity if legitimate need-determined inequalities are taken into account [19,20].

In the present study, we focus on initiation of preventive statin therapy in the high-risk strategy as implemented in Denmark. Due to the social gradient in incidence of CVD we expect an increasing need for CVD preventive drugs with decreasing SEP i.e. unequal needs across socioeconomic groups. In line with other studies focusing on equity in health care delivery [22,42], we assume that equity will be met if care is provided proportionally to the need. To our knowledge no studies has explored to what extent the high-risk strategy to reduce CVD is equitable.

The aim of this study was to examine whether the Danish implementation of the strategy to prevent CVD by initiating statin therapy in high-risk individuals is equitable across socioeconomic groups; hypothesising that this high-risk strategy will not adequately reach groups with a lower SEP, characterised by having a higher risk of CVD.

Methods

Data source and participants

From nationwide Danish registers maintained by the National Board of Health and Statistics Denmark, we retrieved individual-level information on dispensed prescription drugs, hospital discharges, dates of death or emigration, and socioeconomic indicators. Data were linked by means of a unique encrypted person-identifier, allowing authorised researchers to follow individuals in multiple individual-level registries hosted in Statistics Denmark. Register-based studies in Denmark do not require approval by an ethics board.

A cohort corresponding to all Danish citizens aged 20 + without previous register markers of CVD, diabetes or statin treatment were followed, during 2002–2006, in the registries, for dispensed statin prescriptions and the first occurrence of MI (N = 3.3 mill). We applied two different SEP indicators: disposable family income and highest attained education (as of 2002). Table 1 shows the characteristics of the cohort of asymptomatic individuals, by gender, age and highest attained education, demonstrating that historical information on education is poorly covered among persons older than 75.

From the Danish National Patient Registry, we retrieved information on patient discharge from non-psychiatric hospitals since 1977. Records include the admission and discharge dates, discharge diagnoses according to the International Classification of Diseases (ICD), 8th revision until 1993, and 10th revision thereafter (9th revision has not been used in Denmark) along with codes for diagnostic and surgical procedures. We included main and secondary diagnoses for admitted patients and patients in ambulatory care. From the Registry of Causes of Death, we retrieved date and cause of death (primary and contributory).

Information on dispensed prescription drugs was retrieved from the Danish National Prescription Registry (DNPR), containing full information since 1996 on all out-of-hospital purchases of prescription drugs at Danish pharmacies including those of nursing home residents [43]. Records include the person identifier, date of dispensing, and the Anatomical Therapeutic Chemical (ATC) classification code of the dispensed drug [44]. From the DNPR we retrieved information on dispensed cardiovascular drugs (ATC groups C and B01) and antidiabetics (A10).

To identify asymptomatic individuals, we applied historical register data on in/out-patient diagnoses and procedures along with dispensed prescription drugs as register-markers for a range of CVD conditions, including ischemic heart disease (IHD) with or without myocardial infarction (MI), stroke, a range of other atherosclerotic conditions, and diabetes. We define asymptomatic individuals as individuals without register-markers of CVD or diabetes, as defined in a recent publication [45].

Table 1 The cohort as of January 1, 2002, according to gender, age and educational level

Gender	Educational Level ^a	20-39		40-54		55-64		65-74		75-84		85+		All
		N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	
Male	1 (Basic)	215,607	(32)	120,082	(24)	79,745	(30)	53,249	(42)	25,022	(43)	131	(1)	493,837
	2	229,083	(34)	210,144	(42)	108,985	(41)	46,910	(37)	13,966	(24)	0	(0)	609,087
	3	134,755	(20)	50,034	(10)	15,949	(6)	5,071	(4)	1,746	(3)	0	(0)	207,555
	4 (High)	80,853	(12)	95,065	(19)	50,505	(19)	17,750	(14)	6,401	(11)	262	(2)	250,836
	Missing	20,213	(3)	25,017	(5)	10,633	(4)	2,536	(2)	10,474	(18)	12,722	(97)	81,594
	All	673,773		500,343		265,816		126,784		58,190		13,115		1,638,021
Female	1 (Basic)	171,753	(27)	141,319	(29)	109,669	(40)	91,760	(59)	52,508	(55)	320	(1)	567,329
	2	190,837	(30)	160,811	(33)	95,961	(35)	40,437	(26)	15,275	(16)	0	(0)	503,320
	3	146,308	(23)	38,984	(8)	13,709	(5)	4,666	(3)	1,909	(2)	0	(0)	205,576
	4 (High)	114,502	(18)	126,700	(26)	46,609	(17)	17,108	(11)	5,728	(6)	0	(0)	310,647
	Missing	19,084	(3)	24,365	(5)	8,225	(3)	3,111	(2)	20,048	(21)	31,359	(98)	106,192
	All	636,122		487,306		274,173		155,525		95,469		31,999		1,680,594
Both	Total	1,309,895		987,646		539,989		822,298		101,359		45,114		3,318,615

Danish residents without register markers of CVD, diabetes and/or statin treatment.

a) Educational levels:

1) 7–10 years basic school, 2) Vocational training after basic school, 3) High school or 9/10 years basic school with 1 year higher education 4) More than two years higher education.

Study design

While measures such as the Gini-coefficient of inequality, concentration index and the slope index of inequalities provide means for quantifying the degree of for example income-related inequality in health or health care delivery [46], a measure combining potential inequalities both in health care delivery and health care needs is indispensable to quantify inequities in health care delivery – if needs also are unequal across strata. However, measuring the need for preventive health care is a challenge, as such needs not may be captured by for example self-rated health scales. We opted to apply a need-proxy analogous to the underlying presumption of the risk-score chart, namely a measure of CVD- incidence in the background population of asymptomatic individuals, i.e. without CVD, diabetes or statin therapy - stratified by gender, 5-year age groups and SEP indicator.

Due to the high validity of the diagnosis of MI in the Danish registries [47], we applied the incidence of MI as

need-proxy, using two alternative need-proxies in a sensitivity analysis: first stroke or MI as combined CVD endpoint and CVD as cause of death (MI, stroke or aortic aneurysm). Stratum specific MI-incidence rates were calculated, corresponding to number of incident MI cases (hospitalised or fatal) per 10,000 person-years at risk (PYR) during 2002–2006, censoring at death, emigration and register-markers of CVD, diabetes or statin therapy. Analogously, we calculated the ‘observed’ incidence of statin therapy (number redeeming the first statin prescription/10,000 PYR – censoring at death, emigration and register-markers of CVD or diabetes) and the combined MI-stroke endpoint. In order not to confine CVD-mortality to sudden CVD-death, CVD-mortality was calculated without censoring for new events of CVD or diabetes, covering also a longer span of time (2002–2008).

We applied a fixed SEP level corresponding to the beginning of the observation period (primo 2002). In order

Table 2 Observed and need-standardized incidence of statin treatment among asymptomatic individuals, according to gender, age-group and SEP indicators

Gender	SEP	40-54			55-64			65-74			75-84			
		INCOME ^a	Need	Statin incidence ^d		Need	Statin incidence		Need	Statin incidence		Need	Statin incidence	
			Weight ^c	Observed	St. dized ^e	Weight	Observed	St. dized	weight	Observed	St. dized	weight	Observed	St. dized
Male	1 (Low)	1.00	8.0	8.0	1.00	16.0	16.0	1.00	18.6	18.6	1.00	10.2	10.2	
	2	0.90	9.0	10.0	1.01	19.1	18.9	0.96	22.2	23.2	0.81	10.9	13.5	
	3	0.91	9.8	10.7	0.90	20.3	22.5	0.89	23.3	26.1	0.85	11.2	13.2	
	4	0.82	10.2	12.4	0.80	21.2	26.4	0.79	23.2	29.3	0.77	12.6	16.3	
	5 (High)	0.71	10.5	14.9	0.70	21.3	30.5	0.65	23.4	35.8	0.66	15.0	22.8	
Female	1 (Low)	1.00	8.3	8.3	1.00	23.5	23.5	1.00	28.2	28.2	1.00	14.1	14.1	
	2	0.80	8.3	10.4	0.85	25.2	29.8	0.85	29.4	34.6	0.90	15.3	17.1	
	3	0.64	8.4	13.1	0.64	24.7	38.8	0.90	28.8	32.0	1.02	15.6	15.3	
	4	0.60	8.5	14.1	0.57	23.1	40.5	0.69	29.2	42.0	0.82	17.7	21.6	
	5 (High)	0.49	7.6	15.6	0.31	20.7	67.0	0.44	28.2	63.4	0.64	17.5	27.5	
EDUCATION^b														
Male	1 (Basic)	1.00	9.4	9.4	1.00	18.7	18.7	1.00	20.5	20.5	1.00	11.6	11.6	
	2	0.81	10.2	12.6	0.89	20.6	23.2	0.97	23.9	24.6	0.94	14.6	15.6	
	3	0.59	8.3	14.1	0.75	18.5	24.6	0.82	22.1	27.0	0.76	13.9	18.2	
	4 (High)	0.50	8.8	17.8	0.66	19.9	30.2	0.65	23.8	36.7	0.72	15.7	21.8	
Female	1 (Basic)	1.00	9.9	9.9	1.00	25.6	25.6	1.00	28.9	28.9	1.00	17.3	17.3	
	2	0.63	8.8	14.0	0.68	23.8	35.2	0.79	30.0	38.1	0.82	20.2	24.7	
	3	0.43	6.0	14.0	0.47	18.7	39.6	0.54	25.2	46.3	0.68	17.2	25.3	
	4 (High)	0.34	6.2	18.2	0.38	19.1	50.8	0.53	26.4	50.2	0.65	19.0	29.5	

a) Quintiles of disposable family income within gender and 5-years age-groups – inflated average over 1996–2001.

b) Educational levels, cf. notes in Table 1.

c) Incidence Rate Ratio of myocardial infarction (MI-IRR) among asymptomatic individuals, applying lowest SEP group (within gender and age-group) as reference group.

d) Incidence of statin therapy among asymptomatic individuals: 1000*(Number incident statin dispensing/person years at risk).

e) Need-standardized statin incidence ('St.dized'): Observed statin incidence/MI-IRR.

to capture income fluctuations over time, we calculated the average annual income between 1996 and 2001 (inflated to 2000 purchasing power), divided into income quintiles within gender and age group. The highest attained educational levels as of 2002 were divided into four groups according to length of formal education, cf. Table 1. As the analysis covers a time span of five years, individuals were considered to belong to a fixed 5-year age-group (i.e., that of 2002).

To evaluate horizontal inequity in initiation of preventive statin therapy, we adjusted the observed incidence of statin therapy according to the different needs across SEP-groups, applying stratum specific MI-incidence as proxy for needs. By means of indirect standardisation, we calculated the expected incidence of statin therapy ('need-standardized statin incidence'), assuming that incidence of statin therapy must increase proportionally to the need across SEP-groups - for equity to be met. The need-standardized statin incidence was calculated as the observed statin incidence divided by the stratum specific need weights corresponding to the incidence rate ratio of MI (MI-IRR), Table 2. The denominator (PYR) of the observed statin incidence rather than the nominator (number of events) was need-standardized, dividing the observed PYR by MI-IRR.

Based on the need-standardized statin incidence parameters (observed statin events (counts), need-standardized PYR (exposure)), Poisson regression analyses were applied to test the overall horizontal equity across SEP. With the lowest SEP-group (and age group) as reference, a need-standardized statin-IRR > 1 translates

into horizontal inequity favouring the higher SEP-groups. The null hypothesis, horizontal equity, corresponds thus to statin-IRR = 1. We estimated a 'horizontal inequity gradient' (the overall linear trend) reflecting the increase in need-standardized statin-IRR for each increase in SEP. Owing to a gender and age specific pattern of both MI-incidence (need) and incidence of preventive statin therapy, we stratified the analyses according to gender and ages +/-65, cf. Figure 1.

Nonparametric bootstrapping [48] was applied to incorporate the precision of the need-weights in the confidence intervals (CI) of the need-standardized statin IRR. Based on 10,000 bootstrap replications, need-weights were calculated and applied in the Poisson regression analyses of need-standardized statin incidence parameters. Normal based 95% CI from the bootstrapping procedure were applied as CI for the point estimate for statin IRR calculated from the original data.

All analyses were performed using Stata Release 11.1 (StataCorp, College Station, TX, USA). Access to data was provided and secured through collaboration between the University of Copenhagen and Statistics Denmark. Register-based studies in Denmark do not require approval by an ethics board.

Results

Figure 1 depicts the incidence of MI and statin therapy among asymptomatic individuals during 2002–2006 according to age, stratified by gender. Both in men and women, the MI-incidence increases gradually with age, whereas statin incidence increases steeply until the age




Table 3 Result of Poisson regression analysis, testing horizontal equity across levels of socio-economic position (SEP) with differential needs

SEP indicator ^a	Age 40-64				Age 65-84			
	Male		Female		Male		Female	
Income Quintiles	IRR ^b	95% CI ^c	IRR	95% CI	IRR	95% CI	IRR	95% CI
1	1.00		1.00		1.00		1.00	
2	1.22	(1.10-1.35)	1.27	(1.04-1.55)	1.27	(1.05-1.54)	1.25	(1.04-1.51)
3	1.38	(1.29-1.48)	1.63	(1.39-1.92)	1.38	(1.17-1.63)	1.16	(0.93-1.45)
4	1.61	(1.47-1.76)	1.73	(1.40-2.15)	1.58	(1.36-1.84)	1.54	(1.27-1.86)
5	1.89	(1.70-2.10)	2.44	(1.86-3.21)	1.99	(1.68-2.35)	2.26	(1.80-2.80)
HIE gradient ^d	1.17	(1.14-1.19)	1.23	(1.16-1.29)	1.17	(1.14-1.20)	1.20	(1.14-1.27)
Educational level	IRR	95% CI	IRR	95% CI	IRR	95% CI	IRR	95% CI
1	1.00		1.00		1.00		1.00	
2	1.29	(1.17-1.42)	1.38	(1.27-1.49)	1.19	(1.07-1.33)	1.33	(1.13-1.56)
3	1.40	(1.18-1.66)	1.49	(1.33-1.65)	1.34	(1.19-1.52)	1.52	(1.23-1.89)
4	1.74	(1.51-2.01)	1.90	(1.72-2.09)	1.77	(1.59-1.97)	1.70	(1.47-1.97)
HIE gradient	1.19	(1.15-1.24)	1.24	(1.19-1.29)	1.21	(1.17-1.25)	1.21	(1.14-1.28)

a) Indicators of SEP: Quintiles of disposable income and 4 levels of formal education, cf. notes Table 1.

b) Poisson regression analyses of need-standardized statin Incidence Rate Ratio (IRR), applying need-standardized statin incidence parameters within gender and 5-year age-groups, based on indirect standardization with Incidence of myocardial infarction in the background population as need-weights, cf. Table 2.

c) Bootstrapping (10,000 reps.) is applied for 95% confidence intervals (CI).

d) Horizontal Inequity (HIE) gradient: The relative change in need-standardized statin incidence for each increase in SEP, i.e., the estimated overall linear trend.

of 65, decreasing markedly thereafter. While the MI-incidence is highest among men of all ages, the opposite is the case as regards statin incidence.

Table 2 shows that the need-weights (MI-IRR relative to the lowest SEP-group) are decreasing with increasing SEP – independently of gender and age categories. In men aged 55–64, the need (MI-incidence) in the highest income quintile is 70% of that in the lowest income quintile, in women the figure is 30%. Above the age of 75 the gradient is less pronounced. Analogously, when applying four educational levels as an indicator for SEP, the need in men aged 55–64 with the highest educational level is 70% of those with basic education – in

women the figure is 40%. Yet, while the observed statin incidence increases with increasing income in men, only, the need-standardized statin incidence increases steeply with increasing income in both genders – and more so among women due to the steeper gradient in MI-incidence.

Table 3 presents the results of the gender/age stratified Poisson regression analyses on need-standardized statin parameters. In men aged 40–64, the need-standardized statin incidence increases by 17% (IRR 1.17; 95% CI 1.14-1.19) for each increase in income quintile – corresponding to the horizontal inequity gradient (HIE-gradient). In women the HIE gradient is greater 23% (1.23;

Table 4 Sensitivity analysis of horizontal equity in incidence of preventive statin therapy: Three alternative need-weights (need-proxies) in the need-standardized analyses ^a: Myocardial infarction (MI), combined MI-stroke or CVD-death ^b

SEP indicator ^c	Age	Horizontal Inequity gradient (IRR) ^d dependent on need-weights					
		MI		MI-stroke		CVD-death	
		Male	Female	Male	Female	Male	Female
Income	40-64	1.17	1.23	1.18	1.15	1.39	1.36
	65-84	1.17	1.20	1.15	1.10	1.29	1.23
Educational level	40-64	1.19	1.24	1.18	1.11	1.33	1.22
	65-84	1.21	1.21	1.15	1.11	1.23	1.29

a) Need-standardized Poisson analyses see notes Table 3.

b) Myocardial infarction (MI) as need-weight in the basic analyses. Two alternative CVD need: Incidence of MI and stroke as combined endpoint during 2002–2006, and CVD-death (i.e., MI, stroke and aorta-aneurism) during 2002–2008.

c) Indicators of socio-economic position (SEP): cf. notes Tables 1 & 3.

d) Horizontal Inequity (HIE) gradient: The relative change in need-standardized statin incidence for each increase in SEP, i.e., the estimated overall linear trend.

1.16-1.29). Among individuals older than 65, the corresponding HIE-gradient is 17% (1.17; 1.14-1.20) and 20% (1.20; 1.14-1.27), in men and women, respectively.

The HIE-gradient concerning educational level reveals a similar pattern, i.e., 19% (1.19; 1.15-1.24) in men aged 30–64, and 24% (1.24; 1.19-1.29) in women.

Table 4 shows sensitivity analysis of the need-proxy, where the HIE-gradient is calculated applying the original need-proxy, MI incidence, as well as the two alternative need-proxies: MI/stroke as combined end-point and CVD as cause of death, respectively. Independently of need-weights (need-proxy) used, the sensitivity analysis revealed the same pattern of horizontal inequity, favouring the better-off. Yet, the HIE-gradient is attenuated when applying MI/stroke as a combined end-point. This is particularly true for women. Conversely, the HIE-gradient is enhanced when applying CVD-death as need-proxy – especially among men.

Discussion

Principal findings

Applying indirect standardisation and MI-incidence as a proxy for need, we developed a pharmacoepidemiological method to explore horizontal equity in initiation of preventive statin therapy across SEP-groups with unequal needs, adjusting the observed statin incidence according to relative needs across socio-demographic groups. Our study indicates that the high-risk strategy (as implemented in Denmark) to prevent CVD by initiating preventive statin therapy is inequitable, reaching primarily high-risk individuals in low-risk SEP-groups. The favouring of more advantaged groups holds for both genders, independently of applying income quintiles or educational level as SEP indicator. In men aged 30–64, the need-standardized statin incidence increased by 17% for each increase income quintile (the horizontal inequity gradient) – in women the increase was 21%.

Only among men, the *observed* incidence statin therapy tended to increase with increasing SEP, but due to a steeper social gradient in MI-incidence among women, the horizontal inequity gradient of initiating preventive statin therapy was steepest in women.

Strengths and limitations

Given the inverse relationship between SEP and CVD, the challenge of this study examining equity in the medicamental high-risk strategy to prevent CVD was twofold: firstly, to operationalize need and equity in CVD preventive drug therapy across SEP-groups with unequal needs, and secondly, to develop appropriate pharmacoepidemiological methods for testing horizontal equity.

Needs

We opted to apply nationwide register data on MI-incidence in the statin free and asymptomatic background population stratified by gender, age and SEP as need-proxy, instead of calculating individual-level CVD-risk based on survey information on CVD-risk factors and risk scoring,

This is a strength for at least three reasons:

- 1) The nationwide approach is without the well-known selection bias problems from cohort studies/surveys where people are invited to participate.
- 2) The risk-score charts generally have low predictive value – both at the individual and the group level. Various cohort studies indicate that standard risk-score charts tend to underestimate CVD-risk in worse-off groups, overestimating the risk in better-off groups [36,49,50], and attempts have been made to modify risk-score charts according to the actual background populations [51] and/or socioeconomic groups [49,50]. Yet, individual risk prediction is notoriously difficult [52,53], and as most CVD events occur in people with modest risk-factor values, overlapping with those seen in people without CVD, the appropriateness of applying individual risk-factor levels as a screening tool has been questioned [52,54,55]. The risk-score charts applied in Denmark seems, in fact, to have a very poor predictive value [56]. In the present study we applied risk (MI-incidence) at the sub-group level (gender, age and SEP) as need-proxy rather than risk at the individual level. Yet, individual risk estimates based on risk-scoring relies on risk calculated at the group level (age and gender) from historic survey data.
- 3) Register-based information on MI-diagnoses is regarded as valid in-hospital diagnosis information [47]. Thus, the estimated MI-incidences should reveal the actual SEP gradient in MI-incidence. However, due to potential differential symptom awareness and communication barriers, a social gradient in admission for MI may exist, leading to an underestimated MI-incidence in less advantaged groups. The better-off, on the other hand, may be more likely to prevent MI by means of invasive coronary procedures, leading to a potential underestimated MI-incidence here. Yet, including acute invasive coronary procedures as marker for MI revealed the same results (data not shown). Thus, we regard this bias of minor importance. In a sensitivity analysis, stroke/MI as a combined endpoint and CVD-death were tested as two alternative need-proxies. Independently of need-measure, the sensitivity analysis demonstrated

similar patterns of horizontal inequity. Yet, when applying the combined endpoint, the horizontal inequity gradient was – especially in women – attenuated, presumably due to a less pronounced social gradient in the incidence of stroke than in MI-incidence. Applying CVD-death as a need-proxy (corresponding to the risk measure applied in Danish risk-scoring charts), the horizontal inequity was enhanced. Yet the ‘validity’ of actual MI-incidence as the need-proxy rests upon the assumption that the need for statin therapy is a question of CVD-risk rather than a single risk-factor level – and partly on the finding that conventional risk-factors to a large extent mediate the social gradient in CVD-risk (in absolute terms). However, high cholesterol level has not consistently been found to increase with decreasing SEP [6,8,57], potentially explained by the rather imprecise cholesterol parameters applied: risk thresholds for cholesterol levels and measurement of total cholesterol without distinguishing between the various lipid-fractions. Conversely, other studies have demonstrated an association between the metabolic syndrome (caused by physical inactivity, unhealthy diets, and obesity) and elevated low-density lipid cholesterol (LDL) [58], indicating that the inverse social gradient in LDL may follow the social stratification in physical inactivity and obesity.

Pharmacoepidemiological method for testing equity

Analogous to studies within economic equity research [19,22,42], we applied indirect standardisation to evaluate horizontal equity in health care delivery. In a study on equity in US ambulatory care [42], the number of ambulatory visits was adjusted according to differential self-rated health. Applying the need-standardized counts of the dependent variable (ambulatory visits) and a continuous income variable as the explanatory variable, a ‘horizontal inequity index’ (i.e., a need-standardized concentration index) was estimated. In our pharmacoepidemiological approach, we calculated, instead, a need-standardized incidence rate of statin therapy. Applying need-standardized statin incidence parameters as a dependent variable and a SEP indicator as an ordinal explanatory variable (e.g., income quintiles), we estimated a *horizontal inequity gradient*. We consider this methodological analogy to be a strength. Yet, while it is intuitively reasonable to ‘adjust’ for differential health conditions when evaluating horizontal equity in ambulatory visits, it may be less obvious that incidence of preventive CVD-drug therapy should be proportional to the risk of disease for equity to be met.

Interpretation and comparison with other studies

While a range of studies have demonstrated *inequality* in prescription of CVD preventive drugs [23,59-61], no studies have examined and quantified inequities, including both prescribing patterns and needs in a nationwide perspective. In contrast to a Norwegian health survey study [40] showing a decreasing trend of incidence of statin treatment by increasing education in individuals without reported CVD or diabetes at baseline, we found almost the same incidence across educational groups among asymptomatic individuals, censoring for new onset of CVD/diabetes. The lack of censoring for onset of disease in the Norwegian study most likely explains the discrepancy between the studies, as lower SEP individuals are at higher risk of developing disease and may thereby be misclassified as free of CVD or diabetes when initiating statin treatment. Our finding that the high-risk strategy as implemented in Denmark seems to be inequitable may reflect both the poor predictive value of the applied risk-score charts and a selective uptake. The latter being an inherent consequence of applying an opportunistic screening strategy, where uptake depends on the client’s participation and the physician’s general judgement of her/his client. A so-called ‘healthy user effect’ has been shown in pharmacoepidemiological studies, indicating that preventive measures (e.g., vaccinations and drugs) tend to be used by population segments with a broad spectrum of healthier behaviours [62]. With the consistently shown social gradient in CVD in most Western countries, our findings are likely to be applicable in other settings applying an opportunistic screening strategy. Several studies have demonstrated a socioeconomic gradient in screening uptake [63], indicating both financial (due to drug co-payment) and psychosocial barriers in socially deprived groups [64]. Psychosocial barriers to CVD screening may include negative perceptions about screening tests, risk perceptions and the social stress associated with talking about unhealthy lifestyles with the GP – of higher SEP. Our findings may also reflect that high CVD risk in lower SEP at first hand is ‘attacked’ by encouraging individual lifestyle modifications.

In line with other studies [61,65] our study indicates that the high-risk strategy may widen the socioeconomic gradient in CVD owing to the inequitable uptake. However, any widening of the CVD incidence gradient depends on the outcome of therapy and not merely on initiation of therapy. Here two other issues are important: Differential adherence to therapy and differential outcome of therapy. In fact, long-term adherence to statin treatment is disappointing [66] – and is likely to depend on SEP, indication and experienced adverse effects (potentially weighted against the perceived beneficial effect). While the risk of life-threatening adverse effects is low (e.g., rhabdomyolysis), various degrees of muscle

side effects are not unusual, ranging from muscle weakness to rhabdomyolysis [67]. If both incidence and duration of therapy are lowest among less advantaged groups – the social gradient in prevalence and outcome of therapy is likely to be even steeper than the gradient found as to initiation of therapy. However, being exposed to multiple risk factors (known and unknown) acting in concert, socially disadvantaged groups may be more vulnerable to high LDL levels than the better-off. Hence, the *outcome* or beneficial effect of lifelong preventive statin therapy may be greater in less advantaged groups – provided adherence to therapy.

Unanswered questions and future research

Various incentives have been proposed to enhance adherence, requiring often GPs to be more actively involved [68,69]. In a forthcoming study on the same nationwide Danish data we explore potential socioeconomic differences in adherence to statin treatment in asymptomatic individuals.

The incidence of preventive statin treatment in this study was found to peak around the age of 65, and to decrease steeply hereafter. This pattern may reflect the widespread use of the risk-score charts, covering the age range of 40–65, potentially representing an issue of ageism. The finding that statin incidence is considerably higher in asymptomatic women than men although MI-incidence is higher in men (cf. Figure 1) may both reflect a consequence of an opportunistic preventive screening strategy and an overestimation of CVD-risk in Danish women, corresponding to the finding in a Norwegian study [70]. Both matters will require further research.

In contrast to the opportunistic screening strategy applied in Denmark, a universal screening programme to prevent CVD is actually being implemented in the UK [71]. Here, all asymptomatic individuals aged 40–74 are invited for risk-scoring and potential preventive statin therapy. Yet, there is an ongoing discussion about the programme covering both resource implications [72] and the dilemma between a targeted high-risk strategy with a high CVD risk threshold for initiating therapy with potentially non-generic more potent and effective statins versus a “population strategy” in which all invited individuals should be prescribed low-price preventive treatment (generic statins) [71] or eventually a Polypill (a combination of four CVD-preventive drug categories) [73,74], bypassing the risk-scoring charts of poor predicative value and potentially also subsequent GP visits. While recent cost-effectiveness reviews indicate a very high cost effectiveness of Polypill strategies [75], a pilot project may uncover equity concerns before implementation of a general screening strategy. Another strategy could be to focus the high-risk drug strategy on middle-aged asymptomatic men in whom the beneficial effect of preventive statin

treatment is best documented, testing various settings in order to reach lower SEP groups before implementation – potentially also adjusting the reimbursement system accordingly. Yet, by not controlling the causes of high CVD incidence (e.g., low cigarette prices; high levels of saturated and *trans*-fats, sugars, salt hidden in processed foods, and environmental factors) this ‘population’ strategy will be palliative and not radical as structural/population strategies tends to be [55]. Proposing a range of actions to be taking, a newly published Danish report “Health inequality – determinants and policies” [76] demonstrates that reducing health inequality is not primarily a health care task, but a complex task requiring coordinated efforts from different sectors (e.g. the education, social, healthcare, and employment sectors).

Conclusions and implications for policy and practice

Our study indicates that the high-risk strategy to prevent CVD by means of preventive statin therapy – as practiced in Denmark – is inequitable, primarily reaching high-risk individuals in low-risk groups, i.e., individuals in higher SEP-groups. The inequity is likely to be the consequence of using a screening tool with low predictive value and a screening programme with differential socioeconomic uptake. Provided long-term adherence and a beneficial effect of preventive statin therapy independent of SEP, the strategy may contribute to accentuating the inverse relationship between SEP and CVD. Facing the challenges posed by an ageing population, one might question to what extent scarce GP resources should be allocated for better-off, asymptomatic individuals.

Abbreviations

SEP: Socioeconomic position; CVD: Cardiovascular disease; RCTs: Randomised clinical trials; GP: General practitioner; DNPR: Danish National Prescription Registry; ICD: International Classification of Diseases; ATC: Anatomical Therapeutic Chemical classification system; IHD: Ischemic heart disease; MI: Myocardial infarction; PYR: Person-years at risk; IRR: Incidence rate ratio; HIE gradient: Horizontal inequity gradient.

Competing interests

The authors declare no conflicts of interest. Morten Andersen participates, however, in a research project on statins funded by AstraZeneca, Merck Sharp & Dohme and Pfizer.

Authors' contribution

All authors read and approved the final manuscript.

Authors' information

Finn Diderichsen, Allan Krasnik and Theis Lange contributed equally to the work.
Helle Wallach-Kildemoes, first and corresponding author.
Morten Andersen contributed as last author and principal co-author.

Acknowledgements

This study was financed partly by a grant from the Danish Health Foundation and partly by the Nordea Foundation, the latter financing the Centre for Healthy Aging. The register-based study was defined and performed entirely by the researchers. Register-based studies in Denmark do not require approval by an ethics board.

Author details

¹Centre for Healthy Aging, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, Copenhagen 1014, Denmark. ²Social Medicine, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, Copenhagen 1014, Denmark. ³Health Services Research, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, Copenhagen 1014, Denmark. ⁴Biostatistics, Department of Public Health, University of Copenhagen, Øster Farimagsgade 5, Copenhagen 1014, Denmark. ⁵Centre for Pharmacoepidemiology, Karolinska Institutet, Stockholm SE-171 77, Sweden. ⁶Research Unit for General Practice, Institute of Public Health, University of Southern Denmark, B. Winsløvs Vej 9A, Odense 5000, Denmark.

Received: 23 April 2012 Accepted: 20 July 2012

Published: 4 August 2012

References

- Lynch J, Davey SG, Harper S, Bainbridge K: **Explaining the social gradient in coronary heart disease: comparing relative and absolute risk approaches.** *J Epidemiol Community Health* 2006, **60**(5):436–441.
- Marmot MG, Smith GD, Stansfeld S, Patel C, North F, Head J, White I, Brunner E, Feeney A: **Health inequalities among British civil servants: the Whitehall II study.** *Lancet* 1991, **337**(8754):1387–1393.
- Mackenbach JP, Cavalears AE, Kunst AE, Groenhouf F: **Socioeconomic inequalities in cardiovascular disease mortality; an international study.** *Eur Heart J* 2000, **21**(14):1141–1151.
- Mackenbach JP, Bos V, Andersen O, Cardano M, Costa G, Harding S, Reid A, Hemstrom O, Valkonen T, Kunst AE: **Widening socioeconomic inequalities in mortality in six Western European countries.** *Int J Epidemiol* 2003, **32**(5):830–837.
- In *Trends in socioeconomic health differences between 1980–2005*. Edited by Palosuo H, Koskinen S, Lahelma E, Kostiaainen E, Prättälä R, Martelin T, Ostamo A, Keskimäki I, Sihto M, Linnanmäki E. Finland; 2009. http://www.stm.fi/c/document_library/get_file?folderId=39503&name=DLFE-8648.pdf.
- Lynch JW, Kaplan GA, Cohen RD, Tuomilehto J, Salonen JT: **Do cardiovascular risk factors explain the relation between socioeconomic status, risk of all-cause mortality, cardiovascular mortality, and acute myocardial infarction?** *Am J Epidemiol* 1996, **144**(10):934–942.
- Smith GD, Shipley MJ, Rose G: **Magnitude and causes of socioeconomic differentials in mortality: further evidence from the Whitehall Study.** *J Epidemiol Community Health* 1990, **44**(4):265–270.
- Singh-Manoux A, Nabi H, Shipley M, Gueguen A, Sabia S, Dugravot A, Marmot M, Kivimäki M: **The role of conventional risk factors in explaining social inequalities in coronary heart disease: the relative and absolute approaches to risk.** *Epidemiology* 2008, **19**(4):599–605.
- Manrique-Garcia E, Sidorchuk A, Hallqvist J, Moradi T: **Socioeconomic position and incidence of acute myocardial infarction: a meta-analysis.** *J Epidemiol Community Health* 2011, **65**(4):301–309.
- Scandinavian Simvastatin Survival Study group: **Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S).** *Lancet* 1994, **344**(8934):1383–1389.
- Ong HT: **The statin studies: from targeting hypercholesterolaemia to targeting the high-risk patient.** *QJM* 2005, **98**(8):599–614.
- Conroy RM, Pyörälä K, Fitzgerald AP, Sans S, Menotti A, De Backer G, De Bacquer D, Ducimetiere P, Jousilahti P, Keil U, Njolstad I, Oganov RG, Thomsen T, Tunstall-Pedoe H, Tverdal A, Wedel H, Whincup P, Wilhelmsen L, Graham IM: **Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project.** *Eur Heart J* 2003, **24**(11):987–1003.
- Manuel DG, Kwong K, Tanuseputro P, Lim J, Mustard CA, Anderson GM, Ardal S, Alter DA, Laupacis A: **Effectiveness and efficiency of different guidelines on statin treatment for preventing deaths from coronary heart disease: modelling study.** *BMJ* 2006, **332**(7555):1419.
- McElduff P, Jaefarnezhad M, Durrington PN: **American, British and European recommendations for statins in the primary prevention of cardiovascular disease applied to British men studied prospectively.** *Heart* 2006, **92**(9):1213–1218.
- Strandberg-Larsen M, Nielsen M, Vallgård S, Krasnik A, Vrangbæk K: *In Health system review*. Edited by Mossialos E. Denmark: Health Systems in Transition: World Health Organisation; 2007. **9**(6).
- Donaldson C, Gerard K: *Economics of health care financing: The visible hand*. 1st edition. London: Macmillan Press Ltd; 1993.
- Pedersen KM: **Pricing and reimbursement of drugs in Denmark.** *Eur J Health Econom* 2003, **4**:460–465.
- van DE, Wagstaff A, der BH v, Christiansen T, De GD, Duchesne I, Gerdtham UG, Gerfin M, Geurts J, Gross L, Hakkinen U, John J, Klavus J, Leu RE, Nolan B, O'Donnell O, Propper C, Puffer F, Schellhorn M, Sundberg G, Winkelhake O: **Equity in the delivery of health care in Europe and the US.** *J Health Econ* 2000, **19**(5):553–583.
- van Doorslaer E, Koolman X, Jones AM: **Explaining income-related inequalities in doctor utilisation in Europe.** *Health Econ* 2004, **13**(7):629–647.
- Wagstaff A, van Doorslaer E: **Measuring and testing for inequity in the delivery of health care.** *J Hum Resour* 2000, **35**(4):716–733.
- Rasmussen JN, Rasmussen S, Gislason GH, Abildstrom SZ, Schramm TK, Torp-Pedersen C, Kober L, Diderichsen F, Osler M, Madsen M: **Persistent socio-economic differences in revascularization after acute myocardial infarction despite a universal health care system—a Danish study.** *Cardiovasc Drugs Ther* 2007, **21**(6):449–457.
- Gundgaard J: **Income related inequality in prescription drugs in Denmark.** *Pharmacoepidemiol Drug Saf* 2005, **14**(5):307–317.
- Rasmussen JN, Gislason GH, Rasmussen S, Abildstrom SZ, Schramm TK, Kober L, Diderichsen F, Osler M, Torp-Pedersen C, Madsen M: **Use of statins and beta-blockers after acute myocardial infarction according to income and education.** *J Epidemiol Community Health* 2007, **61**(12):1091–1097.
- Gundgaard J: **Income-related inequality in utilization of health services in Denmark: Evidence from Funen County.** *Scand J Public Health* 2006, **34**(5):462–471.
- Pammolli F, Riccaboni M, Magazzini L: **The sustainability of European health care systems: beyond income and aging.** *Eur J Health Econ* 2011, E-pub ahead of print.
- Chappell NL, Hollander MJ: **An evidence-based policy prescription for an aging population.** *Healthc Pap* 2011, **11**(1):8–18.
- Getz L, Sigurdsson JA, Hetlevik I, Kircheng AL, Romundstad S, Holmen J: **Estimating the high risk group for cardiovascular disease in the Norwegian HUNT 2 population according to the 2003 European guidelines: modelling study.** *BMJ* 2005, **331**(7516):551.
- Ward S, Lloyd JM, Pandor A, Holmes M, Ara R, Ryan A, Yeo W, Payne N: **A systematic review and economic evaluation of statins for the prevention of coronary events.** *Health Technol Assess* 2007, **11**(14):1–160.
- Sigurdsson EL, Thorgeirsson G: **Primary prevention of cardiovascular diseases.** *Scand J Prim Health Care* 2003, **21**(2):68–74.
- Rose G: *The strategy of preventive medicine*. Oxford, New York, Tokyo: Oxford University Press; 1992.
- Emberson J, Whincup P, Morris R, Walker M, Ebrahim S: **Evaluating the impact of population and high-risk strategies for the primary prevention of cardiovascular disease.** *Eur Heart J* 2004, **25**(6):484–491.
- Manuel DG, Lim J, Tanuseputro P, Anderson GM, Alter DA, Laupacis A, Mustard CA: **Revisiting Rose: strategies for reducing coronary heart disease.** *BMJ* 2006, **332**(7542):659–662.
- Marmot M, Friel S, Bell R, Houweling TA, Taylor S: **Closing the gap in a generation: health equity through action on the social determinants of health.** *Lancet* 2008, **372**(9650):1661–1669.
- Atkinson T: **Social inclusion and the European Union.** *Journal of Common Market Studies* 2002, **40**(4):625–643.
- Capewell S, Graham H: **Will cardiovascular disease prevention widen health inequalities?** *PLoS Med* 2010, **7**(8):e1000320.
- Tunstall-Pedoe H, Woodward M: **By neglecting deprivation, cardiovascular risk scoring will exacerbate social gradients in disease.** *Heart* 2006, **92**(3):307–310.
- Usher C, Bennett K, Feely J: **Regional variation in the prescribing for diabetes and use of secondary preventative therapies in Ireland.** *Pharmacoepidemiol Drug Saf* 2005, **14**(8):537–544.
- Ward PR, Noyce PR, St Leger AS: **How equitable are GP practice prescribing rates for statins?: an ecological study in four primary care trusts in North West England.** *Int J Equity Health* 2007, **6**:2.
- Ohlsson H, Lynch K, Merlo J: **Is the physician's adherence to prescription guidelines associated with the patient's socio-economic position? An analysis of statin prescription in South Sweden.** *J Epidemiol Community Health* 2010, **64**(8):678–683.
- Selmer R, Sakshaug S, Skurtveit S, Furu K, Tverdal A: **Statin treatment in a cohort of 20 212 men and women in Norway according to cardiovascular risk factors and level of education.** *Br J Clin Pharmacol* 2009, **67**(3):355–362.

41. Reid FD, Cook DG, Whincup PH: Use of statins in the secondary prevention of coronary heart disease: is treatment equitable? *Heart* 2002, **88**(1):15–19.
42. Shin H, Kim J: Differences in income-related inequality and horizontal inequity in ambulatory care use between rural and non-rural areas: using the 1998–2001 U.S. National Health Interview Survey data. *Int J Equity Health* 2010, **9**:17.
43. Kildemoes HW, Sorensen HT, Hallas J: The Danish National Prescription Registry. *Scand J Public Health* 2011, **39**(Suppl 7):38–41.
44. WHO Collaborating centre for Drug Statistics Methodology: *Guidelines for ATC classification and DDD assignment, 2010*. Oslo: Norwegian Institute of Public Health; 2010.
45. Wallach Kildemoes H, Hendriksen C, Andersen M: Drug utilization according to reason for prescribing: a pharmacoepidemiologic method based on an indication hierarchy. *Pharmacoepidemiol Drug Saf* 2011. doi:10.1002/pds.2195.
46. Mackenbach JP, Kunst AE: Measuring the magnitude of socio-economic inequalities in health: an overview of available measures illustrated with two examples from Europe. *Soc Sci Med* 1997, **44**(6):757–771.
47. Madsen M, Davidsen M, Rasmussen S, Abildstrom SZ, Osler M: The validity of the diagnosis of acute myocardial infarction in routine statistics: a comparison of mortality and hospital discharge data with the Danish MONICA registry. *J Clin Epidemiol* 2003, **56**(2):124–130.
48. Efron B: Nonparametric Estimates of Standard Error - the Jackknife, the Bootstrap and Other Methods. *Biometrika* 1981, **68**(3):589–599.
49. Woodward M, Brindle P, Tunstall-Pedoe H: Adding social deprivation and family history to cardiovascular risk assessment: the ASSIGN score from the Scottish Heart Health Extended Cohort (SHHEC). *Heart* 2007, **93**(2):172–176.
50. Hippisley-Cox J, Coupland C, Vinogradova Y, Robson J, Minhas R, Sheikh A, Brindle P: Predicting cardiovascular risk in England and Wales: prospective derivation and validation of QRISK2. *BMJ* 2008, **336**(7659):1475–1482.
51. Thomsen TF, McGee D, Davidsen M, Jorgensen T: A cross-validation of risk-scores for coronary heart disease mortality based on data from the Glostrup Population Studies and Framingham Heart Study. *Int J Epidemiol* 2002, **31**(4):817–822.
52. Brugts JJ, Deckers JW: Statin prescription in men and women at cardiovascular risk: to whom and when? *Curr Opin Cardiol* 2010, **25**(4):484–489.
53. Henderson R, Keiding N: Individual survival time prediction using statistical models. *J Med Ethics* 2005, **31**(12):703–706.
54. Law MR, Wald NJ, Morris JK: The performance of blood pressure and other cardiovascular risk factors as screening tests for ischaemic heart disease and stroke. *J Med Screen* 2004, **11**(1):3–7.
55. Rose G: Sick individuals and sick populations. *Int J Epidemiol* 1985, **14**(1):32–38.
56. Diederichsen AC, Sand NP, Norgaard B, Lambrechtsen J, Jensen JM, Munkholm H, Aziz A, Gerke O, Egstrup K, Larsen ML, Petersen H, Hoiland-Carlson PF, Mickleth H: Discrepancy between coronary artery calcium score and HeartScore in middle-aged Danes: the DanRisk study. *Eur J Prev Cardiol* 2012, **19**(3):558–564.
57. Smith GD, Hart C, Watt G, Hole D, Hawthorne V: Individual social class, area-based deprivation, cardiovascular disease risk factors, and mortality: the Renfrew and Paisley Study. *J Epidemiol Community Health* 1998, **52**(6):399–405.
58. Suadicani P, Hein HO, von Eyben FE, Gyntelberg F: Metabolic and lifestyle predictors of ischemic heart disease and all-cause mortality among normal weight, overweight, and obese men: a 16-year follow-up in the Copenhagen Male Study. *Metab Syndr Relat Disord* 2009, **7**(2):97–104.
59. Dominguez H, Schramm TK, Norgaard ML, Abildstrom SZ, Kober L, Jorgensen C, Guterbaum TJ, Poulsen HE, Torp-Pedersen C, Gislason GH: Initiation and persistence to statin treatment in patients with diabetes receiving glucose-lowering medications. *Open Cardiovasc Med J* 2009, **3**:152–159.
60. Forde I, Chandola T, Raine R, Marmot MG, Kivimaki M: Socioeconomic and ethnic differences in use of lipid-lowering drugs after deregulation of simvastatin in the UK: the Whitehall II prospective cohort study. *Atherosclerosis* 2011, **215**(1):223–228.
61. Stocks NP, Ryan P, McElroy H, Allan J: Statin prescribing in Australia: socioeconomic and sex differences. A cross-sectional study. *Med J Aust* 2004, **180**(5):229–231.
62. Brookhart MA, Patrick AR, Dormuth C, Avorn J, Shrank W, Cadarette SM, Solomon DH: Adherence to lipid-lowering therapy and the use of preventive health services: an investigation of the healthy user effect. *Am J Epidemiol* 2007, **166**(3):348–354.
63. McCaffery K, Wardle J, Nadel M, Atkin W: Socioeconomic variation in participation in colorectal cancer screening. *J Med Screen* 2002, **9**(3):104–108.
64. von Wagner C, Good A, Whitaker KL, Wardle J: Psychosocial Determinants of Socioeconomic Inequalities in Cancer Screening Participation: A Conceptual Framework. *Epidemiol Rev* 2011, **33**(1):135–147.
65. Capewell S: Will screening individuals at high risk of cardiovascular events deliver large benefits? *No. BMJ* 2008, **337**:a1395.
66. Perreault S, Blais L, Dragomir A, Bouchard MH, Lalonde L, Laurier C, Collin J: Persistence and determinants of statin therapy among middle-aged patients free of cardiovascular disease. *Eur J Clin Pharmacol* 2005, **61**(9):667–674.
67. Golomb BA, Evans MA: Statin adverse effects: a review of the literature and evidence for a mitochondrial mechanism. *Am J Cardiovasc Drugs* 2008, **8**(6):373–418.
68. Bates TR, Connaughton VM, Watts GF: Non-adherence to statin therapy: a major challenge for preventive cardiology. *Expert Opin Pharmacother* 2009, **10**(18):2973–2985.
69. Huser MA, Evans TS, Berger V: Medication adherence trends with statins. *Adv Ther* 2005, **22**(2):163–171.
70. Lindman AS, Veierod MB, Pedersen JI, Tverdal A, Njolstad I, Selmer R: The ability of the SCORE high-risk model to predict 10-year cardiovascular disease mortality in Norway. *Eur J Cardiovasc Prev Rehabil* 2007, **14**(4):501–507.
71. Hingorani AD, Hemingway H: How should we balance individual and population benefits of statins for preventing cardiovascular disease? *BMJ* 2011, **342**:c6244.
72. Marshall T: Targeted case finding for cardiovascular prevention. *BMJ* 2010, **340**:c1376.
73. Wald DS, Wald NJ: The polypill in the primary prevention of cardiovascular disease. *Fundam Clin Pharmacol* 2010, **24**(1):29–35.
74. Malekzadeh F, Marshall T, Pourshams A, Gharravi M, Aslani A, Nateghi A, Rastegarpanah M, Khoshnia M, Semnani S, Salahi R, Thomas GN, Larjani B, Cheng KK, Malekzadeh R: A pilot double-blind randomised placebo-controlled trial of the effects of fixed-dose combination therapy ('polypill') on cardiovascular risk factors. *Int J Clin Pract* 2010, **64**(9):1220–1227.
75. Vos T, Carter R, Barendregt J, Mihalopoulos C, Veerman J, Magnus A, Cobiac L, Bertram M, Wallace A: *Assessing cost-effectiveness in prevention (ACE-Prevention): Final report*. Melbourne: University of Queensland, Brisbane and Deakin University; 2010. www.sph.uq.edu.au/bodce-ace-prevention.
76. Diderichsen F, Andersen I, Manuel C: *Health inequalities - determinants and policies*. Copenhagen: University of Copenhagen & Danish Ministry of Health; 2011. http://www.sst.dk/~media/English/Health%20promotion%20and%20disease%20prevention/International/Health%20inequality%20-%20determinants%20and%20policies.ashx.

doi:10.1186/1471-2458-12-610

Cite this article as: Wallach-Kildemoes et al.: Is the high-risk strategy to prevent cardiovascular disease equitable? A pharmacoepidemiological cohort study. *BMC Public Health* 2012 **12**:610.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

