

Research article

Open Access

Portuguese students' knowledge of antibiotics: a cross-sectional study of secondary school and university students in Braga

Maria Manuel Azevedo¹, Céline Pinheiro², John Yaphe³ and Fátima Baltazar^{*2}

Address: ¹School E.B.2, 3 D. Maria II, Vila Nova de Famalicão Portugal, Portugal, ²Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal and ³Community Health, School of Health Sciences, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

Email: Maria Manuel Azevedo - mariamanuel2001@megamail.pt; Céline Pinheiro - cpinheiro@ecsau.de.uminho.pt; John Yaphe - yonahyaphe@hotmail.com; Fátima Baltazar* - fbaltazar@ecsau.de.uminho.pt

* Corresponding author

Published: 23 September 2009

Received: 16 January 2009

BMC Public Health 2009, 9:359 doi:10.1186/1471-2458-9-359

Accepted: 23 September 2009

This article is available from: <http://www.biomedcentral.com/1471-2458/9/359>

© 2009 Azevedo et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Recent surveys show that the knowledge of the general public about the correct use of antibiotics is limited. This contributes to the problem of inappropriate antibiotic use, leading to a progressive loss of bacterial sensitivity to these drugs and the spreading of resistant strains of bacteria.

Methods: In this cross-sectional study, a questionnaire about antibiotic use was given to a sample of students in the 9th and 12th grades of secondary school and in the first year of university in the north of Portugal.

Results: 349 students returned completed questionnaires. Deficits were found in the students' knowledge of antibiotics and their correct use. Only 4% of 9th grade students were aware that antibiotics are used to treat bacteria only, while 14% of 12th grade students and 29% of first-year university students were aware of this. Fewer students were aware that antibiotics are used to treat tuberculosis. There were deficiencies in the knowledge of timing and duration of therapy. However close to 70% of these students are aware that inappropriate use of antibiotics can contribute to resistance to these drugs.

Conclusion: This study has observed a lack of general knowledge on correct antibiotic use in Portugal, as has been found in other countries. Since this may be due to a lack of formal education on this subject, we believe that a teaching unit on infectious diseases should be included in the 9th and 12th grades, in all curricular areas, with emphasis on bacterial and viral pathogens and correct antibiotic use. In addition, education on the correct use of medications may need to begin at much earlier ages.

Background

Antibiotics are among the most commonly prescribed medications, however they are very often misused [1,2].

Among other factors, the indiscriminate use of antibiotics has contributed to the progressive loss of bacterial sensitivity to antibiotics and spreading of resistant strains of

bacteria [3], with substantial clinical and economic impact [4]. The clinical effectiveness of antibiotics depends partially on their correct use, depending on patients, physicians and retailers [5]. Physicians' decisions may be influenced by several factors such as the fear of losing a patient's trust, the lack of correct information on indications for antibiotic use and pressure from patients and families [2]. Patient factors relating to incorrect antibiotic use include self-medication, sharing medication with other people, not taking a full course of treatment and keeping part of the course for another occasion [2,6].

A study performed in 2001 by Eurobarometer revealed that 60% of Europeans do not know that antibiotics are ineffective against viruses [7]. This lack of accurate information may result in high rates of inappropriate consumption. In 2002, Davey and collaborators [8] reviewed several examples of misperceptions regarding antibiotic use in respiratory tract infections. A study by Pechere in 2001 [9], carried out in different countries with more than 5,000 individuals, reported that more than 60% of those studied believed that antibiotics should be prescribed for viral illnesses. Other surveys showed that most of the patients with acute respiratory symptoms expect to receive antibiotics [10-12]. Another study showed that, in Moldova, many adults are unaware that antibiotics do not cure viral infections and the authors believe that many physicians and pharmacists dispense and prescribe antibiotics without regard to the cause of infection [13].

In Portugal, as most countries, antibiotics are dispensed only by prescription however, studies show that people can still obtain antibiotics without prescriptions [9]. McKee found that 26% of patients studied in his sample of American patients had obtained antibiotics without a physician's prescription and this has been observed in European countries as well [14,15].

Even when antibiotics are available only by prescription, the education of the public on correct use of antibiotics is necessary for the success of the treatment and prevention of the spread of bacterial resistance [16]. Information campaigns on antibiotic resistance carried out in Belgium, led to important reduction in antibiotic consumption [17]. Despite the importance of public information campaigns, the effect of an individual campaign is transient, thus these interventions must be sustained.

Following analysis of the Portuguese curriculum in Natural sciences and Biology, it was noted that very little teaching is done in the field of Microbiology. In the 6th grade, among 11 and 12 year-old students, there is a teaching unit on Microbiology. However, the program is limited to microbial classification, characteristics of microorgan-

isms, pathogenicity and control of microbial growth. In later years there are no other courses on Microbiology.

Given the lack of correct information on antibiotics and its association with antibiotic resistance, an evaluation was planned to test the knowledge of Portuguese students in the 9th grade (14 to 15 years old), 12th grade (16 to 17 years old), and first-year university students in different fields of study, regarding antibiotic spectra, indications and their correct use. Differences in the students' knowledge of this subject by grade level and the area of study were to be assessed.

Methods

This survey was conducted between February and April 2007. A convenience sample was used comprising 349 students including 179 from the 9th and 12th grades and 170 first year university students from the district of Braga, the third largest city in Portugal. The sample of 9th year students comprises 80% of the total number of 9th grade students from D. Maria II School, V. N. de Famalicão. In the 12th year, one class was selected at random from each study area from secondary schools in the same city. The sample of first year university students comprises all students from the selected courses taught by the University of Minho (Braga). Ethical approval for the study was obtained from the directors of the schools involved in the study. Participation in the study was voluntary and anonymous. Participants were informed of this and assured that no participant could be identified from pooled presentation of the results. There were no refusals to participate. The data were obtained through the administration of a seven-item questionnaire [see additional file 1], developed by the authors and pre-tested with a sample of 10 subjects for comprehension of the questions. Questionnaires were completed during regular classes with a time limit of 20 minutes for completion. The questionnaire was designed to assess the student's knowledge on the types of organisms sensitive to antibiotics, the types of infectious diseases treated with antibiotics and the correct use of antibiotics.

Data were analysed using the SPSS statistical software Version 14.0. (SPSS Inc. Chicago, IL, USA). Associations between variables were tested with Pearson's Chi-square (χ^2) with significance set at the $p < 0.05$ level.

Results

The response rate was 100% among the 349 students asked to participate. The characteristics of the study population are shown in Table 1.

Data presented in Table 2 show the knowledge of students on the sensitivity of organisms to antibiotics. There is an increase in the number of correct answers by grade level

Table 1: Characteristics of the study population (n = 349).

Grade level	Age (years)	Number of students
9th grade	14-18	100
12th grade		79
Science	16-17	39
Humanities	16-18	7
Informatics	16-19	19
Arts	16-20	14
1st year of university		170
Nursing	18-20	65
Biology/Geology	18-21	14
Law	18-22	49
Engineering	18-22	42

up to university, however with significant heterogeneity in answers among university study areas. The highest scores were obtained by students in the study areas with closer affinity to the topic assessed, such as sciences (12th grade) and nursing (university).

The students' knowledge of the value of antibiotics in the treatment of common diseases of different aetiologies is presented in Table 3. The number of correct answers increases with the grade level, with marked heterogeneity among study areas in 12th grade. There are significant differences between the answers of students from different study areas (12th grade and university). The highest scores were obtained by students in the sciences (12th grade) and

nursing (university). There are high percentages of incorrect answers among all grade levels evaluated.

The responses to questions on correct antibiotic use are shown in Table 4. Significant differences were found in the proportion of correct responses among the students from the three grade levels assessed, with an increase in the number of correct answers by grade level. The highest scores were obtained by students in the study areas with closer affinity to the topic assessed.

Discussion

This study assessed the knowledge of antibiotics among Portuguese school students of 9th and 12th grade and first year of university. We believe this was the first study of its kind to be performed in Portugal.

A convenience sample of high-school and university students was used to allow for rapid collection of data in a short period of time with limited resources. Information about possible confounders, such as socio-economic status and intellectual level, was not collected. These factors may limit the generalizability of our findings. Repeating this study in other areas of the country with random sampling of students and attention to possible confounders will help answer this question.

Students' knowledge on antibiotic spectra and indications for use were limited at all three school levels evaluated, and misconceptions were prevalent among students, however with a lower error rate for nursing students. The differences observed in scores among 12th grade science

Table 2: Percentage of positive answers to question on antibiotic use against bacteria and other organisms.

Options	n	Antibiotics are effective against bacteria only* (%)	Antibiotics are effective against bacteria and other organisms (%)	Antibiotics are effective against other organisms (%)	P value
9th grade	100	4	65	31	
12th grade					
Sciences	39	28	46	26	
Humanities	7	14	14	72	
Informatics	19	0	74	26	
Arts	14	14	65	21	
Mean ± SD		14 ± 11.4	49.8 ± 26.5	36.3 ± 24	0.066
1st year of university					
Biology/Geology	14	29	57	14	
Nursing	65	60	34	6	
Law	49	16	57	27	
Engineering	42	14	62	24	
Mean ± SD		29.8 ± 21.2	52.5 ± 12.6	17.8 ± 9.6	<0.001

SD-Standard deviation, *- Correct answer

Table 3: Percentage of positive answers to questions on antibiotic use against viral illness (influenza, hepatitis, AIDS) and tuberculosis.

Options	n	Antibiotics should be prescribed for tuberculosis* (%)	Antibiotics should be prescribed for tuberculosis and viral illnesses (%)	Antibiotics should be prescribed for viral illnesses (%)	P value
9th grade	100	2	3	95	
12th grade					
Sciences	39	18	8	74	
Humanities	7	0	14	86	
Informatics	19	0	38	62	
Arts	14	0	28	72	
Mean ± SD		4.5 ± 9.0	22 ± 13.6	73.5 ± 9.9	0.049
1st year of university					
Biology/Geology	14	14	43	43	
Nursing	65	38	30	32	
Law	49	4	19	77	
Engineering	42	13	19	68	
Mean ± SD		17.3 ± 14.6	27.8 ± 11.4	55 ± 21.0	0.001

SD-Standard deviation, *- Correct answer

students, nursing students and the remaining students, could be explained by either acquisition of knowledge from different sources other than school, or the result of selection of students, given the higher marks required for admission to science higher degrees.

With respect to correct antibiotic use, our results showed limited knowledge among 9th and 12th grade students and nursing students obtained the best results among university students. This issue is of concern because many students complete their schooling after 9 years of

compulsory education and may receive no further instruction on this topic. This may also influence subsequent generations. Recent studies showed that mothers often influence medical decisions on antibiotic prescription [13]. In this context, paediatricians are often pressured to prescribe antibiotics to children with viral infectious [18,19]. Cebotarenco [20] showed that in times of high incidence of viral infection, half of these infectious diseases are treated with antibiotics, self prescription. In addition, mothers may administer these drugs to children without medical advice [13]. This suggests the need for

Table 4: Percentage of correct answers to questions on antibiotic treatment for bacterial infections.

Study question	n	Correct answer	Grade 9 (%)	Grade 12 (%)	1 st year university (%)	Overall p
			100	79	170	
			Mean ± SD ^a	Mean ± SD ^b		
Antibiotics should be taken with milk	F	57	51.8 ± 13.5	59.2 ± 10.1	0.537	
Antibiotics do not interact with alcohol	F	46	65.5 ± 6.6	75.0 ± 11.3	<0.001	
Antibiotics can be taken at different times each day, if the daily doses are taken.	F	50	49.5 ± 21.9	76.0 ± 9.9	<0.001	
Antibiotic treatment should be stopped as soon as the patient feels better	F	57	66.0 ± 20.4	77.7 ± 8.8	<0.001	
Remaining antibiotic doses can be saved for use on other occasions	F	59	57.5 ± 11.1	79.2 ± 10.7	<0.001	
The incorrect use of antibiotics can lead to development of resistant bacteria.	T	70	68.0 ± 17.8	77.2 ± 5.3	0.570	

SD-Standard deviation; T = true; F = false.

^a, mean score from 4 study areas: sciences, humanities, informatics and arts

^b, mean score from 4 study areas: biology/geology, nursing, law and engineering

more widespread education on the proper use of antibiotics. An extensive school-based educational program in Moldova, which included peer-education sessions, parents' meetings, and distribution of educational newsletters, was successful in reducing antibiotic use for treatment of presumed viral respiratory illnesses [13]. Bush has suggested that these efforts begin as early as kindergarten based on observations of successful programs involving young children and their parents [21]. The United States Pharmacopeia Ad Hoc Advisory Committee on Children and Medicines has guidelines for preparation of educational materials about medication for children as young as three years old. As outlined in the USP Guiding Principles for teaching children and adolescents about medication, education needs to be tailored to their development, capabilities and experience [22].

In the present study, there was a trend towards an increase in the number of correct answers with the grade level. This is in accordance with the study of You and co-workers [23], who suggested that higher education is a positive predictor for adequate knowledge and appropriate attitudes to antibiotic use. Increasing prevalence of antibiotic resistance by bacteria, partly due to indiscriminate widespread use of antibiotics, is a threat to public health. Increasing public awareness of the problem and education of the general public and retailers on proper usage of antibiotics may help to slow this trend [24,25]. Patient demand for antimicrobials might be triggered by mass media, such as TV, internet, magazine or newspaper advertising, behaviours which also contribute to the development of resistance. In a study undertaken in Europe in 1997, physicians elected patients' pressure as the main reason for prescribing inadequate antibiotics [2]. These findings highlight the need to educate people about the appropriate use of antibiotics.

Conclusion

There were marked deficiencies found in the knowledge of Portuguese students of antibiotics and their correct use. This may be due to a lack of formal education on the subject in schools. Education about the correct use of medication may need to begin at very early ages. We believe that a teaching unit on Microbiology should be included in 9th grade curriculum, with emphasis on knowledge of antibiotic spectra/indications and correct antibiotic use. This unit should be reinforced in the 12th grade in all curricular areas. We also believe that it is important to design teaching programs to be tested in schools to improve the knowledge of students on this subject. We suggest that this approach may modify behaviours with regard to antibiotic use with benefits for public health.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MMA and FB designed the study and wrote the manuscript. CP performed data analysis and helped write the manuscript. JY helped to write and edit the manuscript. All authors have read and approved the final version of the manuscript.

Additional material

Additional file 1

Questionnaire. This file contains the questionnaire given to the students concerning antibiotic use.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2458-9-359-S1.doc>]

Acknowledgements

We want to thank the students for their active participation in this study and to thank their teachers for encouraging them to participate. Thanks are due to Pedro Oliveira for statistical advice.

References

1. Chambers HF: **General principles of antimicrobial therapy.** In *Goodman & Gilman's The Pharmacological Basis of Therapeutics* 11th edition. Edited by: Brunton L, Parker K, Blumenthal D, Buxton I. New York, USA, Mc Graw Hill; 2006:1095-110.
2. World Health Organization: **Report on Infectious Diseases. Overcoming antimicrobial resistance.** 2000 [<http://www.who.int/infectious-disease-report/2000/index.html>]. Geneva: World Health Organization
3. Centers for Disease Control and Prevention (CDC) [<http://www.cdc.gov/Features/GetSmart/>]
4. Maragakis LL, Perencevich EN, Cosgrove SE: **Clinical and economic burden of antimicrobial resistance.** *Expert Rev Anti Infect Ther* 2008, **6**:751-63.
5. Radyowijati A, Haak H: **Determinants of antimicrobial use in the developing world.** In *Child Health Special Report Volume 4.* Issue 1 Washington, DC: USAID Bureau of Global Health; 2002.
6. Grigoryan L, Burgerhof JG, Haaijer-Ruskamp FM, Degener JE, Deschepper R, Monnet DL, Di Matteo A, Scicluna EA, Bara AC, Lundborg CS, Birkin J, SAR group: **Is self-medication with antibiotics in Europe driven by prescribed use?** *J Antimicrob Chemother* 2007, **59**:152-6.
7. Eurobarometer 2001 [<http://ec.europa.eu/research/press/2001/pr0612en-report.pdf>].
8. Davey P, Pagliari C, Hayes A: **The patient's role in the spread and control of bacterial resistance to antibiotics.** *European Society of Clinical Microbiology and Infectious Diseases* 2002, **CMI 8(suppl 2)**:43-68.
9. Pechere JC: **Patient's interviews and misuse of antibiotics.** *Clin Infect Dis* 2001, **33(Suppl 3)**:S170-3.
10. Haltinger KA, Hayden GF, Weber T, Evans BA, Possner AB: **Antibiotic-seeking behaviour in college students: what do they really expect?** *J Am Coll Health* 2001, **50**:9-13.
11. Mangione-Smith R, McGlynn EA, Elliot MN, McDonald L, Franz CE, Kravitz RL: **Parent expectations for antibiotics, physician-parent communication, and satisfaction.** *Arch Pediatr Adolesc Med* 2001, **155**:800-6.
12. Cals JW, Boumans D, Lardinois RJM, Gonzales R, Hopstaken RM, Butler CC, Dinant G-J: **Public beliefs on antibiotics and respiratory tract infections: an internet-based questionnaire study.** *Br J Gen Pract* 2007, **57**:942-47.
13. Cebotarenco N, Bush PJ: **Reducing Antibiotics for Colds and Flu: A Student Taught Program.** *Health Educ Res* 2008, **23**:146-157.

14. McKee MD, Mills L, Mainous AGIII: **Antibiotic use for the treatment of upper respiratory infections in a diverse community.** *J Fam Pract* 1999, **48**:993-6.
15. Berzanskyte A, Valinteliene R, Haaijer-Ruskamp FM, Gurevicius R, Grigoryan L: **Self-medication with antibiotics in Lithuania.** *Int J Occup Med Environ Health* 2006, **19**:246-53.
16. Grigoryan L, Burgerhof JG, Degener JE, Deschepper R, Lundborg CS, Monnet DL, Scicluna EA, Birkin J, Haaijer-Ruskamp FM: **Self-Medication with Antibiotics and Resistance (SAR) Consortium. Determinants of self-medication with antibiotics in Europe: the impact of beliefs, country wealth and the healthcare system.** *J Antimicrob Chemother* 2008, **61**:1172-9.
17. Bauraind I, Lopez-Lozano JM, Beyaert A, Marchal JL, Seys B, Yane F, Hendrickx E, Goossens H, Tulkens PM, Verbist L: **Association Between Antibiotic Sales and Public Campaigns for Their Appropriate Use.** *JAMA* 2004, **292**:2468-70.
18. Mangione-Smith R, McGlynn EA, Elliott MN: **The relationship between perceived parental expectations and paediatrician antimicrobial prescribing behaviour.** *Pediatrics* 1999, **103**:711-8.
19. Palmer DA, Bauchner H: **Parents' and Physicians' views on antibiotics.** *Pediatrics* 1997, **99**(6):E6.
20. Cebotarenco N: **Children's knowledge about antibiotics.** *Buletinul Institutului National De Farmacie* 1999, **3**:2-14.
21. Bush PJ: **Can kindergartners help decrease inappropriate antibiotic use?** *J Sch Health* 2007, **77**:650.
22. Bush PJ, Ozias JM, Walson PD, Ward RM: **Ten guiding principles for teaching children and adolescents about medicines.** *Clin Therap* 1999, **21**:1280-1284.
23. You JHS, Yau B, Choi KC, Chau CTS, Huang OR, Lee SS: **Public Knowledge, Attitudes and Behavior on Antibiotic Use: A Telephone Survey in Hong Kong.** *Infection* 2008, **36**:153-7.
24. Wenzel RP, Edmond MB: **Managing antibiotic resistance.** *N Engl J Med* 2000, **343**:1961-3.
25. Wachter DA, Joshi MP, Rimal B: **Antibiotic dispensing by drug retailers in Kathmandu, Nepal.** *Trop Med Int Health* 1998, **4**:782-8.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/9/359/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

