

ERRATUM

Open Access



Erratum to: The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents

Christiane Stock*, Lotte Vallentin-Holbech and Birthe Marie Rasmussen

Erratum

Following publication of this study design article [1], it has come to our attention that some terminology was inconsistently used in the article. The GOOD life trial uses a definition of binge drinking of “5 or more drinks on one occasion” in accordance with the ESPAD study [2]. Both in the abstract and in the methods section, when describing measures of alcohol use or outcomes, any formulation of “more than 5 drinks” becomes “5 or more drinks”. Also, when respondents were asked to identify the amount of occasions they have been drinking 5 or more drinks in the past 30 days, the binge drinking definition was not according to the MULDD study [3], but according to the ESPAD study [2].

Received: 15 May 2017 Accepted: 17 May 2017
Published online: 24 May 2017

References

1. Stock C, Vallentin-Holbech L, Rasmussen BM. The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents. *BMC Public Health*. 2016;16:704.
2. Hibell B, Guttormsson U, Ahlström S, Balakireva O, Bjarnason T, Kokkevi A, Kraus L. The 2011 ESPAD report-substance use among students in 36 European countries. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN); 2012.
3. Nielsen GA, Ringgaard L, Broholm K, Sindballe A, Olsen SF. Unge's livsstil og dagligdag 2000: forbrug af tobak, alkohol og stoffer [Young people's lifestyle and daily life: use of tobacco, alcohol and drugs]. Copenhagen: The Danish Cancer Society and National Board of Health; 2002.

* Correspondence: cstock@health.sdu.dk
Unit for Health Promotion Research, Department of Public Health, University of Southern Denmark, Esbjerg, Denmark

