

RESEARCH

Open Access



Prevalence and distribution of acute gastrointestinal illness in the community of China: a population-based face-to-face survey, 2014–2015

Jikai Liu¹, Baozhang Luo^{2†}, Yijing Zhou^{3†}, Xiaochen Ma^{4†}, Junhua Liang^{5†}, Xianglai Sang^{6†}, Le Lyu^{7†}, Wen Chen^{8†}, Pengyu Fu^{9†}, Hong Liu², Shiqi Zhen³, Chao Wang⁴, Yangbo Wu⁴, Qiong Huang⁵, Xiaocheng Liang⁶, Guangda Bai⁷, Zhen Lan⁸, Shufang Zhang⁹, Yongning Wu¹, Ning Li^{1*} and Yunchang Guo^{1*}

Abstract

Background The true incidence of acute gastrointestinal illness in China is underrecognized by surveillance systems. The aims of this study were to estimate the incidence and prevalence of self-reported AGI in the community of China, and to investigate sociodemographic and epidemiological determinants of AGI.

Methods We conducted a 12-months cross-sectional population-based survey in eight provinces of China during 2014–2015. The survey determined the prevalence and incidence of acute gastrointestinal illness (AGI) in the total permanent resident population in China according to the census of the population in 2010. The random multilevel population sample was stratified by geographic, population, and socioeconomic status. We used a recommended case definition of AGI, with diarrhea (three loose or watery stools) and/or any vomiting in a four-week recall. A face-to-face survey was conducted by selecting the member in the household with the most recent birthday.

Results Among 56,704 sampled individuals, 948 (1,134 person-time) fulfilled the case definition; 98.5% reported diarrhea. This corresponds to 2.3% (95% CI:1.9%–2.8%) of an overall standardized four-week prevalence and 0.3 (95% CI: 0.23–0.34) episodes per person-year of annual adjusted incidence rate. There was no significant difference between males and females. The incidence rates were higher among urban residents, and in the spring and summer. In the whole study period, 50% of the cases sought medical care, of which 3.9% were hospitalized and 14.3% provided a biological sample for laboratory identification of the causative agent. Children aged 0–4 and young adults aged 15–24, people living in rural areas and people who traveled frequently had higher prevalence of AGI.

Conclusion Results showed that AGI represents a substantial burden in China, and will contribute to the estimation of the global burden of AGI. Complemented with data on the etiologies of AGI, these estimates will form the basis to estimate the burden of foodborne diseases in China.

[†]Baozhang Luo, Yijing Zhou, Xiaochen Ma, Junhua Liang, Xianglai Sang, Le Lyu, Wen Chen and Pengyu Fu contributed equally to this work.

*Correspondence:

Ning Li
lining@cfsa.net.cn
Yunchang Guo
gych@cfsa.net.cn

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords Gastrointestinal disease, Incidence, Care seeking, Community, Diarrhoea, Population survey, China

Introduction

Acute gastrointestinal infections (AGI), including foodborne diseases, are common and cause substantial public health and socioeconomic impact globally. The World Health Organization (WHO) estimated that, in 2010, 22 foodborne enteric agents caused two billion illnesses, over one million deaths, and 78.7 million DALYs [1]. Although enteric disease manifests primarily as self-limiting AGI, appearing as diarrhea or vomiting, more severe disease and even death can occur [2].

Knowledge of the magnitude, distribution, and demographic factors associated with AGI is a requirement for estimating the burden of acute gastroenteritis of foodborne origin, the burden caused by specific pathogens commonly transmitted through foods, and the burden caused by specific foods or food groups [3]. Such evidence is key to define strategies for disease prevention. However, data obtained from public health surveillance does not reflect the true burden of AGI, reflecting any failure in the process between a patient becoming ill and the case being reported: patients often do not seek medical care, samples and laboratory analyses may not be performed, and results may go unregistered [4–6]. In China, a pilot study performed in six southern provinces between 2010 to 2011 estimated the incidence of AGI [7]. However, it did not include northern and western provinces, and thus was not representative of the whole China. Inspired by this survey, some provinces estimated their own provincial data in the survey or conducted the similar national survey in the next year follow the China CDC's instruction [8, 9].

The main aims of this study were to estimate the incidence of self-reported AGI in China using a face-to-face method, and to investigate sociodemographic and epidemiological factors as determinants of AGI. Furthermore, we also assessed clinical manifestations and utilization of medical services in cases of AGI. This third nationwide population-based study on AGI covered eight provinces in the period between 2014 and 2015. The estimated AGI incidence also provides an important parameter for estimation the burden of foodborne diseases in China.

Methods

Study design and population

We conducted a nationwide population-based retrospective face-to-face survey, implemented over 12 months. The population under study comprised persons resident in the survey site for more than six months; infants born

within six months were included. The targeted sample size was calculated according to the residential population of selected provinces, expected incidence, and relative risk (allowed error) with 95% confidence intervals and 20% of the loss rate of follow-up using EPI INFO 3.5.1.

We selected the respondents using a multistage stratified random selection procedure. China has five tiers of local government hierarchy above the household: province, prefecture, county, township, and village. In the first sampling stage, eight provinces or municipalities in China were selected according to geographic, population and socioeconomic status, and food preference by a stratified random sampling method. These province were: Jilin province, Northeast China; Beijing, North China; Shanghai and Jiangsu province, East China; Guangdong province, South China; Henan province, Central China; Gansu province; Northwest China; and Sichuan province; Southwest China (Fig. 1). Simple random sampling (SRS) was used to select three prefecture-level cities or city-level areas within each province in the second stage. In the third stage, two county-level units, including one district and one county-level city, were selected from each prefecture-level city in order to maintain a reasonable ratio between urban and rural areas. SRS selected five villages or towns from each district or county in the fourth stage. In the fifth stage, SRS selected two village-level units from each township unit. In the sixth stage, for each village-level unit, the residential community or village




Fig. 1 Distribution of survey sites of the population-based retrospective face-to-face acute gastrointestinal disease survey, corresponding to provinces or municipalities in mainland China selected stratified random sampling method

administration supplied the list of numbers for each household. The local community or village general practitioners (GPs) are the interviewers who used SRS selected eight households each month from the list; households that were previously selected and interviewed would not be selected again. GPs selected 96 households per village or town for twelve months by SRS.

Before the survey, all the involved provincial and cities' CDC experts took a 3-days training to explain the methods and content of the questionnaire. City level CDC experts gave the GPs in the communities in the districts which they selected a similar training but in local dialects and customs.

In the selected communities or villages, GPs acquainted with the residents entirely and have their contact information which is necessary during the daily routine work. Before interview, To allow for the maximum number of people to be contacted, they contacted the household by phone, messaging app or face-to-face to book the time to meet when most of the residents in the household. Contact of the selected households was attempted up to three times. The GPs made an appointment with the members of the household during off-duty hours if some of them were working in the daytime. The household was calculated as a loss to follow-up if the household remained uncontacted up to three times or refused to be interviewed without a neighboring replacement. Then the GPs used standard and unified printed empty questionnaires visited household. The interviewer determined if the household contained two or more adults, and then asked the adult household member who had the most recent birthday [10, 11] in a face-to-face way. If the household was successful interviewed, the rest of the members wouldn't be asked, and household was interviewed only once. Study participants were enrolled from April 2014 to March 2015. All the interviews were conducted in Chinese with local accent, including Cantonese and other dialects to make understandable by the interviewers and the questions were asked in a colloquial but in line with the correct meaning of those AGI symptoms.

The representativeness of the sample was improved by weighing. These weights were generated to avoid deviation of the target population to the whole Chinese population based on data from 2010 (6th) National population census in China [12].

The questionnaire collected self-reported socio-demographic information on the respondent (gender, age, education, occupation of the main earner, family income and jurisdiction), recent travelling history, information on the etiology of infection, AGI symptoms, related health care-seeking behaviors, treatment (available verification like prescriptions, medication boxes or pill strips are suggested), eventual absence

from work associated with illness and direct/indirect expenses.

Case definition

To ensure that the case definition was simple and applicable, as well as comparability with the results from other surveys conducted globally, we used the standard case definition advocated by the international collaboration on enteric disease 'Burden of Illness' Studies [13]. AGI was defined as *having had at least three episodes of loose stools or any vomiting in any 24-h period*. Participants who reported having fewer than three loose stools in a 24-h period were not counted as a case. The same participants who reported another AGI after 7 days with in 28 days before the interview counted as having the AGI twice. Any AGI symptoms caused by a non-infectious gastrointestinal illness like chronic gastrointestinal diseases (Crohn's disease, ulcerative colitis, stomach cancer, and intestinal tumors, irritable bowel or coeliac disease), or alcohol or drugs, or normal physiological phenomena like pregnancy and menstruation were also excluded.

Data analysis

EpiData version 3.1 (EpiData Association, Odense M, Denmark) was used for data collection. All data were weighted by sampling frame and adjusted by age, gender and residency based on the sixth national population census data in 2010. The analysis was performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA).

Four-week period prevalence (14 wk.) and incidence proportion (expressed in %) were calculated, along with 95% confidence intervals (CIs). The annual incidence (I annual) was calculated as $I \text{ annual} = 14 \text{ wk.} \times (365/28)$ and expressed in terms of episodes/person per year [13]. Qualitative data were described using percentages or proportions, and quantitative data were statistically described using mean and 95% confidence intervals. Values of $P < 0.05$ were considered statistically significant. The aim of calculation the four-week period prevalence is to estimate the annual episodes of the AGI. The null hypothesis of no association between prevalence of disease and age group, cultural group, total per capita household income level, and highest education level was tested using χ^2 test. We measured the association between disease determinants as explanatory variables and the defined AGI cases as the outcome variable, defined as odds ratios (ORs), using multivariable logistic regression. For comparison of average means between age groups and gender, the two-tailed P -value from linear regression was used.

Ethical approval

Ethical approvals were granted by China National Center for Food Safety Risk Assessment’s Ethical Board. All data collected including the names, addresses, and telephone numbers of the participant were kept confidential. Each participant was asked to read and sign a consent form that informed about the purpose of the survey before the questionnaire was administered.

Results

Sample and response rates

In total, 58,320 interviews were completed in the period between April 2014 to March 2015. Among these, 56,704 (97%) valid responses were obtained.

AGI occurrence

At least one episode of acute gastroenteritis was reported by 2.3% (95% CI: 1.9–2.8%) of respondents in the four weeks before the interview, corresponding to a rate of 0.3 (95% CI: 0.23–0.3) episodes per person per year. The prevalence was higher in summer months (Fig. 2). The mean duration of illness (the AGI related symptoms) was 11.3 h (95% CI: 3.5–14.5). Half of the respondents sought medical care (49.8%; 95% CI: 43.9–55.6). Nearly half (47.0%; 95% CI: 42.4–51.6%) of the patients who sought medical care were taking some type of medication because of AGI; 17.6% (96% CI: 14.4–21.4%) reported taking anti-diarrheal medication.

Although there were some differences in the prevalence of AGI in some age groups, particularly in the population 15–24 years of age (Fig. 3), results showed there were no significant differences in the incidence of AGI between

males and females. The incidence was higher among children below 5 years (Fig. 3). Results of multivariate analysis showed that occupation of the main earner, and traveling history were significantly associated with acute gastroenteritis (Table 1).

For 17.4% (95% CI: 14.2–21.2%) of those with AGI, they or a member of their family took time off work due to the diarrhea episode. The median number of days off was two, and the mean was 2.7 days.

Socio-demographic characteristics of AGI

Estimates of the four-week period and annual incidence of AGI in China with specific demographic characteristics are presented in Table 1. There was no significant difference in the incidence of AGI between males and females (0.26 vs 0.31). Except for the traits of age groups mentioned above, the overall incidence of AGI was high in Han nationality than in other nationality (0.29 vs 0.08, $P=0.537$). The prevalence of AGI in the north of China was slightly lower than the rate in the south (0.23 vs 0.21, $P=0.097$). There was no significant difference in the incidence of AGI between Urban and Rural residence and per capita household income. The tendency of incidence seemed highest in preschool children and declined with the increase of educational levels; however differences were not significant. There was a borderline significant difference in incidence of AGI between participants with blue collar jobs and with white collar jobs (0.28 vs 0.45 $p=0.053$). The incidence of AGI was significantly higher in participants who had travelled within two weeks than in those who had not (0.89 vs 0.27). Respondents living in a household with two persons were significantly less




Fig. 2 Prevalence of acute gastroenteritis in the overall population by month, 2014–2015




Fig. 3 Prevalence of acute gastroenteritis in mainland China, by age and gender

likely to have experienced AGI than individuals living alone. The prevalence of AGI varied by provinces, with the highest in Gansu (0.41, $P < 0.001$), and the lowest in Beijing (0.10, $P < 0.001$).

Symptoms, healthcare, and impact of AGI

Among the 1,134 cases, 1,122 (98.9%) reported suffering from diarrhea, 238 (21%) reported suffering from vomiting and 226 (19.9%) cases experienced both symptoms. Of the 1,122 cases with diarrhea, 16 (1.4%) had bloody diarrhea. Respiratory symptoms were reported by 10.4% of all the cases, usually not associated with intestinal disease. The information on the duration of illness, presence of concurrent symptoms and recourse to healthcare is reported in Table 2. The average duration of illness was 46.4 h. Children in the 0–4 year group had a significantly higher fever rate (39.5%) than other age groups ($p < 0.001$). On the worst day of symptoms, cases reported diarrhea an average of 4.2 times (range 3–16 times) and vomiting an average of 2.4 times (range 1–11 times).

The healthcare-seeking behaviors of AGI cases, medication had been taken because of the AGI by cases, and the source of the medicine are reported in Table 3. 36.7% of all the cases took outpatient services, while 4% were hospitalized. Outpatient services-seeking behaviors varied by age; the percentage of service-taking for AGI was highest (74.4%) in children aged <5 years. 11.9% of all the cases reported providing a stool sample for laboratory testing. In total, 78.7% of all the cases took medication for AGI. Nearly half (44.8%) of all cases have taken

antibiotics, while 5.1% reported taking pain killers and 2.8% antipyretics. Only 42.7% reported that the medication was prescribed by a doctor.

Self-reported suspect causes of AGI

Among the positive individuals who reported the cause of AGI, 59.3% reported that their AGI could be food related. Of the food related individuals, 17.5% thought it was due to meat and its products (pig meat, beef and mutton), 13.3% thought it was due to fruit and its products, 12.9% thought it was due to vegetable and its products. The distribution of other food categories was scattered. Of the individuals who could recall the food source, 38.7% from home, 12.9% from fast-food restaurant or snack bar, 11.7% from street stalls.

Discussion

We estimated the magnitude and distribution of AGI in the community in China using a large-scale population-based study. Our results showed that the prevalence of AGI in the population was 2.3%, corresponding to an incidence of 0.3 episodes per person per year. In other words, our study suggests that one in every three Chinese residents suffers from AGI every year. This was the first survey provide nationwide representative estimates of the prevalence and incidence of AGI in China.

Applying the synchronized interviewing cycle and the same questionnaire, this survey covered not only north and south representative provinces in China, but also provinces representing the northwest (Gansu

Table 1 Incidence estimates and determinants for acute gastrointestinal illness (AGI) in China, 2014–2015 (weighted) ($n=56,704$). Statistically significant values are highlighted in bold

Determinants	Sample size	No. of AGI cases (unweighted)	four-week period prevalences	Annual incidence ^b	95% CI ^a	OR	95% CI	P value
Total	56,704	1,134	2.16	0.28	(0.23–0.34)	-	-	-
Gender								
Male ($n=28,047$)	28,047	547	1.96	0.26	(0.19–0.32)	1.07	0.94–1.22	0.298
Female ($n=28,657$)	28,657	587	2.36	0.31	(0.22–0.40)	Ref	Ref	Ref
Age								
< 5	1,493	44	2.93	0.38	(0.22,0.55)	0.61	(0.44,0.85)	0.003
5–14	2,777	55	1.46	0.19	(0.10,0.30)	1.03	(0.76,1.41)	0.844
15–24	3,431	66	2.49	0.33	(0.17,0.50)	1.06	(0.79,1.41)	0.713
25–44	14,870	318	2.34	0.3	(0.21,0.40)	Ref	Ref	Ref
45–64	22,323	426	1.8	0.23	(0.17,0.30)	1.15	(0.98,1.35)	0.085
≥ 65	11,810	225	2	0.26	(0.17,0.36)	1.12	(0.93,1.35)	0.233
Ethnicity								
Han	55,210	1098	2.22	0.29	(0.23,0.35)	1.127	(0.77,1.65)	0.537
Other	1,494	36	0.58	0.08	(0.01,0.15)	Ref	Ref	Ref
Geographical								
North	28,075	555	1.8	0.23	(0.15,0.32)	1.12	(0.98–1.27)	0.097
South	28,629	579	2.4	0.31	(0.24,0.38)	Ref	Ref	Ref
Residence								
Urban	26,178	505	2.65	0.23	(0.15,0.32)	1	(0.88, 1.14)	0.982
Rural	30,526	629	1.81	0.31	(0.24,0.38)	Ref	Ref	Ref
Education								
Preschool children	1900	49	2.41	0.31	(0.18,0.45)	0.72	(0.52,1.00)	0.052
Illiterate	4065	96	2.24	0.29	(0.12,0.48)	0.95	(0.72,1.25)	0.696
Primary school	13,301	286	1.89	0.25	(0.17,0.33)	1.01	(0.82,1.24)	0.913
Middle school	18,388	340	1.74	0.23	(0.14,0.33)	1.22	(1.00,1.49)	0.055
High school	10,579	196	2.25	0.29	(0.16,0.43)	1.16	(0.93,1.45)	0.19
Undergraduate and above	8,471	167	3.14	0.41	(0.27,0.55)	Ref	Ref	Ref
Occupation								
White collar job	7,402	175	3.49	0.45	(0.27,0.66)	Ref	Ref	Ref
Blue collar job	17,624	377	2.14	0.28	(0.17,0.38)	0.82	(0.68,1.00)	0.053
Others	31,678	582	1.85	0.24	(0.19,0.30)	1.08	(0.94,1.25)	0.281
Per capita household income								
< 20,000 RMB/person	28,822	617	1.82	0.24	(0.17,0.31)	Ref	Ref	Ref
20,000–50,000RMB/person	17,347	351	3.01	0.39	(0.27,0.53)	1.01	(0.88,1.17)	0.864
> 50,000 RMB/person	5,270	92	2.16	0.28	(0.14,0.43)	1.17	(0.92,1.48)	0.205
Non-reported	5,265	74	1.95	0.25	(0.12,0.38)	1.38	(1.07,1.78)	0.015
No. of persons in household								
1	7,382	122	2.53	0.33	(0.17,0.53)	Ref	Ref	Ref
2	19,689	426	2.75	0.36	(0.24,0.48)	0.8	(0.64,1.00)	0.046
3	16,188	317	1.93	0.25	(0.16,0.35)	0.88	(0.70,1.10)	0.27
≥ 4	13,445	269	1.94	0.25	(0.17,0.34)	0.88	(0.70,1.12)	0.3
Travel within two weeks								
Yes	1,001	36	6.81	0.89	(0.39,1.46)	Ref	Ref	Ref
No	55,701	1092	2.08	0.27	(0.22,0.33)	0.41	(0.30,0.50)	< 0.001
Non-reported	1	-	-	-	-	-	-	-
Province								
Beijing	9,885	131	0.73	0.1	(0.05,0.15)	Ref	Ref	Ref

Table 1 (continued)

Determinants	Sample size	No. of AGI cases (unweighted)	four-week period prevalences	Annual incidence ^b	95% CI ^a	OR	95% CI	P value
Jilin	5,906	167	3.04	0.4	(0.24,0.56)	0.48	(0.37, 0.62)	<0.001
Shanghai	8,407	149	1.62	0.21	(0.14,0.28)	0.73	(0.57, 0.95)	0.018
Jiangsu	6,948	187	3.58	0.47	(0.31,0.63)	0.47	(0.37, 0.59)	<0.001
Henan	5,433	63	1.14	0.15	(0.07,0.22)	1.06	(0.77, 1.46)	0.717
Guangdong	5,745	108	1.99	0.26	(0.16,0.36)	0.66	(0.50, 0.86)	0.003
Sichuan	7,529	135	2.25	0.29	(0.20,0.39)	0.72	(0.55, 0.94)	0.014
Gansu	6,851	194	3.15	0.41	(0.21,0.64)	0.54	(0.42, 0.69)	<0.001

^a CI Confidence interval^b Annual incidence: specified as episodes per person-year**Table 2** Self-reported proportions and average means for associated factors, care seeking and treatment of acute gastrointestinal illness cases in China, by age and gender ($n = 948$)

	Age group (years)							P value for age	Male	Female	P value for gender
	Total	0–4	5–14	15–24	25–44	45–64	65–				
Diarrhea (%)	98.94	97.67	97.87	98.28	98.87	99.42	98.94	0.283	99.11	98.79	0.624
Vomiting (≥ 1 times/day) (%)	20.95	39.53	36.17	22.41	22.26	17.44	17.02	<0.001	18.4	23.28	0.066
Bloody diarrhea (%)	1.39	2.38	0	0	2.29	0.58	2.15	0.665	0.89	1.84	0.225
Fever (> 38 °C) (%)	11.85	39.53	29.79	6.9	10.94	9.88	7.45	<0.001	11.97	11.74	0.912
Respiratory symptoms (%)	10.36	27.91	12.5	6.9	8.3	10.17	10.11	0.173	11.5	9.31	0.27
Stool sample (%)	11.88	15.63	3.57	20	8.65	11.11	17.74	0.408	8.7	14.67	0.09
Outpatients (%)	36.65	74.42	59.57	34.48	39.62	28.86	32.98	<0.001	35.7	37.53	0.561
Hospitalized (%)	4.02	0	0	5	4.76	4	6.35	0.059	1.84	5.95	0.066
Work absenteeism (%)	13.21	9.3	14.89	15.52	20.08	11.5	6.38	0.001	12.22	14.11	0.394
Medication taken (%)	78.69	95.35	85.11	72.41	75.85	76.61	82.98	0.838	78.44	78.9	0.863
Medication prescribed (%)	42.72	68.29	57.5	42.86	47.76	35.5	37.82	<0.001	42.78	42.67	0.978
Diarrhea medication (%)	45.6	69.23	35.56	34.48	45.95	43.15	50.54	0.557	46.22	45.04	0.719
Antibiotics medication (%)	44.75	25.64	66.67	44.83	46.54	44.35	41.62	0.315	44.65	44.83	0.954
Painkiller medication (%)	5.12	5.13	8.7	8.62	6.59	2.69	5.46	0.089	5.52	4.76	0.604
Antipyretic medication (%)	2.84	15.79	6.52	5.17	3.49	1.2	0.55	<0.001	2.53	3.11	0.6
Mean duration of using medication (hours)	46.43	49.95	35.92	31.1	45.63	42.47	n.a	0.497	43.56	49.03	0.061
Mean duration of symptoms (hours)	42.19	50.11	37.52	35.19	41.39	38.49	n.a	0.068	40.92	43.37	0.398
Mean duration of hospitalization (hours)	121.71	n.a	n.a	72	115.2	156	108	0.814	56	139.64	0.151
Mean duration of work absenteeism (days)	2.47	n.a	1.84	1.44	2.1	3.13	n.a	0.001	2.36	2.56	0.572
Mean duration before seeing doctor (hours)	15.36	15.56	12.61	10.1	16.1	13.56	19.82	0.218	15.5	15.24	0.897

province) and southwest (Sichuan province), thus providing national representative data on the incidence and distribution of AGI in the population. We applied a face-to-face methodology like what has been used in other studies [14, 15]. Similar to those studies, we achieved a high average response rate of 97.2% when compared to telephone-administered surveys conducted in most developed countries. The high response rate was not only determined by the face-to-face interviewing, but also

greatly benefited from cooperative street residents' committee from selected residential district or villages.

Seasonality and geography

The results showed a seasonal distribution of AGI during the study period that is in line with the normally expectation- of high incidence of AGI in summer and low incidence in winter. The data showed its peak in spring and summer, and lower incidence in autumn and late

Table 3 Distribution of medical institutions utilized by acute gastroenteritis cases

Hospitals	Outpatients		Hospitalized	
	n = 346	%	n = 14	%
Grades of hospital				
Tertiary hospital	26	7.51	0	0.00
Secondary hospital	48	13.87	8	57.14
Primary hospital	81	23.41	5	35.71
Other hospital	191	55.20	1	7.14
Types of hospital				
General hospital	113	32.66	10	71.43
Children's hospital	4	1.16	0	0.00
Chinese medicine hospital	12	3.47	2	14.29
Other hospital	217	62.72	2	14.29
Classification of hospital				
Provincial and ministerial hospital	6	1.73	0	0.00
City hospital	31	8.96	1	7.14
County hospital	28	8.09	4	28.57
Community health center	110	31.79	6	42.86
Health service station	130	37.57	0	0.00
Private hospital	9	2.60	3	21.43
Individual clinic	31	8.96	0	0.00
Other hospital	1	0.29	0	0.00

winter, but relative high in early winter. The variation is likely to reflect the seasonal variation in infections with viral enteric pathogens in colder months, as seen in other studies [2, 16–18]. China is so large that temperatures and other meteorological factors typical of different seasons vary by province during the same period. For example, when the southern region is in summer, the northern region is in early spring, and in topography descends from the west to the east. As a result, the seasonality of AGI of bacterial origin is not synchronized. An analysis stratified by province and season would be needed to clarify such variations.

Determinants

We did not observe significant differences in the likelihood of having AGI in individuals with different levels of education, with different levels of income, or living in urbanized or rural areas. Travelling within two weeks was a significant risk factor, which is in accordance with what seen elsewhere [19].

Healthcare utilization

Care seeking behavior and utilization of healthcare services was high in all age groups. Half of surveyed individuals that reported AGI went to primary and lower health care institutions for medical care, which represents a higher rate of seeking medical care than estimated by

other countries such as the United States (19%) [19], Ireland (19.5%) [20], Canada (20.4%) [21], Australia (19.5%) [22] and Denmark (12%) [16], Cuba (17.1%–38.1%) [23] and Argentina (26%) [14]. It remains to be investigated if this disproportion is caused by the attention to disease, the attitude towards medicine, different medical insurance systems or differences in healthcare systems. The collection rate of biological samples in this survey was 14.3%, which is close to Canada (14.4%) [21] and Ireland (14.9%) [20], but lower than Australia (18.4%) [22] and the United States (21.1%) [19]. These results showed that the attention of hospitals in China to investigating the causes of AGI still needs to be improved and suggest that the central and local governments should increase awareness and investment to improve the sampling rate of patients' and the rate of laboratory diagnostics, to better grasp the etiology of acute gastroenteritis.

The hospitalization rate of AGI in our survey was 7.9%, suggesting a high proportion of severe cases of AGI in China. This index was not included in the previous domestic survey [7].

Nearly 80% of cases were treated with drugs, among which 62% use antibiotics, a much higher proportion estimated in other countries (US: 8.3%, Ireland: 5.6%, Canada: 3.8%, Australia: 3.6%, Italy: 6.5% Argentina: 7%, Cuba: 6.5–18.9%) [2, 14, 19–21]. Overuse or misuse of antimicrobials may lead to antimicrobial resistance, an important public health concern globally. These estimates suggest that there is still a long way to go to popularize the legal knowledge of the scientific use of antibiotics in China.

Domestic comparison

The estimated incidence of AGI in China of 0.28 (95%CI: 0.23–0.34) episodes per person was lower than the rate of 0.56 (95% CI: 0.56–0.57) episodes per person per year estimated by the previous survey conducted between 2010 to 2011 [7]. However, they were like the estimation of 0.31 episodes per person per year calculated by reviewing scientific literature [24]. The differences in the estimates of the different surveys may be explained by several factors. First, differences in the selection of provinces in the survey design, since China has great population diversity, which may also be reflected in diarrhea incidence and risk factors. All the provinces selected by the previous survey were southern provinces. On the contrary, our study took place in eight provinces that represent all the traditional seven regions of China [25, 26]. Second, in the past several years, the Chinese government has continuously improved the food safety regulation systems, developed innovative regulation mechanisms, established a structure for developing food safety standards and successfully dealt with intentional

food safety issues [27]. Risk communication was also enhanced. As examples, many ministries and departments jointly organized the Food Safety Awareness Week and Open Day. Several activities based on data generated from greatly improved foodborne disease surveillance since 2011 were also implemented. In 2013 alone, 120,000 supervision staff, more than 4,000 experts and scholars, and 35 million employees participated in activities. Hundreds of media issues nearly 20,000 news reports and over 300,000 micro-blogging topics [27]. These activities have also had an impact on public awareness and engagement in food safety practices. The overall food safety status has improved steadily and may be reflected in the burden of diarrheal diseases.

International comparison

Several cross-sectional surveys have been conducted in other countries to estimate the prevalence and distribution of AGI. In America [15, 28–38], Europe [2, 16–18, 39–44], Oceania [45, 46], Asia [47] and Africa [48].

The estimated incidence of AGI in China is comparable to similar retrospective studies conducted in Sweden and France, which reported 0.31 and 0.33 episodes of AGI each year, respectively, despite slightly different case definitions and recall periods. It is also similar to estimates of prospective follow-up studies in The Netherlands (0.28) (ranging from 0.42 to 1.66 episodes/person-year) [49]. However, it is lower than observations in almost all the similar cross-sectional studies conducted in other countries or regions, no matter what interviewing methods they used, and higher than rate from England and Wales (0.19) [50]. Nevertheless, comparisons between countries need to consider the varying case definitions, interviewing and sampling methods, and general differences in populations.

Our study used criterion for the identification of AGI recommended by an international collaboration focusing on burden of foodborne illnesses [51]. However, most studies, unlike ours, used a telephone survey; some of them used random digit dialing techniques or modified version to select participants [2, 16, 18, 47]. Other possible explanations for the differences in AGI incidence include cultural aspects and likelihood to answer questions openly. Chinese people are not used to talk about their health status in front of strangers, no matter if in public or in private, especially in big cities like Beijing and Shanghai. Furthermore, our survey had a very high response rate. With the help of social workers in resident's committees or villager's committees, almost all the selected household participated, and we had a response rate of 97%. In contrast, the response rate in other countries was below 70%, but there is a possibility that people who have AGI recently are more inclined to respond.

Other explanations for differences in reported incidence of diarrhea may include differences in risk factors such as food consumption and preparation habits, food contamination, or environmental factors. Furthermore, 59.3% of individuals reported that consuming suspected food was the cause of their AGI which is much higher than the proportion in Hong Kong (45.0%) [47] and Ireland (18.5%) [20]. The most self-reported suspected food categories (meat, fruit and vegetable) was correlated with the tendency of foodborne outbreaks [52] and diseases surveillance [53] in the past 10 years.

Limitations and strengths

In this survey, the samples were weighted and standardized according to the result of 6th national population census in 2010 with the structure of age and gender directly, to make the survey sample representative. A sample weight was assigned to each sample person. Sample weights were considered measures of the number of people represented by the particular sample person. However, there are still some biases and lack of consideration in the survey: (i) We did not consider design effects (DEFF) when designing the survey, so the results may be limited to the sample and cannot be extrapolated to the entire population. However, standardizing the sample and overestimating the loss rate of follow-up and relative risk could neutralize the bias of ignoring the DEFF; (ii) Inevitably, young and middle aged subgroup whose residency is in rural area but worked in the cities were not sampled according to the characteristics of the census, the occurrence of left-behind elderly and children in the sample were higher than reality. We generated the weights to adjust the target population. However, the adjustment works only if there is someone were sampled in the subgroup in someplace, for those didn't, there is no adjustment which might result in bias. (iii) In addition, the selection of survey sites failed to give full consideration to China's western minority areas, such as Tibet and Xinjiang. At the same time, some groups of group-life accommodation (such as inpatients in hospitals, elderly people in nursing homes, students in long-term accommodation, prisoners in prisons and officers and soldiers in barracks) were not covered. This might have introduced selection bias. (iv) No question about food preparation preferences and personal hygiene were involved in the questionnaire, and further research should examine these so that appropriate prevention measures can be put in place. (v) Our study suffered from the limitations of all cross-sectional surveys. Its results reflect only real time data from the survey sample and is difficult to make causal inferences. (vi) As a cross-sectional survey, this study should use prevalence rate ratios instead of odds ratios. Furthermore, to describe those AGI-like

acute diseases, we should also choose the prevalence rate ratios, since using odds ratios might amplifying the numbers.

Impact

Despite these limitations mentioned above, our estimates provide evidence on the incidence of AGI in the population of mainland China in seven regions for the first time. Due to underdiagnosis and underreporting, such data are not available from public health surveillance. However, it is crucial to demonstrate the true burden of diarrheal diseases in the population. This burden is not only of a health nature, but also social and economic. For example, our estimates demonstrated that at least 17% of respondents or caretakers with AGI loss a mean 3 days of work due to the illness. If extrapolated to the population, this equates to approximately 1.5 million working days loss due to AGI annually. Together with healthcare costs, this represents a substantial burden nationally. When combined with evidence on the contribution of different causative agents for the overall incidence of AGI, these estimates form the basis to estimate the burden of food-borne diseases in China.

Abbreviations

AGI	Acute gastrointestinal illness
SRS	Simple random sampling
CDC	Centers for Disease Control and Prevention
CI	Confidence interval
REF	References
OR	Odds ratio
SAS	Statistics Analysis Systems

Acknowledgements

The authors thank the participating Provincial, including municipal Centers for Disease Control and their Regional Centers for Disease Control and Prevention in Beijing, Jilin, Shanghai, Jiangsu, Henan, Guangdong, Sichuan, and Gansu, for their support in doing the face-to-face survey and data collection. The authors gratefully thank Dr. Yan Chen, for providing advice on the survey, and thank for the China Scholarship Council (No. 202009110012)

Authors' contributions

Jikai Liu: Writing-Introduction, partial Discussion, and partial Result. Baozhang Luo: Data curation and statistics, Funding acquisition, Writing-Methodology and Visualization. Yijing Zhou: Writing: partial Result, Investigation. Xiaochen Ma: Data curation, Resources, Investigation, and Writing—review. Junhua Liang: Resources, Data curation, Investigation, and Writing—partial Discussion. Xianglai Sang: Data curation, Resources, Investigation, and Writing—review. Le Lyu: Data curation, Resources, Investigation, and Writing—review. Wen Chen: Data curation, Resources, Investigation, and Writing—review. Pengyu Fu: Data curation, Resources, Investigation, and Writing—review. Hong Liu: Survey and questionnaire design. Data curation, Supervision. Shiqi Zhen: Data curation, Supervision, Resources, Investigation. Chao Wang: Data collection and curation and Resources, Investigation. Yangbo Wu: Data collection and curation and Resources, Investigation. Qiong Huang: Data curation, Resources, Investigation, and Writing—review. Xiaocheng Liang: Data curation, Supervision. Resources, Investigation. Guangda Bai: Data curation, Supervision. Resources, Investigation. Zhen Lan: Data curation, Supervision, Resources, Investigation. Shufang Zhang: Data curation, Supervision, Resources, Investigation. Ning Li: Validation. curation, supervision, Project administration. Yongning Wu: Curation, Supervision, Yunchang Guo: Conceptualization, Project administration, Funding acquisition. The author(s) read and approved the final manuscript.

Funding

This work was supported by Food Safety Research Unit (grant number 2019RU014) of Chinese Academy of Medical Science and Excellent Young Talents Project of Shanghai Public Health Three-year Action Plan (grant number GWV-10.2-YQ21).

Availability of data and materials

All data generated or analyzed during this study are included in this article.

Declarations

Ethics approval and consent to participate

Ethical approvals were obtained from the China National Center for Food Safety Risk Assessment's Ethical Board & Ethical approval number 2013–01. The study which has no interventions and treatment measures was in Accordance with *Measures for the Ethical Review of Biomedical Research Involving Humans* issued by the Ministry of Health. All participants gave written informed consent to participate in the study. All data collected including the names, addresses, and telephone numbers of the participant were kept confidential. Each participant was asked to read and sign a consent form that informed about the purpose of the survey before the questionnaire was administered.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interest.

Author details

¹NHC Key Laboratory of Food Safety Risk Assessment, Food Safety Research Unit (2019RU4) of Chinese Academy of Medical Science, China National Center for Food Safety Risk Assessment, Beijing, China. ²Shanghai Municipal Center for Disease Prevention and Control, Shanghai, China. ³Jiangsu Provincial Center for Disease Prevention and Control, Nanjing, China. ⁴Beijing Center for Disease Prevention and Control, Beijing, China. ⁵Guangdong Provincial Center for Disease Control and Prevention, Guangzhou, China. ⁶Gansu Provincial Center for Disease Control and Prevention, Lanzhou, China. ⁷Jilin Provincial Center for Disease Control and Prevention, Changchun, China. ⁸Sichuan Provincial Center for Disease Control and Prevention, Chengdu, China. ⁹Henan Provincial Center for Disease Control and Prevention, Zhengzhou, China.

Received: 9 February 2022 Accepted: 28 February 2023

Published online: 08 May 2023

References

- Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleeschauwer B, et al. World Health Organization Estimates of the Global and Regional Disease Burden of 22 Foodborne Bacterial, Protozoal, and Viral Diseases, 2010: A Data Synthesis. *PLoS Med*. 2015;12:e1001921.
- Scavia G, Baldinelli F, Busani L, Caprioli A. The burden of self-reported acute gastrointestinal illness in Italy: a retrospective survey, 2008–2009. *Epidemiol Infect*. 2012;140:1193–206.
- Flint JA, van Duynhoven YT, Angulo FJ, DeLong SM, Braun P, Kirk M, et al. Estimating the burden of acute gastroenteritis, foodborne disease, and pathogens commonly transmitted by food: an international review. *Clin Infect Dis*. 2005;41:698–704.
- Haagsma JA, Geenen PL, Ethelberg S, Fetsch A, Hansdotter F, Jansen A, et al. Community incidence of pathogen-specific gastroenteritis: reconstructing the surveillance pyramid for seven pathogens in seven European Union member states. *Epidemiol Infect*. 2012;141:1–15.
- Pires SM, Jakobsen LS, Ellis-Iversen J, Pessoa J, Ethelberg S. Burden of Disease Estimates of Seven Pathogens Commonly Transmitted Through Foods in Denmark, 2017. *Foodborne Pathog Dis*. 2019;17:322–339.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, et al. Foodborne illness acquired in the United States—Major pathogens. *Emerg Infect Dis*. 2011;17:7–15.

7. Chen Y, Yan WX, Zhou YJ, Zhen SQ, Zhang RH, Chen J, et al. Burden of self-reported acute gastrointestinal illness in China: a population-based survey. *BMC Public Health*. 2013;13:1–10.
8. Zhou YJ, Dai Y, Yuan BJ, Zhen SQ, Tang Z, Wu GL, et al. Population-based estimate of the burden of acute gastrointestinal illness in Jiangsu province, China, 2010–2011. *Epidemiol Infect*. 2013;141:944–52.
9. Sang XL, Liang XC, Chen Y, Li JD, Li JG, Bai L, et al. Estimating the burden of acute gastrointestinal illness in the community in Gansu Province, north-west China, 2012–2013. *BMC Public Health*. 2014;14:1–9.
10. Velu R, Naidu GM. A review of current survey sampling methods in marketing research (telephone mall intercept and panel surveys). Elsevier. p. 533–554.
11. Watson EK, Firman DW, Heywood A, Hauquitz AC, Ring I. Conducting regional health surveys using a computer-assisted telephone interviewing method. *Aust N Z J Public Health*. 1995;19:508–511.
12. Population Census Office under the State Council (PCO). Tabulation on the 2010 Population Census of the People's Republic of China by Township. China Statistics Press; Beijing, Book. 2012;3:1–915.
13. Majowicz SE, Hall G, Scallan E, Adak GK, Gauci C, Jones TF, et al. A common, symptom-based case definition for gastroenteritis. *Epidemiol Infect*. 2008;136:886.
14. Thomas MK, Perez E, Majowicz SE, Reid-Smith R, Albil S, Monteverde M, et al. Burden of acute gastrointestinal illness in Galvez, Argentina, 2007. *J Health Popul Nutr*. 2010;28:149–58.
15. Thomas MK, Perez E, Majowicz SE, Reid-Smith R, Olea A, Diaz J, et al. Burden of acute gastrointestinal illness in the Metropolitan region, Chile, 2008. *Epidemiol Infect*. 2011;139:560–71.
16. Müller L, Korsgaard H, Ethelberg S. Burden of acute gastrointestinal illness in Denmark 2009: a population-based telephone survey. *Epidemiol Infect*. 2012;140:290–8.
17. Baumann-Popczyk A, Sadkowska-Todys M, Rogalska J, Stefanoff P. Incidence of self-reported acute gastrointestinal infections in the community in Poland: a population-based study. *Epidemiol Infect*. 2012;140:1173–84.
18. Wilking H, Spitznagel H, Werber D, Lange C, Jansen A, Stark K. Acute gastrointestinal illness in adults in Germany: a population-based telephone survey. *Epidemiol Infect*. 2013;141:2365–75.
19. Jones TF, McMillan MB, Scallan E, Frenzen PD, Cronquist AB, Thomas S, et al. A population-based estimate of the substantial burden of diarrhoeal disease in the United States; FoodNet, 1996–2003. *Epidemiol Infect*. 2007;135:293.
20. Scallan E, Fitzgerald M, Collins C, Crowley D, Daly L, Devine M, Igoe D, Quigley T, Robinson T, Smyth B. Acute gastroenteritis in northern Ireland and the Republic of Ireland: a telephone survey. *Commun Dis Public Health*. 2004;7:61–7.
21. Majowicz SE, Horrocks J, Bocking K. Demographic determinants of acute gastrointestinal illness in Canada: a population study. *BMC Public Health*. 2007;7:1–8.
22. Scallan E, Majowicz SE, Hall G, Banerjee A, Bowman CL, Daly L, et al. Prevalence of diarrhoea in the community in Australia, Canada, Ireland, and the United States. *Int J Epidemiol*. 2005;34:454–60.
23. Gabriel OO, Jaime A, Mckensie M, Auguste A, Pérez E, Indar L. Estimating the Burden of Acute Gastrointestinal Illness: A Pilot Study of the Prevalence and Underreporting in Saint Lucia, Eastern Caribbean. *J Health Popul Nutr*. 2013;31(4 Suppl 1):S3.
24. Mao XD, Hu JF, Liu X. Epidemiological burden of bacterial foodborne diseases in China—Preliminary study - CNKI. *Chin J Food Hyg*. 2011;02:132–6.
25. Zhao J. *New Chinese Natural Geography*, first edition. Higher Education Press; Beijing, Book. 2015:1–464.
26. Jiehua L, Qin L. The characteristics, influence and coping strategies of the new form of aging society in China: interpretation based on the data of the seventh census. *Popul Econ*. 2021;248:13–24.
27. Jen JJC. Food safety in China. 2017.
28. Almaro CV, Ballal ML, Chey WD, Nordstrom C, Khanna D, Spiegel BMR. Burden of Gastrointestinal Symptoms in the United States: Results of a Nationally Representative Survey of Over 71,000 Americans. *Am J Gastroenterol*. 2018;113:1701–10.
29. Herikstad H, Yang S, van Gilder TJ, Vugia D, Hadler J, Blake P, et al. A population-based estimate of the burden of diarrhoeal illness in the United States: FoodNet, 1996–7. *Epidemiol Infect*. 2002;129:9–17.
30. Prieto PA, Finley RL, Muchaal PK, Guerin MT, Isaacs S, Domínguez AC, et al. Burden of self-reported acute gastrointestinal illness in Cuba. *J Health Popul Nutr*. 2009;27:345–57.
31. Gabriel OO, Jaime A, Mckensie M, Auguste A, Pérez E, Indar L. Estimating the burden of acute gastrointestinal illness: a pilot study of the prevalence and underreporting in Saint Lucia, Eastern Caribbean. *J Health Popul Nutr*. 2013;31:4 SUPPL.1.
32. Sargeant JM, Majowicz SE, Snelgrove J. The burden of acute gastrointestinal illness in Ontario, Canada, 2005–2006. *Epidemiol Infect*. 2008;136:451–60.
33. Thomas MK, Murray R, Nesbitt A, Pollari F. The Incidence of Acute Gastrointestinal Illness in Canada, Foodbook Survey 2014–2015. *Can J Infect Dis Med Microbiol*. 2017;2017:5956148.
34. Thomas MK, Majowicz SE, MacDougall L, Sockett PN, Kovacs SJ, Fyfe M, et al. Population distribution and burden of acute gastrointestinal illness in British Columbia, Canada. *BMC Public Health*. 2006;6:307.
35. Glasgow LM, Forde MS, Antoine SC, Pérez E, Indar L. Estimating the burden of acute gastrointestinal illness in Grenada. *J Health Popul Nutr*. 2013;31:4 SUPPL.1.
36. Thomas MK, Perez E, Majowicz SE, Reid-Smith R, Albil S, Monteverde M, et al. Burden of Acute Gastrointestinal Illness in Galvez, Argentina, 2007. 2007.
37. Persuad S, Mohamed-Rambaran P, Wilson A, James C, Indar L. Determining the community prevalence of acute gastrointestinal illness and gaps in surveillance of acute gastroenteritis and foodborne diseases in Guyana. *J Health Popul Nutr*. 2013;31:4 SUPPL.1.
38. Harper SL, Edge VL, Ford J, Thomas MK, Pearl DL, Shirley J, et al. Acute gastrointestinal illness in two Inuit communities: Burden of illness in Rigolet and Iqaluit. *Canada Epidemiol Infect*. 2015;143:3048–63.
39. Hansdotter FI, Magnusson M, KuhlmannBerenzon S, Hulth A, Sundstrom K, Hedlund KO, et al. The incidence of acute gastrointestinal illness in Sweden. *Scand J Public Health*. 2015;43:540–7.
40. Kuusi M, Aavitsland P, Gondrosen B, Kapperud G. Incidence of gastroenteritis in Norway - A population-based survey. *Epidemiol Infect*. 2003;131:591–7.
41. Viviani L, van der Es M, Irvine L, Tam CC, Rodrigues LC, Jackson KA, et al. Estimating the incidence of acute infectious intestinal disease in the community in the UK: a retrospective telephone survey. *PLoS One*. 2016;11:e0146171.
42. van Cauteren D, de Valk H, Vaux S, le Strat Y, Vaillant V. Burden of acute gastroenteritis and healthcare-seeking behaviour in France: a population-based study. *Epidemiol Infect*. 2012;140:697–705.
43. Gauci C, Gilles H, O'Brien S, Mamo J, Stabile I, Ruggeri FM, et al. The magnitude and distribution of infectious intestinal disease in Malta: a population-based study. *Epidemiol Infect*. 2007;135:1282–9.
44. Sävje-Söderbergh M, Toljander J, Bylund J, Simonsson M. Burden of gastrointestinal illness in Sweden-SMS as a tool for collecting self-reported gastrointestinal illness. *Epidemiol Infect*. 2019. <https://doi.org/10.1017/S0950268819002103>.
45. Lake RJ, Adlam SB, Perera S, Campbell DM, Baker MG. The disease pyramid for acute gastrointestinal illness in New Zealand. *Epidemiol Infect*. 2010;138:1468–71.
46. Hall GV, Kirk MD, Ashbolt R, Stafford R, Lalor K, Bell R, et al. Frequency of infectious gastrointestinal illness in Australia, 2002: Regional, seasonal and demographic variation. *Epidemiol Infect*. 2006;134:111–8.
47. Ho SC, Chau PH, Fung PK, Sham A, Nelson EA, Sung J. Acute gastroenteritis in Hong Kong: a population-based telephone survey. *Epidemiol Infect*. 2010;138:982–91.
48. Patterson K, Clark S, Berrang-Ford L, Lwasa S, Namanya D, Twebaze F, et al. Acute gastrointestinal illness in an African Indigenous population: The lived experience of Uganda's Batwa. *Rural Remote Health*. 2020;20:5141.
49. de Wit M, Koopmans M, Kortbeek L, Wannet WJB, Vinjé J, van Leusden F, et al. Sensor, a population-based cohort study on gastroenteritis in the Netherlands, incidence and etiology. *Am J Epidemiol*. 2001;154:666–74.
50. Wheeler JG, Sethi D, Cowden JM, Wall PG, Rodrigues LC, Tompkins DS, et al. Study of infectious intestinal disease in England: rates in the community, presenting to general practice, and reported to national surveillance. The Infectious Intestinal Disease Study Executive. *BMJ*. 1999;318:1046–50.

51. Chen Y, Ford L, Hall G, Dobbins T, Kirk M. Healthcare utilization and lost productivity due to infectious gastroenteritis, results from a national cross-sectional survey Australia 2008–2009. *Epidemiol Infect.* 2016;144:241–6.
52. Li W, Pires SM, Liu Z, Ma X, Liang J, Jiang Y, et al. Surveillance of foodborne disease outbreaks in China, 2003–2017. *Food Control.* 2020;118:107359.
53. Liu J, Bai L, Li W, Han H, Fu P, Ma X, et al. Trends of foodborne diseases in China: lessons from laboratory-based surveillance since 2011. *Front Med.* 2018;12:48–57.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

