

RESEARCH ARTICLE

Open Access

Gender-specific differences in risk for intimate partner violence in South Korea

Minjee Lee¹, Katherine M Stefani² and Eun-Cheol Park^{3*}

Abstract

Background: Various risk factors of intimate partner violence (IPV) have been found to vary by gender. South Korea has one of the highest prevalences of IPV in the world; however, little is known about potential risk factors of IPV and whether gender influences this relationship.

Methods: Using data from the 2006 Korea Welfare Panel Study, 8,877 married participants (4,545 men and 4,332 women) aged ≥ 30 years were included. Reported IPV was categorized as verbal or physical IPV and the association between IPV and related factors was assessed by multivariate logistic regression analysis.

Results: Women were significantly more likely than men were to report IPV victimization (verbal 28.2% vs. 24.4%; physical 6.9% vs. 3.4%). Women reported perpetrating verbal IPV against their partner more often than men did (26.7% vs. 25.3%). However, more men reported perpetrating physical violence against their wife (5.1% vs. 3.4%). A low perceived level of satisfaction with family and life were the strongest predictors of physical victimization and perpetration in IPV among men and women. Men dissatisfied with their family relationships had a 5.49 higher odds (95% CI, 2.91-10.37) of physical victimization than men satisfied with their family. Women dissatisfied with their family relationships had a 9.46 (5.21-17.19) higher odds of physical perpetration than women satisfied with their family. Moreover, alcohol intake was significantly associated with IPV perpetration and victimization in both genders.

Conclusion: Significant gender-specific differences were found among factors related to perpetrating violence and being a victim of violence among adults in heterosexual relationships in South Korea.

Keywords: Gender, Intimate partner violence, South Korea

Background

Intimate partner violence (IPV) is a worldwide public health problem as well as a serious social problem in South Korea [1-3]. IPV is characterized as any behavior within an intimate relationship that causes physical, psychological, or sexual harm of one partner to another [1]. According to the 2010 national survey in South Korea, the prevalence of reported IPV was 53.8%, and 81.9% of this violence was perpetrated by husbands against their wives [4].

IPV causes a wide range of negative effects on the health of women [5-7] and children, such as injury, chronic pain, gastrointestinal problems, sexually transmitted diseases, depression, and post-traumatic stress disorder [8-10]. In

addition, a significant number of deaths among women are considered to result from IPV [11,12]. Moreover, children who witness IPV in their home are also significantly more likely to experience or perpetrate IPV than are children who do not [13].

The prevalence of reported partner violence varies greatly (15%-71%) among various countries [1,14,15]. In South Korea, the special law to Prevent Family Violence and Protect the Victim mandates that Korean nationals be surveyed triennially. According to this survey, the prevalence of reported IPV dropped from 53.8% in 2010 to 45.5% in 2013 [16].

IPV typically results due to gender inequality and is frequently considered a form of gender-based violence [1]. Violence against women has been the focus of most research, yet little is known about the prevalence of violence perpetrated against men who are in a heterosexual

* Correspondence: ecpark@yuhs.ac

³Department of Preventive Medicine and Public Health, Institute of Health Services Research, Yonsei University College of Medicine, 134 Shinchon-dong, Seodaemun-gu, 120-752 Seoul, South Korea
Full list of author information is available at the end of the article

relationship [17,18]. Some studies have found the prevalence of violence against men to be equivalent to that against women [18,19]. Similarly, in South Korea, previous research has highlighted the scope and risk factors of violence against women; however, few studies have investigated the prevalence of violence against men or the possible risk factors associated with male perpetration and victimization [20,21].

Factors associated with IPV victimization among women include pregnancy, depressive symptoms, smoking, alcohol consumption, low socioeconomic status, experiencing IPV during childhood, and witnessing IPV perpetration against their mother [22-28]. Tumwesigye and colleagues [27] investigated IPV victimization among women in Uganda and found socioeconomic status including education, employment status, income, and education level as well as the employment status of the partner to be potential risk factors. Furthermore, Lemon and colleagues [28] reported that current smoking and frequent alcohol use are related to IPV victimization among women living in the US (Rhode Island). Although few studies have examined IPV victimization among men, alcohol consumption, low socioeconomic status, experiencing IPV during childhood, and witnessing IPV perpetration against their mother were factors associated with IPV perpetration [29-33]. In addition, a study that compared both men and women found alcohol dependency to be associated with severe physical IPV perpetration in the New Zealand Birth Cohort [33].

In addition, lacking a social network, emotional support, and having low perceived life satisfaction were found to be related with IPV among men and women [2,33-37]. For example, South Korean women with poor social/support networks were more likely to experience IPV victimization [2,35,36] as well as continued abuse [37]. Among Chinese men and women, life dissatisfaction was related with IPV victimization [34]. Furthermore, a weak social support system was strongly associated with physical IPV victimization among women in New Zealand [33].

Although the aforementioned studies identified some factors associated with IPV, the limited sample size, lack of gender-specific analyses, and a lack of consideration for IPV perpetration among women and IPV victimization among men warrant further study. Furthermore, previous research has not considered factors related to each type of IPV (verbal and physical). To this end, we investigated whether gender-specific differences exist in the prevalence of IPV as well as the type of the violence that was perpetrated or experienced.

Methods

Data and sample

Data from the nationally representative 2006 Korea Welfare Panel Study (KOWEPS) performed by the Korean Institute

of Social and Health Affairs in conjunction with Social Welfare Research Institute of Seoul National University were used for this study. Details of this study have been published elsewhere [38,39]. Briefly, KOWEPS is a comprehensive dataset that provides a variety of information on families and individuals with respect to their social service needs, health care utilization patterns, economic and demographic background, sources of income, and subjective emotional and behavioral health status. A stratified, multistage, probability design was used. Men and women older than 19 were selected from sampling units using household registries. In total, 7,072 households participated in the survey. Trained interviewers conducted all surveys at participants' homes, and all participants provided informed consent before participating in the survey.

In the 2006 dataset, 18,856 men and women were recruited. Of them, 16,084 (85.29%) men and women aged 19 or older participated in the survey. For our analyses, only those who were older than 30 years old and married were included leaving 9,667 men and women. The 16 participants who were younger than 30 and married were excluded from our analysis because, according to the National Statistical Office in South Korea, the average age of marriage among males and females was 32.1 was and 29.4 in 2012, respectively [40]. By limiting our analysis to those 30 and older, we hoped to include a more representative population of married couples. After exclusion for those missing any relevant data ($n = 790$), a total of 8,877 participants (4,545 men and 4,332 women) who reported being married at the time of the survey were included in the analysis.

Measurement of IPV

During data collection, participants were asked 13 questions pertaining to the level and type of violence experienced in their marriage over the past 12 months. Verbal IPV was assessed by asking how often in the past 12 months their spouse was (1) insulting, (2) made a malicious remark, or (3) threatened them. Physical IPV was assessed across ten violent activities. Respondents were asked how often in the past 12 months their spouse perpetrated the following physically violent activities at him/her: (1) threw something, (2) pushed, (3) slapped, (4) kicked or punched, (5) used an object to hit, (6) beat, (7) threatened using a weapon like a knife, (8) choked, (9) caused a sprain or bruise, or (10) caused him/her to be hospitalized after a violent encounter. These questions were adopted from the Conflict Tactics Scale [39]. In addition, participants were also asked how often they perpetrated any of these acts against their spouse, and their answers were recorded as never, 1-2 times, 3-5 times, 6-10 times, or >10 times throughout the previous 12 months. A person was considered to have experienced or perpetrated IPV if a violent event occurred once or more over the past 12 months,

and this was recorded as the binary outcome variable (yes or no) in our analysis.

Independent variables

The statistical models used in this study were created based on variables reported in previous studies [15,41]. Nine variables from three domains (related to socio-demographic factors, family/life satisfaction, and health behaviors) that have been robustly linked to IPV in epidemiological studies for IPV victimization among women and IPV perpetration among men were selected [27,28,42]. The socio-demographic factors include age, education, household income, employment status, and the perceived wealth during childhood. Participants' subjective levels of satisfaction in either family relationships or one's personal life was also measured along with health behaviors such as smoking and alcohol intake [27,28,42].

Participants were divided into five age groups, 30–39, 40–49, 50–59, 60–69, and ≥ 70 years, for the analysis. In addition, education level was stratified into three groups based on the highest level of education achieved as elementary school, middle or high school, or university or higher. Income was calculated according to the equivalized household income equation as the sum of the total household income from all sources including earned income, income from assets, and miscellaneous income divided by the square root of the number of household members. The equivalized income was then divided into quartiles [39]. Employment status was categorized as either being employed full time, part time, being self-employed, or unemployed. Participants' perceived level of wealth during childhood was categorized as either poor, average or wealthy. The perceived level of satisfaction with family and life was recorded as not satisfied, neutral, or satisfied. Smoking was categorized as never or ever. Moreover, participants were divided into four groups based on the average number of drinks consumed at one time as a nondrinker, light drinker (1–4 drinks), moderate drinker (5–9 drinks), or heavy drinker (>10 drinks).

Statistical analysis

Descriptive statistics were used to describe participant characteristics and report the number and percentage of participants for each variable. In addition, the prevalence of IPV was calculated for all variables. Odds ratios (OR) with 95% confidence intervals (CI) were calculated to measure the strength of the association between IPV and all possible IPV-related factors in this study population. Multivariate logistic regression models, with IPV as the dependent variable, were used to calculate gender-specific ORs. Fully adjusted ORs were calculated after controlling for all potential confounders (age, education, household income, employment status, perceived wealth during

childhood, satisfaction with family and life, smoking, and alcohol intake). Sampling weights were also added to aid in generalizing our findings to the entire population of South Korea. All statistical analyses were performed using SAS version 9.2 (SAS Inc., Cary, NC, USA), and a p -value > 0.05 was considered statistically significant.

Results

Gender-specific data on the characteristics of the study population and the prevalence of IPV are shown in Tables 1 and 2, respectively.

The majority of men reported working full time (47.6%), had at least a middle or high school education (47.2%), were non-smokers (51.7%), and moderate drinkers (30.4%). For men, the prevalence of verbal IPV was 24.4% for victimization and 25.3% for perpetration. The prevalence of physical IPV was 3.4% for victimization and 5.1% for perpetration (Table 1).

The majority of women in this study population were employed full time (48.3%) and had at least a middle or high school education (47.7%). In addition, most women were nonsmokers (98.2%) and nondrinkers (69.9%). For women, the prevalence of verbal IPV was 28.2% for victimization and 26.7% for perpetration. The prevalence of physical IPV was 6.9% for victimization and 3.4% for perpetration (Table 2). For both men and women, the prevalence of victimization and perpetration may be overlapped, thus should not be considered independent measures.

Women were significantly more likely than men were to report being a victim of IPV (verbal: 28.2% vs. 24.4% [$p < .0001$], physical: 6.9% vs. 3.4% [$p < .0001$]). In addition, women tended to perpetrate verbal violence against their spouse more often than men were (26.7% vs. 25.3%). However, 5.1% of men and 3.4% of women reported perpetrating physical violence against their spouse. A similar proportion of men and women reported male-to-female violence, yet more women than men reported female-to-male violence.

Among males, household income, perceived wealth during childhood, and smoking were not significantly associated with IPV victimization nor with IPV perpetration. Men with a middle or high school education were significantly less likely to report perpetrating verbal IPV than were men with only an elementary school education. In addition, men employed part time were significantly more likely to perpetrate physical IPV than men employed full time were. Moreover, having a high level of satisfaction with family relationships and one's personal life made men less likely to perpetrate both verbal and physical IPV. Compared with men who were satisfied with their family relationships, the adjusted OR for men dissatisfied with their family relationships was 2.44 (95% CI: 1.66-3.58) for verbal IPV victimization, 5.49 (95% CI: 2.91-10.37) for

Table 1 Descriptive data across the type of intimate partner violence among men

Variables	Total %		Victim						Perpetrator					
			Verbal %	p-value	Physical %	p-value	Verbal %	p-value	Physical %	p-value				
Age (years)														
30-39	401	8.82	104	25.9	<.0001	26	6.5	<.0001	105	26.2	<.0001	37	9.2	<.0001
40-49	1103	24.27	322	29.2		69	6.3		317	28.7		77	7.0	
50-59	955	21.01	254	26.6		35	3.7		267	28.0		54	5.7	
60-69	750	16.5	159	21.2		12	1.6		176	23.5		28	3.7	
70+	1336	29.39	269	20.1		13	1.0		286	21.4		35	2.6	
Education														
None or elementary school	1056	23.23	253	24.0	0.936	23	2.2	0.021	278	26.3	0.645	38	3.6	0.034
Middle or high school	2143	47.15	525	24.5		75	3.5		541	25.2		123	5.7	
University or higher	1346	29.61	330	24.5		57	4.2		332	24.7		70	5.2	
Household income														
Quartile 1	1086	23.89	233	21.5	0.058	25	2.3	0.080	263	24.2	0.777	40	3.7	0.032
Quartile 2	1106	24.33	276	25.0		46	4.2		280	25.3		53	4.8	
Quartile 3	1162	25.57	305	26.2		45	3.9		303	26.1		74	6.4	
Quartile 4	1191	26.2	294	24.7		39	3.3		305	25.6		64	5.4	
Employment status														
Employed full time	2161	47.55	521	24.1	0.006	86	4.0	0.022	523	24.2	0.015	107	5.0	<.0001
Employed part time	989	21.76	276	27.9		36	3.6		288	29.1		77	7.8	
Self employed	491	10.8	121	24.6		17	3.5		127	25.9		17	3.5	
Unemployed	904	19.89	190	21.0		16	1.8		213	23.6		30	3.3	
Perceived wealth in childhood														
Poor	2146	47.22	520	24.2	0.843	67	3.1	0.319	559	26.0	0.431	110	5.1	0.920
Average	1878	41.32	465	24.8		73	3.9		470	25.0		93	5.0	
Wealthy	521	11.46	123	23.6		15	2.9		122	23.4		28	5.4	
Satisfaction with family relationships														
Satisfied	3682	81.01	814	22.1	<.0001	107	2.9	<.0001	839	22.8	<.0001	148	4.0	<.0001
Neutral	732	16.11	238	32.5		30	4.1		253	34.6		62	8.5	
Not satisfied	131	2.88	56	42.7		18	13.7		59	45.0		21	16.0	
Satisfaction with one's personal life														
Satisfied	1668	36.7	332	19.9	<.0001	37	2.2	<.0001	339	20.3	<.0001	55	3.3	<.0001
Neutral	2167	47.68	547	25.2		71	3.3		575	26.5		114	5.3	
Not satisfied	710	15.62	229	32.3		47	6.6		237	33.4		62	8.7	
Smoking														
Never	2351	51.73	512	21.8	<.0001	57	2.4	0.000	525	22.3	<.0001	91	3.9	0.000
Ever	2194	48.27	596	27.2		98	4.5		626	28.5		140	6.4	
Alcohol intake														
Nondrinker	1373	30.21	243	17.7	<.0001	25	1.8	<.0001	253	18.4	<.0001	42	3.1	<.0001
Light drinker (1-4 drinks)	1053	23.17	249	23.6		28	2.7		242	23.0		38	3.6	
Moderate drinker (5-9 drinks)	1381	30.39	380	27.5		53	3.8		422	30.6		77	5.6	
Heavy drinker (10 ≤ drinks)	738	16.24	236	32.0		49	6.6		234	31.7		74	10.0	
Total	4545	100	1108	24.4		155	3.4		1151	25.3		231	5.1	

Table 2 Descriptive data across the type of intimate partner violence among Women

Variables	Total %		Victim						Perpetrator					
			verbal %	p-value	physical %	p-value	verbal %	p-value	physical %	p-value				
Age (years)														
30-39	658	15.19	176	26.7	0.001	65	9.9	<.0001	197	29.9	<.0001	42	6.4	<.0001
40-49	1175	27.12	372	31.7		96	8.2		363	30.9		61	5.2	
50-59	877	20.24	269	30.7		71	8.1		244	27.8		25	2.9	
60-69	762	17.59	194	25.5		39	5.1		174	22.8		11	1.4	
70+	860	19.85	211	24.5		27	3.1		180	20.9		9	1.0	
Education														
None or elementary school	1400	32.32	404	28.9	0.448	78	5.6	0.061	358	25.6	0.185	21	1.5	<.0001
Middle or high school	2068	47.74	589	28.5		157	7.6		549	26.5		79	3.8	
University or higher	864	19.94	229	26.5		63	7.3		251	29.1		48	5.6	
Household income														
Quartile 1	1038	23.96	300	28.9	0.766	59	5.7	0.239	273	26.3	0.932	22	2.1	0.027
Quartile 2	1055	24.35	306	29.0		74	7.0		288	27.3		36	3.4	
Quartile 3	1098	25.35	304	27.7		87	7.9		297	27.0		39	3.6	
Quartile 4	1141	26.34	312	27.3		78	6.8		300	26.3		51	4.5	
Employment status														
Employed full time	2094	48.34	582	27.8	0.072	161	7.7	<.0001	591	28.2	0.000	97	4.6	<.0001
Employed part time	875	20.2	277	31.7		75	8.6		263	30.1		28	3.2	
Self employed	189	4.36	50	26.5		15	7.9		44	23.3		10	5.3	
Unemployed	1174	27.1	313	26.7		47	4.0		260	22.1		13	1.1	
Perceived wealth in childhood														
Poor	1548	35.73	463	29.9	0.179	118	7.6	0.316	440	28.4	0.050	47	3.0	0.052
Average	2219	51.22	605	27.3		146	6.6		587	26.5		89	4.0	
Wealthy	565	13.04	154	27.3		34	6.0		131	23.2		12	2.1	
Satisfaction with family relationships														
Satisfied	3396	78.39	841	24.8	<.0001	168	4.9	<.0001	811	23.9	<.0001	88	2.6	<.0001
Neutral	780	18.01	294	37.7		88	11.3		264	33.8		34	4.4	
Not satisfied	156	3.6	87	55.8		42	26.9		83	53.2		26	16.7	
Satisfaction with one's personal life														
Satisfied	1451	33.49	303	20.9	<.0001	58	4.0	<.0001	296	20.4	<.0001	37	2.5	0.000
Neutral	2216	51.15	654	29.5		154	6.9		596	26.9		71	3.2	
Not satisfied	665	15.35	265	39.8		86	12.9		266	40.0		40	6.0	
Smoking														
Never	4254	98.2	1199	28.2	0.800	292	6.9	0.775	1134	26.7	0.416	141	3.3	0.006
Ever	78	1.8	23	29.5		6	7.7		24	30.8		7	9.0	
Alcohol intake														
Nondrinker	3026	69.85	778	25.7	<.0001	179	5.9	0.000	721	23.8	<.0001	81	2.7	<.0001
Light drinker (1-4 drinks)	1046	24.15	345	33.0		88	8.4		342	32.7		44	4.2	
Moderate drinker (5-9 drinks)	213	4.92	81	38.0		24	11.3		75	35.2		17	8.0	
Heavy drinker (10 ≤ drinks)	47	1.08	18	38.3		7	14.9		20	42.6		6	12.8	
Total	4332	100	1222	28.2		298	6.9		1158	26.7		148	3.4	

physical IPV victimization, 2.54 (95% CI: 1.74-3.71) for verbal IPV perpetration, and 4.68 (95% CI: 2.66-8.27) for physical IPV perpetration. However, the adjusted OR among men who were dissatisfied with their personal life was 1.76 (95% CI: 1.39-2.22) for verbal IPV victimization, 3.07 (95% CI: 1.79-5.26) for physical IPV victimization, 1.69 (95% CI: 1.34-2.13) for verbal IPV perpetration, and 2.51 (95% CI: 1.60-3.95) for physical IPV perpetration when compared to men who were satisfied with their personal life. Among the measured health behaviors, men reporting high alcohol intake were more likely to have experienced verbal IPV victimization, physical IPV victimization, verbal IPV perpetration, and physical IPV perpetration. Compared with non-drinkers, the adjusted OR of heavy drinkers was 2.06 (95% CI: 1.65-2.58) for verbal IPV victimization, 2.54 (95% CI: 1.49-4.31) for

physical IPV victimization, 2.00 (95% CI: 1.60-2.50) for verbal IPV perpetration, and 2.81 (95% CI: 1.84-4.30) for physical IPV perpetration (Table 3).

Among women, age was significantly associated with all types of IPV, but household income, employment status, perceived wealth during childhood, and smoking were not. In addition, women with at least a middle or high school education were significantly less likely to report verbal IPV victimization and physical IPV perpetration than women with less education were. Similar to men, increased alcohol intake was associated with an increased odds of verbal and physical IPV perpetration. However, unlike men, alcohol drinking was not significantly associated with physical IPV victimization. Compared with non-drinkers, the adjusted ORs of moderate drinkers for verbal IPV victimization, verbal IPV perpetration, and physical

Table 3 Factors associated with intimate partner violence reported by men

Variables	Victimization				Perpetration			
	Verbal		Physical		Verbal		Physical	
Age (years)								
30-39	1.00		1.00		1.00		1.00	
40-49	1.16	(0.89 - 1.51)	0.92	(0.57 - 1.50)	1.11	(0.85 - 1.45)	0.70	(0.45 - 1.07)
50-59	0.98	(0.74 - 1.30)	0.46	(0.26 - 0.81)	1.02	(0.77 - 1.35)	0.52	(0.32 - 0.83)
60-69	0.75	(0.55 - 1.03)	0.18	(0.08 - 0.39)	0.81	(0.60 - 1.11)	0.36	(0.20 - 0.64)
70+	0.74	(0.54 - 1.02)	0.08	(0.04 - 0.20)	0.73	(0.53 - 1.00)	0.27	(0.15 - 0.51)
Education								
None or elementary school	1.00		1.00		1.00		1.00	
Middle or high school	0.81	(0.66 - 1.01)	0.69	(0.38 - 1.26)	0.81	(0.66 - 1.00)	1.09	(0.69 - 1.71)
University or higher	0.82	(0.62 - 1.07)	0.74	(0.37 - 1.50)	0.83	(0.63 - 1.08)	0.97	(0.56 - 1.70)
Employment status								
Employed full time	1.00		1.00		1.00		1.00	
Employed part time	1.13	(0.94 - 1.36)	0.73	(0.48 - 1.13)	1.18	(0.98 - 1.42)	1.45	(1.05 - 2.02)
Self employed	1.13	(0.88 - 1.45)	1.22	(0.68 - 2.19)	1.16	(0.90 - 1.48)	0.83	(0.47 - 1.44)
Unemployed	0.93	(0.74 - 1.17)	0.82	(0.44 - 1.54)	1.02	(0.82 - 1.28)	0.99	(0.61 - 1.60)
Satisfaction with family relationships								
Satisfied	1.00		1.00		1.00		1.00	
Neutral	1.68	(1.39 - 2.02)	1.58	(1.01 - 2.47)	1.73	(1.43 - 2.08)	2.38	(1.70 - 3.35)
Not satisfied	2.44	(1.66 - 3.58)	5.49	(2.91 - 10.37)	2.54	(1.74 - 3.71)	4.68	(2.66 - 8.27)
Satisfaction with one's personal life								
Satisfied	1.00		1.00		1.00		1.00	
Neutral	1.30	(1.10 - 1.54)	1.62	(1.05 - 2.51)	1.31	(1.11 - 1.55)	1.55	(1.08 - 2.21)
Not satisfied	1.76	(1.39 - 2.22)	3.07	(1.79 - 5.26)	1.69	(1.34 - 2.13)	2.51	(1.60 - 3.95)
Alcohol intake								
Nondrinker	1.00		1.00		1.00		1.00	
Light drinker (1-4 drinks)	1.50	(1.22 - 1.83)	1.46	(0.84 - 2.57)	1.36	(1.11 - 1.67)	1.17	(0.74 - 1.84)
Moderate drinker (5-9 drinks)	1.71	(1.40 - 2.08)	1.62	(0.97 - 2.70)	1.92	(1.58 - 2.33)	1.58	(1.05 - 2.38)
Heavy drinker (10 ≤ drinks)	2.06	(1.65 - 2.58)	2.54	(1.49 - 4.31)	2.00	(1.60 - 2.50)	2.81	(1.84 - 4.30)

Results are presented as adjusted odds ratios and (95% confidence intervals), and adjusted for age, education, household income, employment status, perceived wealth during childhood, satisfaction with family relationships, satisfaction with one's personal life, smoking, and alcohol intake.

IPV perpetration were 1.68 (95% CI: 1.24-2.27), 1.53 (95% CI: 1.13-2.08), and 2.15 (95% CI: 1.21-3.83), respectively. Moreover, women with a high level of family and life satisfaction were less likely to report verbal IPV victimization, physical IPV victimization, verbal IPV perpetration, and physical IPV perpetration; these findings are similar to that among men. Compared with women who were satisfied with their family relationships, the adjusted ORs among women dissatisfied with their relationships were 3.17 (95% CI: 2.23-4.52) for verbal IPV victimization, 6.68 (95% CI: 4.23-10.56) for physical IPV victimization, 2.78 (95% CI: 1.95-3.96) for verbal IPV perpetration, and 9.46 (95% CI: 5.21-17.19) for physical IPV perpetration. Moreover, the adjusted ORs among women dissatisfied with their personal life was 1.96 (95% CI: 1.55-2.49) for verbal IPV victimization, 2.52 (95% CI: 1.64-3.89) for physical IPV victimization, 2.36 (95% CI: 1.85-3.00) for verbal IPV perpetration, and 2.23 (95% CI: 1.22-4.08) for physical IPV perpetration when compared to women who reported being satisfied with their personal life (Table 4).

Discussion

In this nationally representative population of Korean men and women, significant gender-specific differences were evident for the prevalence of IPV and its associated factors. Factors significantly associated with an increased likelihood of IPV victimization and perpetration among men and women were the subjective measures of family and personal life satisfaction as well as alcohol intake. In addition, the prevalence of IPV victimization was significantly higher among women than among men.

Although gender-specific data on IPV are limited, our findings are consistent with estimates gathered from previous studies with smaller samples of South Korean adults and population-based studies conducted in other Asian countries [20,42-45], which found higher exposures to IPV among women than men.

Investigation into the physical perpetration of IPV revealed that males had slightly higher rates of perpetration than that of females (males, 5.1% vs. females, 3.4%). In addition, women reported a lower rate of experiencing

Table 4 Factors associated with intimate partner violence reported by women

Variables	Victimization				Perpetration			
	Verbal		Physical		Verbal		Physical	
Age (years)								
30-39	1.00		1.00		1.00		1.00	
40-49	1.23	(0.99 - 1.54)	0.75	(0.53 - 1.07)	1.04	(0.84 - 1.30)	0.79	(0.52 - 1.22)
50-59	1.06	(0.82 - 1.36)	0.63	(0.42 - 0.96)	0.86	(0.66 - 1.10)	0.43	(0.24 - 0.77)
60-69	0.69	(0.51 - 0.93)	0.31	(0.18 - 0.52)	0.57	(0.42 - 0.78)	0.22	(0.09 - 0.50)
70+	0.53	(0.38 - 0.75)	0.15	(0.08 - 0.28)	0.43	(0.30 - 0.61)	0.14	(0.05 - 0.39)
Education								
None or elementary school	1.00		1.00		1.00		1.00	
Middle or high school	0.77	(0.61 - 0.96)	0.70	(0.46 - 1.07)	0.73	(0.57 - 0.92)	1.07	(0.53 - 2.17)
University or higher	0.76	(0.56 - 1.02)	0.67	(0.39 - 1.13)	0.87	(0.64 - 1.17)	1.38	(0.62 - 3.11)
Satisfaction with family relationships								
Satisfied	1.00		1.00		1.00		1.00	
Neutral	1.66	(1.39 - 1.99)	2.43	(1.80 - 3.28)	1.48	(1.23 - 1.78)	1.94	(1.24 - 3.04)
Not satisfied	3.17	(2.23 - 4.52)	6.68	(4.23 - 10.56)	2.78	(1.95 - 3.96)	9.46	(5.21 - 17.19)
Satisfaction with one's personal life								
Satisfied	1.00		1.00		1.00		1.00	
Neutral	1.48	(1.25 - 1.76)	1.63	(1.16 - 2.29)	1.45	(1.22 - 1.73)	1.45	(0.92 - 2.27)
Not satisfied	1.96	(1.55 - 2.49)	2.52	(1.64 - 3.89)	2.36	(1.85 - 3.00)	2.23	(1.22 - 4.08)
Alcohol intake								
Nondrinker	1.00		1.00		1.00		1.00	
Light drinker (1-4 drinks)	1.38	(1.17 - 1.62)	1.21	(0.91 - 1.61)	1.45	(1.23 - 1.70)	1.13	(0.76 - 1.67)
Moderate drinker (5-9 drinks)	1.68	(1.24 - 2.27)	1.58	(0.98 - 2.53)	1.53	(1.13 - 2.08)	2.15	(1.21 - 3.83)
Heavy drinker (10 ≤ drinks)	1.47	(0.80 - 2.70)	1.78	(0.75 - 4.18)	1.83	(1.00 - 3.35)	3.18	(1.23 - 8.20)

Results are presented as adjusted odds ratios and (95% confidence intervals), and adjusted for age, education, household income, employment status, perceived wealth during childhood, satisfaction with family relationships, satisfaction with one's personal life, smoking, and alcohol intake.

male perpetration than that of experiencing victimization. However, the rate of female perpetration was roughly aligned with the rate of victimization that was reported by men. One interpretation for this discrepancy may be that men underreported IPV perpetration due to a social desirability bias. Similar disparities in reporting have been found in studies conducted in other countries [46].

The majority of research surrounding IPV has addressed violence against women, examining prevalence estimates of female victimization and male perpetration. In the current study, rates of verbal IPV victimization among females (28.2%) fell within the range of previous estimates (24.6%-55.0%) [47,48], although male rates of verbal perpetration (25.3%) were slightly lower than previous estimates (31.8%-42.3%) [49,50]. However, limitations between survey methods make comparing our results with previous studies difficult; inconsistencies in defining violence and variations in survey periods may have created discrepancies between these estimates.

Among men, low education level was positively associated with perpetrating verbal violence; however, among women, low education level was positively associated with both victimizing and perpetrating verbal IPV. Previous studies have reported similar associations and have strongly suggested that future violence prevention programs should aim to increase levels of education and understanding about IPV [47].

Our results also confirm the finding of other studies that IPV victimization and perpetration are likely to occur regardless of one's drinking habits [46,47]. However, we cannot determine whether alcohol is a risk factor of or a result of IPV due to the cross-sectional design of our study. Nevertheless, our findings support previous assertions that alcohol interventions may be a crucial component of future violence prevention programs [19,51,52].

Our study has important limitations. First, the prevalence of IPV was measured over the previous 12 months; therefore, these data may have underestimated the actual prevalence of IPV in this study population. Second, the KOWEPS dataset did not measure experiences with IPV during childhood such as any exposures to domestic violence or witnessing IPV perpetration among their parents. Third, the KOWEPS dataset did not ask participants who were married whether they live with their spouse. We cannot assume that all married couples live with their spouse and this factor may influence the prevalence of IPV; therefore, future studies should investigate whether this factor is associated with IPV. Fourth, IPV related to sexual abuse was not included in the KOWEPS survey. Last, the confidence intervals estimated in our analyses were wide; therefore, future prospective studies with large sample sizes are needed to better understand these relationships.

In this nationally representative study, we found that the prevalence of IPV and its associated factors were

gender specific. In particular, alcohol intake, family, and life satisfaction were strongly associated with both verbal and physical IPV among men and women. In addition, the prevalence of IPV victimization was significantly higher among women than among men. Moreover, older men who were dissatisfied with their family and personal life as well as heavy drinkers were more likely to be victims of physical IPV than their counterparts were. Furthermore, older women who were dissatisfied with their family and personal life as well as heavy drinkers were more likely to perpetrate physical IPV than their counterparts were. Large, prospective studies are needed to understand the etiology of these factors for the proper implementation of preventative measures to reduce IPV in South Korea.

Conclusions

Worldwide, gender differences in IPV have been reported and, among those countries, South Korea has one of the highest prevalences of IPV. However, the factors related to IPV in Korean adults are unknown.

We found significant gender-specific differences among the factors related to IPV. In addition, the prevalences for each the type of violence perpetrated and victimized varied significantly by gender.

This is the first study to utilize nationally representative data to investigate the prevalence and risk factors of IPV in South Korea. This study measured IPV as a self-reported experience over the previous 12 months, yet further data such as sexual IPV and violence during childhood were not collected; therefore, further studies are needed.

Abbreviation

IPV: Intimate partner violence.

Competing interests

The authors have no conflicts of interest to disclose.

Authors' contributions

ML developed the research question and performed the analysis, ML and KMS drafted the manuscript and interpreted the data, ML and ECP participated in the design and planning of the study and ECP is guarantor. All authors have read and approved the final manuscript.

Author details

¹Department of Health Management and Policy, College of Public Health, University of Iowa, 105 River Street, N265 CPHB, 52242 Iowa City, IA, USA.

²Department of Research Affairs, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, 120-752 Seoul, South Korea. ³Department of Preventive Medicine and Public Health, Institute of Health Services Research, Yonsei University College of Medicine, 134 Shinchon-dong, Seodaemun-gu, 120-752 Seoul, South Korea.

Received: 5 November 2013 Accepted: 25 April 2014

Published: 1 May 2014

References

1. Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Watts CH, WHO Multi-country Study on Women's Health and Domestic Violence against Women Study Team: Prevalence of intimate partner violence: findings

- from the WHO multi-country study on women's health and domestic violence. *Lancet* 2006, **368**(9543):1260–1269.
2. Hong JS, Kim SM, Yoshihama M, Byoun SJ: **Wife battering in South Korea: an ecological systems analysis.** *Child Youth Serv Rev* 2010, **32**(12):1623–1630.
 3. Abramsky T, Watts CH, Garcia-Moreno C, Devries K, Kiss L, Ellsberg M, Jansen HA, Heise L: **What factors are associated with recent intimate partner violence? findings from the WHO multi-country study on women's health and domestic violence.** *BMC Public Health* 2011, **11**:109.
 4. Ministry of Gender Equality & Family: **The 2010 Domestic Violence Survey of South Korea.** In *Domestic Violence Survey of South Korea.* Ministry of Gender Equality & Family; 2010. <http://www.mogef.go.kr/>.
 5. Ellsberg M, Jansen HA, Heise L, Watts CH, Garcia-Moreno C, WHO Multi-country Study on Women's Health and Domestic Violence against Women Study Team: **Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study.** *Lancet* 2008, **371**(9619):1165–1172.
 6. Campbell JC: **Health consequences of intimate partner violence.** *Lancet* 2002, **359**(9314):1331–1336.
 7. Ishida K, Stupp P, Melian M, Serbanescu F, Goodwin M: **Exploring the associations between intimate partner violence and women's mental health: evidence from a population-based study in Paraguay.** *Soc Sci Med* 2010, **71**(9):1653–1661.
 8. Suglia SF, Enlow MB, Kullowatz A, Wright RJ: **Maternal intimate partner violence and increased asthma incidence in children: buffering effects of supportive caregiving.** *Arch Pediatr Adolesc Med* 2009, **163**(3):244–250.
 9. Ackerson LK, Subramanian SV: **Intimate partner violence and death among infants and children in India.** *Pediatrics* 2009, **124**(5):e878–e889.
 10. Breiding MJ, Ziembroski JS: **The relationship between intimate partner violence and children's asthma in 10 US states/territories.** *Pediatr Allergy Immunol* 2011, **22**(1 Pt 2):e95–e100.
 11. Biroscak BJ, Smith PK, Post LA: **A practical approach to public health surveillance of violent deaths related to intimate partner relationships.** *Public Health Rep* 2006, **121**(4):393–399.
 12. Martin SL, Macy RJ, Sullivan K, Magee ML: **Pregnancy-associated violent deaths: the role of intimate partner violence.** *Trauma Violence Abuse* 2007, **8**(2):135–148.
 13. Wood SL, Sommers MS: **Consequences of intimate partner violence on child witnesses: a systematic review of the literature.** *J Child Adolesc Psychiatr Nurs* 2011, **24**(4):223–236.
 14. Krug EG, Mercy JA, Dahlberg LL, Zwi AB: **The world report on violence and health.** *Lancet* 2002, **360**(9339):1083–1088.
 15. Djikanovic B, Jansen HA, Otasevic S: **Factors associated with intimate partner violence against women in Serbia: a cross-sectional study.** *J Epidemiol Community Health* 2010, **64**(8):728–735.
 16. Ministry of Gender Equality & Family: **The 2013 Domestic Violence Survey of South Korea.** In *Domestic Violence Survey of South Korea.* Ministry of Gender Equality & Family; 2014. <http://www.mogef.go.kr/>.
 17. Carmo R, Grams A, Magalhaes T: **Men as victims of intimate partner violence.** *J Forensic Leg Med* 2011, **18**(8):355–359.
 18. Gass JD, Stein DJ, Williams DR, Seedat S: **Gender differences in risk for intimate partner violence among South African adults.** *J Interpers Violence* 2011, **26**(14):2764–2789.
 19. Ansara DL, Hindin MJ: **Exploring gender differences in the patterns of intimate partner violence in Canada: a latent class approach.** *J Epidemiol Community Health* 2010, **64**(10):849–854.
 20. Shim WS, Nelson-Becker H: **Korean older intimate partner violence survivors in North America: cultural considerations and practice recommendations.** *J Women Aging* 2009, **21**(3):213–228.
 21. Chung GH, Oswald RF, Hardesty JL: **Enculturation as a condition impacting Korean American physicians' responses to Korean immigrant women suffering intimate partner violence.** *Health Care Women Int* 2009, **30**(1–2):41–63.
 22. Hellmuth JC, Gordon KC, Stuart GL, Moore TM: **Risk factors for intimate partner violence during pregnancy and postpartum.** *Arch Womens Ment Health* 2013, **16**(1):19–27.
 23. Woolhouse H, Gartland D, Hegarty K, Donath S, Brown SJ: **Depressive symptoms and intimate partner violence in the 12 months after childbirth: a prospective pregnancy cohort study.** *BJOG* 2012, **119**(3):315–323.
 24. Ackerson LK, Kawachi I, Barbeau EM, Subramanian SV: **Effects of individual and proximate educational context on intimate partner violence: a population-based study of women in India.** *Am J Public Health* 2008, **98**(3):507–514.
 25. Bensley L, Van Eenwyk J, Wynkoop Simmons K: **Childhood family violence history and women's risk for intimate partner violence and poor health.** *Am J Prev Med* 2003, **25**(1):38–44.
 26. Nguyen DV, Ostergren PO, Krantz G: **Intimate partner violence against women in rural Vietnam—different socio-demographic factors are associated with different forms of violence: need for new intervention guidelines?** *BMC Public Health* 2008, **8**:55.
 27. Tumwesigye NM, Kyomuhendo GB, Greenfield TK, Wanyenze RK: **Problem drinking and physical intimate partner violence against women: evidence from a national survey in Uganda.** *BMC Public Health* 2012, **12**:399.
 28. Lemon SC, Verhoek-Oftedahl W, Donnelly EF: **Preventive healthcare use, smoking, and alcohol use among Rhode Island women experiencing intimate partner violence.** *J Womens Health Gend Based Med* 2002, **11**(6):555–562.
 29. Testa M, Kubiak A, Quigley BM, Houston RJ, Derrick JL, Levitt A, Homish GG, Leonard KE: **Husband and wife alcohol use as independent or interactive predictors of intimate partner violence.** *J Stud Alcohol Drugs* 2012, **73**(2):268–276.
 30. Gil-Gonzalez D, Vives-Cases C, Ruiz MT, Carrasco-Portino M, Alvarez-Dardet C: **Childhood experiences of violence in perpetrators as a risk factor of intimate partner violence: a systematic review.** *J Public Health* 2008, **30**(1):14–22.
 31. Roberts AL, Gilman SE, Fitzmaurice G, Decker MR, Koenen KC: **Witness of intimate partner violence in childhood and perpetration of intimate partner violence in adulthood.** *Epidemiology* 2010, **21**(6):809–818.
 32. Basile KC, Hall JE: **Intimate partner violence perpetration by court-ordered men: distinctions and intersections among physical violence, sexual violence, psychological abuse, and stalking.** *J Interpers Violence* 2011, **26**(2):230–253.
 33. Magdol L, Moffitt TE, Caspi A, Newman DL, Fagan J, Silva PA: **Gender differences in partner violence in a birth cohort of 21-year-olds: bridging the gap between clinical and epidemiological approaches.** *J Consult Clin Psychol* 1997, **65**(1):68–78.
 34. Parish WL, Wang T, Laumann EO, Pan S, Luo Y: **Intimate partner violence in China: national prevalence, risk factors and associated health problems.** *Int Fam Plan Perspect* 2004, **30**(4):174–181.
 35. Chung H: **Variables influencing the adaptation to wife abuse: Based on the double ABCX model.** *J Korean Home Econ Assoc* 1999, **37**:107–122.
 36. Kim JR, Kim KS: **The effects of social support and coping styles on quality of life in abused wives.** *Kor Liv Sci Assoc* 2007, **16**:1–11.
 37. Sullivan CM, Bybee DI: **Reducing violence using community-based advocacy for women with abusive partners.** *J Consult Clin Psychol* 1999, **67**(1):43–53.
 38. Kim M: *The Korean Welfare Panel study and Its Advantages*, Volume 158. Korea Institute for Health and Social Affairs; 2009.
 39. Kim J, Lee J: **Prospective study on the reciprocal relationship between intimate partner violence and depression among women in Korea.** *Soc Sci Med* 2013, **99**:42–48.
 40. Korean National Statistical Office: *National Statistics for Marriage*; 2012.
 41. Vives-Cases C: **Intimate partner violence against women in Spain.** *J Epidemiol Community Health* 2006, **60**(8):652–653.
 42. Yoshihama M, Horrocks J, Kamano S: **Experiences of intimate partner violence and related injuries among women in Yokohama, Japan.** *Am J Public Health* 2007, **97**(2):232–234.
 43. Vung ND, Ostergren PO, Krantz G: **Intimate partner violence against women, health effects and health care seeking in rural Vietnam.** *Eur J Public Health* 2009, **19**(2):178–182.
 44. Hasegawa M, Bessho Y, Hosoya T, Deguchi Y: **Prevalence of intimate partner violence and related factors in a local city in Japan.** *Nihon Koshu Eisei Zasshi* 2005, **52**(5):411–421.
 45. Xu X, Zhu F, O'Campo P, Koenig MA, Mock V, Campbell J: **Prevalence of and risk factors for intimate partner violence in China.** *Am J Public Health* 2005, **95**(1):78–85.
 46. O'Leary KD, Tittle N, Bromet EJ, Gluzman SF: **Descriptive epidemiology of intimate partner aggression in Ukraine.** *Soc Psychiatry Psychiatr Epidemiol* 2008, **43**(8):619–626.
 47. Jewkes R: **Intimate partner violence: causes and prevention.** *Lancet* 2002, **359**(9315):1423–1429.

48. Dunkle KL: Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *Lancet* 2004, **9419**(2):1415–1421.
49. Abrahams N, Jewkes R, Laubscher R, Hoffman M: Intimate partner violence: prevalence and risk factors for men in Cape Town, South Africa. *Violence Vict* 2006, **21**(2):247–264.
50. Dunkle KL, Jewkes RK, Nduna M, Levin J, Jama N, Khuzwayo N, Koss MP, Duvvury N: Perpetration of partner violence and HIV risk behaviour among young men in the rural Eastern Cape, South Africa. *AIDS* 2006, **20**(16):2107–2114.
51. Foran HM, O'Leary KD: Alcohol and intimate partner violence: a meta-analytic review. *Clin Psychol Rev* 2008, **28**(7):1222–1234.
52. Caetano R, Schafer J, Cunradi CB: Alcohol-related intimate partner violence among white, black, and Hispanic couples in the United States. *Alcohol Res Health* 2001, **25**(1):58–65.

doi:10.1186/1471-2458-14-415

Cite this article as: Lee et al.: Gender-specific differences in risk for intimate partner violence in South Korea. *BMC Public Health* 2014 **14**:415.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

