

Research article

Interactions of pathogens and irritant chemicals in land-applied sewage sludges (biosolids)

David L Lewis*^{1,2}, David K Gattie³, Marc E Novak², Susan Sanchez⁴ and Charles Pumphrey⁵

Address: ¹US Environmental Protection Agency, National Exposure Research Laboratory, Athens, GA, USA, ²Departments of Marine Sciences University of Georgia, Athens, GA, USA, ³Biological & Agricultural Engineering University of Georgia, Athens, GA, USA, ⁴Medical Microbiology University of Georgia, Athens, GA, USA and ⁵Prime Care of Sun City, Menifee, CA

E-mail: David L Lewis* - LewisDavel@aol.com; David K Gattie - dgattie@enr.uga.edu; Marc E Novak - uziel@arches.uga.edu; Susan Sanchez - ssanchez@cvm.vet.uga.edu; Charles Pumphrey - schoulten@msn.com

*Corresponding author

Published: 28 June 2002

Received: 4 April 2002

BMC Public Health 2002, 2:11

Accepted: 28 June 2002

This article is available from: <http://www.biomedcentral.com/1471-2458/2/11>

© 2002 Lewis et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Fertilisation of land with processed sewage sludges, which often contain low levels of pathogens, endotoxins, and trace amounts of industrial and household chemicals, has become common practice in Western Europe, the US, and Canada. Local governments, however, are increasingly restricting or banning the practice in response to residents reporting adverse health effects. These self-reported illnesses have not been studied and methods for assessing exposures of residential communities to contaminants from processed sewage sludges need to be developed.

Methods: To describe and document adverse effects reported by residents, 48 individuals at ten sites in the US and Canada were questioned about their environmental exposures and symptoms. Information was obtained on five additional cases where an outbreak of staphylococcal infections occurred near a land application site in Robesonia, PA. Medical records were reviewed in cases involving hospitalisation or other medical treatment. Since most complaints were associated with airborne contaminants, an air dispersion model was used as a means for potentially ruling out exposure to sludge as the cause of adverse effects.

Results: Affected residents lived within approximately 1 km of land application sites and generally complained of irritation (e.g., skin rashes and burning of the eyes, throat, and lungs) after exposure to winds blowing from treated fields. A prevalence of *Staphylococcus aureus* infections of the skin and respiratory tract was found. Approximately 1 in 4 of 54 individuals were infected, including 2 mortalities (septicaemia, pneumonia). This result was consistent with the prevalence of *S. aureus* infections accompanying diaper rashes in which the organism, which is commonly found in the lower human colon, tends to invade irritated or inflamed tissue.

Conclusions: When assessing public health risks from applying sewage sludges in residential areas, potential interactions of chemical contaminants with low levels of pathogens should be considered. An increased risk of infection may occur when allergic and non-allergic reactions to endotoxins and other chemical components irritate skin and mucus membranes and thereby compromise normal barriers to infection.

Background

Class B biosolids, the most common form of processed sewage sludges used in agriculture, are treated to reduce levels of pathogens by various processes including anaerobic digestion and pH elevation (lime stabilisation). The finished product may contain a wide variety of contaminants with a potential for adverse health effects [1]. This includes enteric bacteria, viruses, endotoxins, inorganic and organic sulfides, volatile fatty acids, alkyl amines, and ammonia [2,3]. No formal pathogens risk assessment has been carried out for public exposure to Class B biosolids [4]; and, complaints of sludge-related illnesses reported by residents have not been previously studied.

Retrospective studies have been done on workers exposed to sewage sludges who have reported illnesses. Gregersen et al [5], for example, investigated five workers who experienced fever and flu-like symptoms after repairing a decanter used for concentrating sludge at a sewage treatment plant. High concentrations of *Legionella pneumophila* (serogroup 1) found in the sludge along with positive antibody titers in all five workers indicated that they developed Pontiac fever from contact with sludge. Similarly, NIOSH investigated five workers complaining of headaches, stomach cramps, and diarrhoea at a sludge processing operation [6]. Investigators isolated a variety of enteric, opportunistic pathogens from bulk sludge samples and airborne particulates, including species of the genera *Mycobacterium*, *Pseudomonas*, and *Staphylococcus*. Based on the nature and timing of their symptoms, investigators concluded that ingestion or inhalation of pathogens in sewage sludge probably caused the illnesses.

Dorn et al, on the other hand, found no elevated risks of infection in the only epidemiological study done with residents exposed to sewage sludges [7]. The authors of this prospective study, however, did not determine whether any pathogens were present in the material. Also, because of the low application rates used in the study, they cautioned against extrapolating their data to other sites.

Our retrospective investigation was designed to use a survey for describing adverse effects reported by residents, develop an approach for assessing their environmental exposures, and evaluate whether any infections associated with exposure to sewage sludges were prevalent within the group.

Methods

Surveying residents

To indicate the general symptomatology, we surveyed 48 affected residents at ten land application sites in Alabama, California, Florida, New Hampshire, Ohio, Ontario, Pennsylvania, and Texas (Table 1). Questionnaires were patterned after a previous epidemiological study in Ohio

and based on symptoms associated with exposure to sewage sludges [6,7]. More specifically, residents were queried about their symptoms and how much time elapsed between exposures and onset. Individuals who were previously questioned by environmental activists were excluded from the survey and medical records were obtained in cases involving death or serious illnesses (those requiring medical attention).

Using chi-squared analyses, proportions of individuals reporting a particular symptom at each site were compared with the combined reports for that symptom at all sites. *P* indicated how consistently a symptom was reported from site to site, that is, the degree to which numbers of individuals reporting a particular symptom at a given site could be expected based on the frequency of that symptom for all sites combined. Low *P* values were indicative of symptoms that were largely independent of confounding health-related and environmental factors (e.g., previous medical history, demographics, sources of biosolids, and environmental conditions).

This survey, which did not include an unexposed control group, was primarily intended to describe and document self-reported illnesses. It also provided guidance on which exposure routes (e.g., contaminated drinking water, dermal contact, or inhalation of aerosols) should be the focus of the research. Exposure, on the other hand, was evaluated using a dose-response curve in a separate part of the study concerning environmental assessments.

Assessing environmental conditions

Sludges used at all but one agricultural site (Ontario), according to county records, were limed or applied to alkaline soils then exposed to weathering. Whether sludges used for mining reclamation in Osceola Mills were lime stabilised, however, was disputed. Residents reported that the area was limed after sludging.

County records indicated that biosolids-related complaints for individual patients described in this study were concurrent with land application of Class B biosolids. In the case of one family (Residents 19–22), however, records indicated that dairy wastes, rather than biosolids, had been applied. Unconfirmed reports that dairy wastes had been mixed with Class B biosolids prior to application at this site, however, precluded the authors from ruling out biosolids as a possible cause of the symptoms. According to county officials, soil in this area is naturally alkaline such that irritation similar to that observed with exposure to lime-stabilised sludge may occur.

The NH site was used to develop an approach for assessing exposures to windborne contaminants. At the site, approximately 132 metric tons (dry wt) of Class B biosolids

Table 1: Survey of residents complaining of adverse health effects.

Res.	Site, distance away	(km)	Age/sex	Smoker	Symptoms ^a
1	Grand Bay, AL	0.4	28♀	N	Be, Bt, Bl, Cn, Db, N
2	Grand Bay, AL	0.4	4♀	N	Be, Bt, Bl, Cn, Db, H, N
3	Grand Bay, AL	0.8	48♀	N	C, Be, Bt, Cn, Db, R, N, D, Ft
4	Grand Bay, AL	0.8	19♀	N	C, Bt, Bl, Cn, Db, S, R, H, N, D, Ft
5	Grand Bay, AL	0.4	71♂	N	C, Cn, Db, H
6	Grand Bay, AL	<0.1	39♀	Y	C, Cn, Db, R, H, N
7	Grand Bay, AL	<0.1	19♂	N	C, Be, Bl, Cn, Db, R
8	Grand Bay, AL	<0.1	51♀	N	Be, Bt, Bl, Cn, Db, D
9	Grand Bay, AL	0.2	45♂	N	C, Be, Bt, Bl, Cn, H, N
10	Menifee, CA	0.4	42♂	N	C, Cn, Db, Si, N, F, D, Ft
11	Menifee, CA	0.1	37♀	N	Be, Bt, Cn, R, H, Si, N, Nb
12	Menifee, CA	0.1	4♂	N	Bt, Cn, Db, Si, S, P
13	Menifee, CA	0.8	40♀	N	C, Db, R, H, N, D, Si
14	Menifee, CA	0.4	52♀	Y	Cn, Db, H, N, Nb, Ft
15	Menifee, CA	0.4	7♀	N	C, Bl, Cn, Db, Fv, F
16	Menifee, CA	0.4	28♂	N	C, Cn, Db, H, S
17	Menifee, CA	0.2	29♀	N	C, Be, Bt, Cn, Db, R, H, Si, Ft
18	Menifee, CA	0.2	5♀	N	C, Be, Cn, Db, S, P, R, H, Fv, F, Ft
19	Menifee, CA	<0.1	43♀	Y	Be, Bt, Cn, Db, R, H, N, Fv, Nb, Ft
20	Menifee, CA	<0.1	10♂	N	Be, Bt, Cn, Db, S, H, N, Fv, Nb, Ft
21	Menifee, CA	<0.1	68♀	Y	Cn, Db, R, N, Fv, F, Ft
22	Menifee, CA	<0.1	46♂	Y	Cn, Db, R, D, F, Ft
23	Menifee, CA	0.4	44♀	Y	C, Be, Bt, Bl, Cn, H, F, Ft
24	Winchester, CA	0.8	69♂	Y	C, Be, Bt, Bl, Db, P
25	Winchester, CA	0.8	46♀	Y	C, Be, Bt, Bl, Db, H, N, Ft
26	Arcadia, FL	<0.2	38♀	Y	Be, Bt, Bl
27	Arcadia, FL	<0.2	9♀	N	Db, N, D
28	Arcadia, FL	<0.2	8♂	N	C, Cn, Db
29	Greenland, NH	0.2	53♀	N	C, Cn, Db, H, N, D
30	Greenland, NH	0.2	70♀	N	C, Be, Bt, Cn, Nb
31	Greenland, NH	<0.1	32♀	Y	Be, Bt, Bl, Cn, Db, H, N, F, Nb
32	Greenland, NH	<0.1	9♀	N	C, Be, Cn, Db, N, D, F, Nb
33	Greenland, NH	<0.1	14♀	N	Be, Bt, Cn, Db, N, D, F, Ft
34	Greenland, NH	<0.2	48♀	Y	C, Be, Bt, Cn, Db, H, N, F, Ft
35	Greenland, NH	<0.2	26♂	N	C, Cn, H, F
36	Waynesville, OH	0.2	60♀	N	C, Be, Bt, Bl, Db, S, R
37	Waynesville, OH	0.1	78♂	N	C, Db, S
38	Cedarville, ON	0.2	51♂	N	Be, Bt, Bl
39	Robesonia, PA	0.2	15♂	N	C, Cn, Db, P, H, Fv, F, Si, Ft
40	Robesonia, PA	0.2	17♂	N	C, Bt, Cn, H, S, Si,
41	Robesonia, PA	0.2	46♂	N	C, Bt, Si
42	Osceola Mills, PA	0.8	11♂	N	Be, Bt, Cn, H, Si, N, F, Ft
43	Guy, TX	1.6	55♂	N	Bt, Bl, Db
44	Guy, TX	0.4	2♂	N	C, Cn, Db, Si
45	Guy, TX	0.4	67♀	N	C, Be, Bt, Bl
46	Guy, TX	0.4	9♂	N	Be, Bl, Cn, N
47	Guy, TX	0.4	6♀	N	C, Cn, Be, Bt, Bl, N
48	Guy, TX	0.4	68♂	N	C, Be, Cn, N, S, Ft

Residents complaining of adverse health effects from exposure to land-applied sewage sludge were surveyed at ten sites in North America to describe and document self-reported illnesses.

^a (C) cough; (Be) burning eyes/(Bt)throat/(Bl)lungs; (H) headache; (Cn) congestion; (Db) difficulty breathing; (F) flu-like; (Fv) fever; (N) nausea/vomiting (D) diarrhoea; (S) sinusitis; (Si) staphylococcal infection (skin, oral cavity); (P) pneumonia; (R) skin rash; (Nb) nosebleed; (Ft) fatigue

containing 12 metric tons of lime and 318 kg of Stockhausen 644BC cationic polymer (Stockhausen, Inc. Greensboro, NC) were applied to a 10-ha field in late October 1995. The field was located at the end of a paved road with eight houses (A-H) and 28 residents.

Historical records of symptoms experienced in the neighbourhood were kept by one of the residents. Records began approximately 2 weeks after sludges were applied and continued for approximately two years (1995–96). Descriptions of symptoms included coughing, burning eyes, burning throat, congestion, and difficulty in breathing.

To evaluate wind conditions at the site at the time sewage sludge was applied, we used a computer program (WRPLOT Version 3.5, Lakes Environmental Corp., Waterloo, Ontario), topographic information, and historical weather data from Pease Air Force Base located approximately 5 km from the site. Exposure times (h), which were based on duration of wind passing directly from the field toward each house, were determined from wind frequency data averaged for three 8-day periods after sludges were applied (between October 31 – November 23, 1995). Distances (m) each household resided from the nearest edge of the treated field were 23 (A, B); 77 (C), 170 (D); 130 (E); 280 (G); and 320 (H).

Concentrations of air pollutants to which residents were exposed, relative to concentrations over the treated field, were predicted using an Industrial Source Complex (ISC3) Dispersion Model [8]. This prediction was not specific for any pollutant. Instead, it was used as a means for possibly ruling out the effects of ammonia, organic amines, or other gaseous or volatile pollutants. Should airborne chemical contaminants be present only at a fraction of a percent of the concentrations present over the treated field, the result would argue against airborne chemical pollutants as a cause of adverse effects. If, on the other hand, pollutants are sufficiently concentrated to cause burning eyes and other symptoms of irritation at the application site, these same effects may occur among residents in areas where pollutant concentrations are not highly diluted.

The ISC3 air dispersion model is designed to predict concentrations of pollutants off site based on emission rates of the pollutants at their sources and local meteorological conditions. This approach is unreliable when emission rates cannot be accurately determined. Microbiological process rates, which largely control emission rates of airborne pollutants produced by sewage sludges, are highly unpredictable. To circumvent this problem, the model was used only to predict the relative concentrations of air pollutants between the treated field and potentially affected residences. This approach is based on the dispersion of

air pollutants as winds carry them over land surfaces, irrespective of their emission rates at the source.

Results

Residents survey

Demographic data for the 48 residents surveyed, along with their symptoms, are listed in Table 1. The average age of the group was 35 years; 56% were female; 23% were smokers. Residents who complained of adverse effects lived within approximately 1 km of the treated fields. Symptoms were primarily associated with sites where freshly applied, lime-stabilised sewage sludges remained on the surface and were exposed to wind erosion.

The primary complaints (a fourth or more reporting the symptom) within 1 h of exposure were coughing (63%, 8 sites), burning throat (56%, 10 sites), burning eyes (56%, 9 sites), and headaches (46%, 6 sites). Within 24 h, residents developed nasal and chest congestion (77%, 7 sites), difficulty in breathing (71%, 8 sites), nausea/vomiting (46%, 7 sites), fatigue (33%, 7 sites), and flu-like symptoms (29%, 5 sites). In some cases, symptoms of chemical irritation (e.g., burning eyes, burning throat) recurred for over a year after applications ceased.

Proportions of individuals reporting coughing and burning throats exhibited the greatest consistency from site to site (P 0.02, 0.03, respectively). Half of the residents reported bacterial, viral or fungal infections within 1 month of exposure (50 %, 8 sites). Most residents also reported symptoms indicating exposure to infectious micro-organisms or microbial products (endotoxins). These included flu-like symptoms (29%, 5 sites), nausea/vomiting (46%, 7 sites), diarrhoea (21%, 4 sites), rashes (25%, 3 sites), and sinusitis (19%, 5 sites).

The most common infections identified by medical records were caused by *S. aureus* (14 residents at 3 sites, including the additional patients identified in Robesonia). Records indicated that one infection (*S. lugdunensis*, Resident 44) was probably hospital-acquired. Affected residents in Greenland, NH recalled that respiratory infections experienced by individuals in several households treated by the same physician responded to erythromycin, an antibiotic primarily used for treating gram positive bacterial infections.

In Menifee, CA, two family members complaining of chemical irritation developed different staphylococcal infections. Resident 17 experienced recurrent *S. aureus* infections of the nasal passages and oral cavity. The spouse was hospitalised for an invasive *S. epidermitis* infection that developed approximately two days after an undressed surgical incision became contaminated with sand blowing from an adjacent field treated with Class B biosolids.




Figure 1
Exposure to gases and volatile chemicals Concentrations of gases and volatile pollutants at Household C relative to concentrations 2 m directly above the biosolids-treated field.

Mild chronic maxillary sinusitis in one child (Resident 18) living in this household deteriorated into recurrent (unidentified) bacterial infections. The patient responded poorly to oral and IV antibiotic therapy, tonsillectomy, and removal of the adenoids. Rigorous antibiotic therapy, including intramuscular injections of rocephrine, also yielded unsatisfactory results.

County records showed that Class B biosolids were applied to an adjacent field at the time the above-mentioned illnesses developed. The paediatrician treating patients in the area (author C. Pumphrey) noted that children living near the land application sites appeared to develop abnormally persistent sinusitis.

Environmental assessment

Samples of sludges collected at the NH site, which were frozen since 1995, retained a strong faecal odour. Microbiological assays showed that they contained $(3.2 \pm 3.5) \times 10^5$ colony-forming units of bacteria per gram but almost no culturable fungi. Predominant culturable bacteria were identified as common gram-negative environmental isolates and included *Alcaligenes faecalis* and *Brevundimonas diminuta*, which occasionally serve as opportunistic human pathogens.

Exposure times (h) for each household when initial symptoms of chemical irritation were reported (Oct 31 – Nov 23) were: 28 ± 25 (A); 48 ± 17 (B, C), 21 ± 12 (D), 88 ± 17 (E); and 13 ± 08 (G, H) (mean \pm standard deviation, $n = 3$). The air-dispersion model indicated that adverse effects from exposure to airborne pollutants from the land application site could not be ruled out as a cause of ad-

verse health effects. Air pollutants at Household C, for example, were frequently present at concentrations of approximately 5 – 40% compared with concentrations 2 m directly above the treated field (Fig. 1).

Numbers of individuals who reported adverse effects compared with the total numbers of individuals per household, respectively, were: 2, 2 (A); 2, 2 (B); 5, 5 (C), 2, 3 (D); 4, 4 (E); 1, 3 (G); and 1, 5 (H). Data from one household (F) were unavailable. Based on a least-squares analysis, proportions of individuals with symptoms increased linearly from 40 to 80 h (r^2 0.98) with time exposed to wind blowing from the field; all occupants in households with exposure ≥ 80 h reported symptoms (Fig. 2). Proportions of individuals with symptoms also decreased linearly with distance from the field from 130 to 320 m (r^2 0.95); all occupants in households living ≤ 130 m from the field reported symptoms.

Prevalence of *S. aureus* infections

Two cases involving mortality from *S. aureus* occurred with individuals who directly contacted freshly applied biosolids. In the first case, an 11-year-old male in Osceola Mills, PA with an unremarkable medical history died of staphylococcal septicaemia. The patient developed a sore throat, headaches, and furuncles on one leg and one arm within several days after riding a motorbike through sewage sludges applied nearby for mine reclamation purposes. A primary care physician prescribed antibiotics and the patient was admitted to the hospital the following day with difficulty in breathing and high fever.

S. aureus was isolated from skin lesions and IV antibiotics were administered. The patient developed septicaemia and expired six days after contacting the biosolids. Mine workers in the same area requested a health hazard evaluation after experiencing similar respiratory and skin irritation symptoms that they attributed to biosolids exposure.

The second mortality was associated with an outbreak of *S. aureus* in Robeson, PA. According to state records, Class B, lime-stabilised sewage sludges from a local wastewater treatment facility were applied to ten fields comprising a total of 300 acres. Applications began in 1988 and continued through December 1995. Sewage sludges were applied five times a week at a rate of approximately 1,300 tons (dry wt) per year.

Of nine individuals living in or frequenting the house where the outbreak occurred, eight developed *S. aureus* infections over a five-year period beginning in February 1993 (Table 2). The house was located across a paved road approximately 200 m from the treated field. Three individuals (Patients 1, 7, 8) were granted hunting rights on




Figure 2
Dose-response relationship Proportions of individuals experiencing symptoms in households versus exposure to gaseous and volatile pollutants from the biosolids-treated field.

the farmland and frequently traversed treated areas during the application period.

Eight additional individuals lived in four other houses that were also located in close proximity to the field (<50 m). These individuals complained of odours, inundation with houseflies, and respiratory symptoms. They associated these problems with application of biosolids; however, they could not recall having any infections.

Residents reported that feral cats occupying a barn near the treated field had to be destroyed after developing multiple boils and weeping sores after sewage sludge applications began. Similarly, residents at the New Hampshire site reported that one individual and numerous pets (dogs, cats) developed boils after sewage sludges were applied. Pets may have served as a reservoir of staphylococcal infections in these neighbourhoods and, in some cases, transmitted their infections to humans.

Medical records were available for three of the four infected residents: Patients 1, 3, 8. Patient 1, a 17 year-old male with a history of excellent health, was treated for a furuncle of the knee in February 1993 then succumbed to *S. aureus* pneumonia in March 1995 after contracting a rotavirus infection and viral pneumonia. Each of four relatives who frequented the house (≥ 2 d/wk) was also infected with *S. aureus*. The first of these (Patient 3) developed furuncles concurrently with dermatophytic infections (identified as tinea cruris/tinea corpora). All but two individuals (Patients 6, 7) sought medical attention. These patients were self-treated using antibiotics prescribed for other family members. The last *S. aureus* infec-

tion at Robesonia (Patient 8) occurred approximately two years after biosolids applications ceased.

Overall, the prevalence of *S. aureus* infections in this study was approximately twenty-five times higher than infections among hospitalised patients, a recognised risk group for *S. aureus*[9].

Discussion

Chemical irritants of concern with biosolids include endotoxins, lime, ammonia, and alkyl amines. These contaminants may cause allergic and non-allergic reactions that could contribute to broncho-obstructive and inflammatory responses [3]. The senior author and two accompanying individuals, for example, experienced coughing, burning eyes, burning throat, headaches, congestion and difficulty in breathing within 1 h while conducting a site visit in Menifee, CA. The visit occurred during a time when residents were reporting these same symptoms. Although high winds were blowing sand from the treated fields, no odour was evident at that time. This experience argued against attributing the symptoms to odour-related psychosomatic responses, which are thought to play an important role in self-reported illnesses associated with human and animal waste-treatment operations [2].

The lower human colon is the primary reservoir for *S. aureus*; therefore, the organism is a common sewage contaminant. It is known to survive desiccation for days to weeks and can be aeri ally transmitted [10]. It tends to invade irritated tissue and infections are a common complication of diaper rash after prolonged contact with urine and faeces [11,12]. Although it can be part of the natural skin and nasal microflora, risks of infection vary widely among different strains and increase with exposure to virulent strains from other sources. It is a leading cause of nosocomial pneumonia and surgical site and bloodstream infections [9]; however, the EPA considered it to be a low risk to public health with land applied sewage sludges [13].

In the case of recurrent *S. aureus* infections in Robesonia (Table 1), it was evident that a persistent source of *S. aureus* was present. It was unclear, however, whether the strain(s) responsible for the infections had a common environmental source lasting for several years or if certain individuals or pets became persistent carriers and continued to expose others even after land application ceased. Residents reported that tilling the heavily treated soil still, seven years later, draws large swarms of houseflies to the area.

A lack of any documented risk of staphylococcal infections among workers processing and handling wastewater and sewage sludges may reflect the fact that workers are more immunocompetent than the general population, or that adequate epidemiological studies have not been

Table 2: Outbreak of *S. aureus* in Robesonia, PA.

Patient	Onset of disease	Age/sex	Occupancy	Presentation
1	Feb 93	15♂	resident	furuncle ^a
2	Jul/Aug 93	87♀	resident	furuncle ^b
3	Jul 94	40♂	visitor	furuncles ^{a,c}
4	Aug 94	66♀	visitor	foliculitis
1	Mar 95	17♂	resident	pneumonia
5	Jun/Jul 95	37♀	visitor	furuncles ^d
6	Aug 95	7♀	visitor	furuncle ^b
7	Dec 95	16♂	resident	furuncle ^c
8	Sep/Oct 97	46♂	resident	furuncles ^{e,f}

Recurrent *Staphylococcus aureus* infections developed in a Robesonia, PA household over a 5-yr period after Class B biosolids were applied several times a week on a yearly basis to a nearby farm. Patients 1, 7, and 8 correspond with Residents 39, 40, and 41, respectively (Table 1). Patient 1 developed a *S. aureus* furuncle on one knee then, two years later, succumbed to *S. aureus* pneumonia.

^a knee; ^b leg; ^c thigh; ^d nose, arm, abdomen, buttocks; ^e neck, chin; ^f chest

done. It may also reflect differences in exposure routes; or, it may suggest that incubation in the field can lead to increased concentrations or virulence of the organism.

Risks of infection with *S. aureus* may also depend on environmental processes that enhance the ability of sewage sludges to irritate the skin and respiratory tract. As gram negative bacteria die off after sludge application, endotoxin concentrations (a potential source of bronco-pulmonary irritation) increase. As sludges dry in the field, lime and other irritant chemicals become concentrated and, therefore, may be more irritating upon dermal contact and inhalation.

Conclusions

Federal guidelines advise wearing protective gear and exercising good personal hygiene when handling Class B biosolids or exposed to the aerosols [6,14]. The nature and timing of symptoms reported by residents suggest that steps should also be taken to protect the public from exposure to airborne contaminants from land application areas. Air-dispersion models, such as the one used in this study, could provide guidance in selecting application sites where exposure of residents would be minimal.

The Osceola Mills case, Robesonia outbreak, and the occurrence of staphylococcal infections among residents at other land application sites suggest that risks land application practices pose from these pathogens may be higher than expected. Chemical contaminants in sludges, which irritate the skin and mucous membranes, may be responsible for an increased host susceptibility.

Further research is needed to address illnesses among residents exposed to Class B biosolids, especially airborne dusts, and the role irritant chemicals may play in risks posed by low levels of pathogens.

Competing interests

None declared.

Authors' contributions

DL was responsible for the study's concept and design, supervised the study, participated in the analysis and interpretation of data, drafted the manuscript, and participated in the critical revision of the manuscript. DG supervised and participated in the data collection using air dispersion modelling, participated in the analysis and interpretation of that and other data, contributed to the technical support of the project, and participated in the critical revision of the manuscript. MN participated in the dispersion modelling, conducted the survey of self-reported symptoms, contributed to the computer support and data management of the project, and participated in the revision of the manuscript. SS supervised the isolation and identification of microbial isolates and contributed to the analysis and interpretation of that data. CP treated children living near land application sites in Menifee, CA and participated in the acquisition and interpretation of medical data.

Acknowledgements

Support in identifying microbial isolates was provided by Pamela Currin, Eric Wommack (DNA sequencing of *Brevundimonas diminuta*), G. E. Michaels, Scott Russell, and Lynda Jones.

Disclaimer This work was conducted in part under an Intergovernmental Personnel Act assignment between the U.S. Environmental Protection Agency (EPA or Agency) and the University of Georgia. Although this paper has been subjected to an Agency review process and approved for publication, the views expressed are those of the authors and do not necessarily reflect the views or policies of the EPA. Mention of trade names or commercial products does not constitute an endorsement or recommendation for use.

Technical assistance with air dispersion modeling was provided by Dennis G. Atkinson.

References

- Lewis D, Gattie D: **Pathogen risks from applying sewage sludge to land.** *Environ Sci Technol* 2002, **36**(13):286A-293A
- Schiffman SS, Walker JM, Dalton P, Lorig TS, Raymer JH, Shusterman D, Williams CM: **Potential health effects of odor from animal operations, wastewater treatment facilities, and recycling byproducts.** *J Agromed* 2000, **7**(1):1-80
- Norn S, Clementsen P, Kristensen KS, Skov PS, Bisgaard H, Gravensten S: **Examination of mechanisms responsible for organic dust-related diseases – mediator release induced by microorganisms – a review.** *Indoor Air* 1994, **4**(4):217-222
- Committee on the Use of Treated Municipal Wastewater Effluent and Sludge in the Production of Crops for Human Consumption: **Use of reclaimed water and sludge in food crop production.** *Water Science and Technology Board, National Academy of Sciences Press, Washington* 1996
- Gregersen PT, Grunnet K, Uldum SA, Anderson BH, Madson H: **Pontiac fever at a sewage treatment plant in the food industry.** *Scand. J. Work Environ. Health* 1999, **25**(3):291-295

6. **Centers for Disease Control & Prevention, National Institute for Occupational Health & Safety (NIOSH).** *Health Hazard Evaluation Report No. 98-0118-2748.*
7. Dorn RC, Reddy CS, Lamphere DN, Gaeuman JV, Lanese R: **Municipal sewage sludge application on Ohio farms: health effects.** *Environ Res* 1985, **38**:332-359
8. **Industrial Source Complex (ISC3) Dispersion Model. ISCST3 Version 99155.** *Office of Air Quality Planning and Emissions, Monitoring, and Analysis Standards Division. EPA-454/B-95-0032. US Environmental Protection Agency: Research Triangle Park, NC. 1995*
9. Rubin RJ, Harrington CA, Poon A, Dietrich K, Greene JA, Moiduddin A: **The economic impact of *Staphylococcus aureus* in New York City hospitals.** *Emerg Infect Dis* 1999, **5**:9-17
10. Sherertz RJ, Bassetti S, Bassetti-Wyss B: **"Cloud" health-care workers.** *Emerg Infect Dis* 2001, **7**:241-244
11. Leyden JJ, Kligman AM: **The role of microorganisms in diaper dermatitis.** *Arch Dermatol* 1978, **114**(1):56-59
12. Brook I: **Microbiology of secondarily infected diaper dermatitis.** *Int J Dermatol* 1992, **31**:700-702
13. U.S. Environmental Protection Agency: **Preliminary risk assessment for bacteria in municipal sewage sludge applied to land.** *EPA/600/6-91/006* 1991
14. US Department of Health and Human Services: **Workers exposed to Class B biosolids during and after field application.** *DHHS (NIOSH) Publication No. 158* 2000

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/2/11/preview>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>



BioMedcentral.com

editorial@biomedcentral.com