

RESEARCH ARTICLE

Open Access



Individual and family level factors associated with physical and mental health-related quality of life among people living with HIV in rural China

Yongkang Xiao¹, Chunqing Lin², Li Li² and Guoping Ji^{3*} 

Abstract

Background: One of the major challenges faced by people living with HIV (PLH) is the compromised quality of life due to the negative impact of HIV on their health. HIV/AIDS control effort should go beyond laboratory and lay more emphasis on improving the health-related quality of life (HRQoL) for PLH. The objective of this study is to evaluate the physical and mental HRQoL of PLH in rural China, and explore the relationship between HRQoL and individual- and family-level factors.

Methods: A cross-sectional study was conducted among 522 PLH in Anhui, China. Participant's sociodemographic characteristics, family status, and HIV-related factors were collected. Physical health summary score (PHS) and mental health summary score (MHS) of quality of life were measured. Multiple linear regressions were conducted to estimate the association of the individual- and family-level factors with MHS and PHS.

Results: Male were more likely to report a higher level of PHS and MHS than female ($\beta = 0.123$, $P = 0.009$; $\beta = 0.150$, $P = 0.002$). Age was significantly negatively associated with the PHS ($\beta = -0.232$, $P < 0.001$) when other variables were controlled. Family size remained negatively correlated with PHS ($\beta = -0.105$, $P = 0.021$). Family annual income was significantly positively associated with PHS and MHS ($\beta = 0.126$, $P = 0.003$; $\beta = 0.135$, $P = 0.002$).

Conclusions: Future intervention should be carefully tailored to the specific needs of sub-populations (such as female and older PLH) considering their physical and mental HRQoL conditions. More attention and care should be provided to PLH with left-behind children in the family.

Keywords: HIV, Physical health, Mental health, Health-related quality of life, China

Background

The availability of antiretroviral therapy (ART) and advancement of primary care system have prolonged the lifespan of people living with HIV (PLH), and HIV has become a chronic illness [1]. However, one of the significant challenges faced by PLH is the compromised quality of life due to the negative impact of HIV on their physical health and mental health [2, 3]. Literature has suggested that suboptimal quality of life may in turn have adverse influences on the patients' ART retention,

medication adherence, and treatment outcomes [4, 5]. Therefore, HIV/AIDS control effort should go beyond objective measures such as CD4 count and viral load, and lay more emphasis on improving the health-related quality of life (HRQoL) of PLH.

Previous studies have identified multilevel factors associated with HRQoL of PLH. Social-demographic characteristics, such as gender, age, and education, were all found to be correlated with HRQoL [6]. Older age was independently associated with lower physical HRQoL for both genders [7]. Disease-related factors associated with HRQoL included stage and time since diagnosis, current CD4 count/viral load, and severity of HIV/AIDS-related symptoms [8, 9], and being on ART was associated with

* Correspondence: ji990066@163.com

³Anhui Provincial Center for Woman and Child Health, 38 Yonghong Road, Hefei 230032, China

Full list of author information is available at the end of the article



better physical and mental health [10]. Psychological factors, such as depression [11] and social support [6, 9], were also found to be correlated with lower HRQoL of PLH. In addition, family factors play an essential role in the HRQoL of PLH in Asia because of the cultural norms with strong family values [12]. Family income [13] and family support [14] were the associating factors identified in previous studies, as family presents a source of support for the individual with HIV/AIDS, contributing to their physical and mental well-being [5]. Therefore, the study of PLH's HRQoL should focus more on family-related factors.

In China, there were 501,000 reported cases of PLH by the end of 2014, most of whom lived in rural areas [15]. However, few studies reported the HRQoL of PLH in rural China. Existing studies in China mainly focus on the relationship between HRQoL and the PLH's individual level factors, such as age [16, 17], employment status [18], income [19], social support [19–23], CD4 count [16], depression [20] and stigma [23–25]. Recently, mental HRQoL was found to be negatively associated with unprotected anal intercourse among Chinese men who have sex with men (MSM) [26]. Studies examining the association between HRQoL and family-related factors were scant in China. A few exceptions included Hidru and colleagues' (2016) study among MSM with HIV/AIDS, which reported that family income has an influence on PLH's HRQoL [27]. A study conducted in Nanning, China reported that the lower HRQoL among older Chinese PLH was due to their limited social networks and heavy dependence on family to cope with HIV infection and ageism [28]. Wu and colleagues (2015) studied HIV clinic attendees and reported that family stigma was inversely correlated with most of the HRQoL domains [23]. On the contrary, family support was found to contribute to PLH's HRQoL [29, 30]. The study aimed to evaluate the physical and mental HRQoL of PLH in rural China, and explore the relationship between HRQoL and individual- and family-level factors. The findings will provide insights to develop strategies to improve the HRQoL for PLH in rural China.

Methods

Study sample

The study used baseline data from a randomized controlled trial for HIV-affected families in Anhui Province, China. Though the number of new HIV cases through sexual infection has rapidly increased in recent years, the proportion of PLH who were caused by contaminated commercial plasma donations in the last century was still dominant among the existing total PLH population in certain areas, especially in rural areas [31–34]. A total of 522 PLH were recruited in this study based on the following criteria: (1) being aged 18 or over;

(2) being local residents; and (3) having a sero-negative family member and a child in the family.

Data collection

Data were collected between October 2011 and March 2013 from 32 villages in four counties (Lixin, Funan, Yinzhou, and Linqian) of Anhui. The four counties were similar in terms of demographics and HIV epidemiological profiles. Detailed study procedure was documented elsewhere [35, 36]. This study used MOS-HIV questionnaires as the main instrument to measure the quality of life, the questionnaire has been widely used in HIV-related studies and previously published in Wu's article [37, 38].

Written informed consent was obtained from the participants. The study was reviewed and approved by the institutional review boards of the Anhui Provincial Center for Disease Control and Prevention and the University of California, Los Angeles.

Measures

Quality of life

Quality of life was assessed with the Medical Outcomes Study HIV Health Survey (MOS-HIV) [39], which has 11 subscales. With these subscales, the physical health summary score (PHS) and mental health summary score (MHS) were generated. According to the unified algorithm, the physical functioning, pain and role functioning subscales contributed most to PHS, the mental health, health distress, quality of life, and cognitive functioning subscales contributed most to MHS, and the vitality, general health, and social functioning subscales contributed partially to both summary scores. Scores for these subscales and two aggregate measures (PHS and MHS) were calculated following the instruction and ranged from 0 (worst) to 100 (best) [37]. The MOS-HIV has high internal consistency and test-retest reliability [40] and has been tested among Chinese population [41]. The internal consistency reliability of the 11 subscales in our sample ranged from 0.75 to 0.91.

Sociodemographic characteristics

The PLH participants' sociodemographic characteristics, including age, gender, marital status, and education, were collected.

Family factors

Three family factors were measured based on the following questions: (1) Family size: "How many people are there in your family, including those who live with you?", (2) Family annual income: "What is your total yearly family income (yuan)?", and (3) Spouse's HIV sero-status; combination of three questions: "What is the health status of your spouse? (Ill/Dead/Healthy)"

“What is his/her illness? (HIV/AIDS or Other- Specify)”
 “What was the cause of death?”

HIV- related factors

Two HIV-related factors were assessed based on the following questions: (1) Time of HIV diagnosis: “When did you first know your HIV infection diagnosed by a doctor? (Specify month/year)”; (2) ART status, “Are you currently on ARV treatment?” (Yes or No).

Statistical analysis

All analyses were performed using SAS 9.4 software (SAS Institute Inc., Cary, NC, USA). First, we calculated descriptive statistics and frequencies (i.e., frequency, mean, and standard deviation) to describe the participants’ demographic characteristics, family and HIV-related conditions, and MHS and PHS scores. Secondly, we compared MHS and PHS on every variable of interest among the participants using T-test or ANOVA to determine whether MHS and PHS were different among different subgroups. Finally, multiple linear regression models were fitted to estimate the association of factors with the MHS and PHS adjusting for control variables. MHS and PHS were treated as continuous outcome variables in the multiple linear regression models.

Results

Sample characteristics

A total of 522 participants were included in this study. The average age was 48.57 years (range: 22–73 years). Approximately 45% were male. The majority of participants (81.4%) were married or cohabiting, and 40.2% had no education. Approximately one third (36.7%) of the participants had less than 10,000 yuan (approximately USD 1450) annual family income, 87.5% had four or more members in the family, and 26.8% reported their spouses were also HIV positive. The majority of participants (86.0%) had diagnosed as HIV-positive for six years or above, and 92.7% were on ART (Table 1).

HRQoL summary scores (PHS and MHS)

The mean score of PHS was 35.75 ± 9.26 (range: 12.10–62.97), and the mean score of MHS was 36.15 ± 8.86 (range: 8.24–60.92). Univariate analysis indicated that the mean scores of PHS and MHS were both different among groups by gender, education, family annual income, spouse’s HIV status, and ART status (all $P < 0.05$). In addition, PHS scores were significantly lower among older participants. PLH who were married/cohabiting reported significantly higher PHS score than those who were single, divorced, or widowed ($P < 0.05$) (Table 1).

Factors associated with HRQoL summary scores (PHS and MHS)

All nine candidate variables were included in the multiple linear regression models. Five variables (gender, age, family size, family annual income, and ART condition) were significantly associated with PHS score in the final model. Male were more likely to report a higher level of PHS than female ($\beta = 0.123$; $P = 0.009$). Age was significantly negatively associated with the PHS ($\beta = -0.232$; $P < 0.001$) when other variables were controlled. Family size remained negatively correlated with PHS ($\beta = -0.105$; $P = 0.021$). Family annual income was significantly positively associated with PHS ($\beta = 0.126$; $P = 0.003$). PLH who were not on ART reported better PHS than those who were on ART while other variables were held constant ($\beta = 0.141$; $P = 0.001$) (Table 2).

Three variables, gender, family annual income, and ART status, were significantly associated with MHS in the final model. Male were also more likely to report a higher level of MHS than female ($\beta = 0.150$; $P = 0.002$). Family annual income was significantly positively associated with MHS ($\beta = 0.135$; $P = 0.002$). PLH who were not on ART reported better MHS than those who were on ART in the model ($\beta = 0.098$; $P = 0.021$) (Table 2).

Discussion

In this study, two sociodemographic factors were found to be associated with HRQoL, including gender and age. Previous research on the relation between gender and HRQoL of PLH had mixed results. Some studies reported that mental HRQoL of male PLH was better than that of female PLH [42], while others generated opposite results [8, 13]. Other research showed no apparent correlation between gender and HRQoL [10]. This study showed that male PLH in rural China had better physical and mental HRQoL than their female counterparts, and it may be due to cultural differences. China has historically been a male-dominated country, and female PLH may bear more burden originated from social position and role orientation [43, 44]. Age is also a major factor that affects PLH’s physiological functioning. Natural aging of body organs can lead to declined body functions. Various HIV/AIDS comorbidities, such as tuberculosis, tumor, hepatitis, and/or opportunistic infections, all severely compromise the HRQoL of PLH [45, 46]. In this study, older age was associated with lower physical HRQoL. This finding is consistent with reports of other studies [13, 42, 47, 48]. However, age is associated with declining physical health, but not necessarily mental health. The association between age and MHS was not statistically significant in regression analysis, which suggests that older PLH may not always have worse mental health compared to younger PLH. Hence, researchers should carefully consider tailored approaches

Table 1 PHS and MHS by background demographic variables

Variables	n (%)	Physical health summary score		Mental health summary score	
		Mean (SE)	P	Mean (SE)	P
Gender					
Female	288 (55.2)	34.54(9.31)	<0.001	34.67(8.65)	0.001
Male	234(44.8)	37.24(8.99)		37.98(8.79)	
Age groups (year)					
< 45	192(36.8)	38.45(9.46)	<0.001	37.37(9.08)	3.079
45–54	182(34.9)	35.39(8.77)		35.70(9.06)	
≥ 55	148(28.4)	32.69(8.59)		35.12(8.19)	
Marital status					
Married/Cohabiting	425 (81.4)	36.14 (9.24)	0.042	36.44(8.84)	0.119
Divorced/Widowed/Other	97 (18.6)	34.03 (9.18)		34.89(8.91)	
Education					
Yes	312 (59.8)	36.94 (9.33)	< 0.001	37.43(8.83)	<0.001
None	210 (40.2)	33.99 (8.89)		34.26(8.59)	
Family size (people)					
3	65(12.5)	37.59(10.62)	0.094	36.26(9.74)	0.455
4–5	207(39.7)	36.12(9.47)		36.71(8.89)	
≥ 6	250(47.8)	34.96(8.64)		35.66(8.60)	
Family annual income (yuan)					
< 10,000	191(36.7)	34.78(9.12)	0.020	35.06(8.39)	0.008
10,000–19,999	177(34.0)	35.41(8.92)		35.84(9.33)	
≥ 20,000	152(29.2)	37.51(9.59)		37.97(8.68)	
Spouse's HIV status					
HIV–/death/other	382(73.2)	36.23(9.42)	0.049	36.69(9.20)	0.013
HIV+	140(26.8)	34.43(8.72)		34.69(7.70)	
Diagnosis time (year)					
<3	23(4.4)	36.43(12.97)	0.252	34.96(9.80)	0.553
3–5	50(9.6)	37.97(10.38)		37.48(9.64)	
6–8	277(53.3)	35.19(8.95)		35.83(8.92)	
≥ 9	170(32.7)	35.95(8.83)		36.46(8.47)	
ART condition					
On ART	484(92.7)	35.33(9.06)	<0.001	35.88(8.83)	0.012
Not on ART	38(7.3)	41.14(10.13)		39.62(8.69)	

to address the differentiated health needs of specific sub-population (such as female and older PLH) to improve their HRQoL.

China has a distinct family-centered culture and custom, that most PLH receive or require support from their families [49, 50]. In addition, China's law stipulated that grown-up children are duty-bound to support their parents. It is a common practice for generations of Chinese to live in the same household, especially in rural areas. Therefore, the family forms a crucial safety umbrella for PLH, and family members are the primary caregivers for daily care, treatment, psychosocial and financial support,

and childcare [25]. We found that several family characteristics, including family size and family annual income, were associated with PLH's HRQoL. PLH's physical health was lower among those who had larger families, which could be explained by two possible reasons: first, the more members in a family, the fewer shares of limited family resources for every member; second, due to urbanization in the last three decades, a large number of young adults migrate to major cities to seek higher income, leaving behind their young child (ren) to the elder parents. We speculated that the families of PLH in the study mainly consisted of grandchildren of PLH, who could not provide family

Table 2 The multiple linear regression analysis results of PHS and MHS

Model	Physical health summary score (n = 518)		Mental health summary score (n = 518)	
	Standardized β	P	Standardized β	P
Gender (ref. Female)	0.123	0.009	0.150	0.002
Age	-0.232	<0.001	-0.087	0.066
Marital status (ref. Married/Cohabiting)	-0.026	0.562	-0.044	0.329
Education (ref. Yes)	-0.023	0.635	-0.063	0.201
Family size	-0.105	0.021	-0.052	0.265
Family annual income	0.126	0.003	0.135	0.002
Spouse HIV status (ref. HIV-/death/other)	-0.082	0.065	-0.087	0.056
Diagnosis time	0.048	0.253	0.061	0.157
ART condition (ref. on ART)	0.141	0.001	0.098	0.021

support and at the same time increase child-care burden of PLH. The finding indicates that more service and care should be provided to PLH with left-behind children in the family. The associations between higher income and better HRQoL was supported by many previous studies, as good family economic condition means better living conditions and better health care for PLH [13, 42, 47, 48]. Thus, one of the important mechanisms to improve HRQoL of PLH in rural areas is to carry out effective strategies for poverty alleviation, which need a collaborative effort of the whole society.

Another family status deserves attention is spousal HIV status, that having an HIV-positive spouse was marginally associated with lower physical and mental HRQoL. In rural China, the spouse is usually the primary source of caring and support for PLH [42, 51]. Spouse's HIV infection can to a great extent undermine the daily life care, material support, and psychological comfort a family can provide to PLH [52]. In addition, when a family has more than one HIV positive member, the burden of the family will often be more than doubled due to the escalated societal stigma and shame [53, 54]. Hence, HIV seroconcordant couples are in greater need of care and support to cope with aggravating impact of HIV.

Interestingly, current ART use was found to be associated with lower levels of physical and mental health. The finding should be interpreted with caution. At the time of the study, the test-and-treat policy has not been implemented in China, and ART was only provided to PLH with a CD4 lower than 500 (cells/mm³) and/or those with clinical symptoms [35]. The lower level of HRQoL found in PLH who were under ART was due to the advanced disease stages [47, 55, 56].

Limitations

Some limitations of the study should be interpreted. First, we cannot draw causal inferences based on the cross-sectional data. Second, because the participants

were recruited from Anhui province in China, findings may not be generalizable to PLH in other areas. Finally, the self-report data may be subject to socio-desirability bias.

Conclusions

The study added to the current understanding of levels of HRQoL among PLH and the individual- and family-level factors associated with physical and mental HRQoL. The findings identified the family-related factors had a strong relationship on HRQoL of PLH in rural China. The results may have implications for intervention design and future research. Future intervention should be carefully tailored to the specific needs of sub-populations considering their physical and mental HRQoL conditions. For example, PLH with left-behind children in the family would need greater care and support. More intervention efforts should be paid to HIV seroconcordant couples to help them cope with aggravating impact of HIV.

Abbreviations

ART: Antiretroviral therapy; CAPI: computer-assisted personal interview; HIV: Human immunodeficiency virus; HRQoL: Health-related quality of life; MHS: Mental health summary; MOS-HIV: Medical outcomes study HIV health survey; PHS: Physical health summary; PLH: People living with HIV

Acknowledgments

We would like to thank the study participants and the project team members.

Funding

This study was funded by the National Institute of Children Health & Human Development/NIH (Grant number R01HD068165). The funders had no role in the design of the study, collection, analysis, interpretation of data or writing of the manuscript.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YX collected the data, reviewed the literature and drafted the manuscript. CL contributed to data analysis and helped to draft the paper. LL and GJ

designed the study and reviewed the final paper. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The study was reviewed and approved by the Institutional Review Boards of the University of California, Los Angeles in U.S. and the Anhui Provincial Center for Disease Control and Prevention in China. Written informed consent was obtained from all individual participants included in the study.

Consent for publication

Not Applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of HIV/AIDS Prevention, Anhui Provincial Center for Disease Control and Prevention, Hefei, China. ²Semel Institute for Neuroscience and Human Behavior, Center for Community Health, University of California at Los Angeles, Los Angeles, CA, USA. ³Anhui Provincial Center for Woman and Child Health, 38 Yonghong Road, Hefei 230032, China.

Received: 21 May 2018 Accepted: 20 December 2018

Published online: 03 January 2019

References

- UNAIDS. global AIDS update 2016[R].2016.
- Miners A, Phillips A, Kreif N, et al. Health-related quality-of-life of people with HIV in the era of combination antiretroviral treatment: a cross-sectional comparison with the general population. *Lancet HIV*. 2014;1(1):e32–40.
- Charles B, Jeyaseelan L, Pandian AK, et al. Association between stigma, depression and quality of life of people living with HIV/AIDS (PLHA) in South India - a community based cross sectional study. *BMC Public Health*. 2012;12:463.
- Geocz L, Mucci S, De Marco MA, et al. Quality of life and adherence to HAART in HIV-infected patients. *Rev Saude Publica*. 2010;44(4):743–9.
- Da SL, Tavares JS. The family's role as a support network for people living with HIV/AIDS: a review of Brazilian research into the theme. *Cien Saude Colet*. 2015;20(4):1109–18.
- Abrefa-Gyan T, Cornelius LJ, Socio-Demographic Factors OJ, Support S. Quality of life, and HIV/AIDS in Ghana. *J Evid Inf Soc Work*. 2016;13(2):206–16.
- Van der Kop ML, Muhula S, Patel A, et al. Gender differences in health-related quality of life at the time of a positive HIV test - a cross-sectional study in a resource-poor, high prevalence setting in Nairobi, Kenya. *AIDS Care*. 2017;30:1–7.
- Imam MH, Karim MR, Ferdous C, et al. Health related quality of life among the people living with HIV. *Bangladesh Med Res Counc Bull*. 2011;37(1):1–6.
- George S, Bergin C, Clarke S, et al. Health-related quality of life and associated factors in people with HIV: an Irish cohort study. *Health Qual Life Outcomes*. 2016;14(1):115.
- Mbalinda SN, Kiwanuka N, Kaye DK, et al. Reproductive health and lifestyle factors associated with health-related quality of life among perinatally HIV-infected adolescents in Uganda. *Health Qual Life Outcomes*. 2015;13:170.
- Degroote S, Vogelaers DP, Vermeir P, et al. Socio-economic, behavioural, (neuro) psychological and clinical determinants of HRQoL in people living with HIV in Belgium: a pilot study. *J Int AIDS Soc*. 2013;16:18643.
- Chen WT, Barbour R. Life priorities in the HIV-positive Asians: a text-mining analysis in young vs. old generation. *AIDS Care*. 2016;29:1–4.
- Yang Y, Thai S, Choi J. An evaluation of quality of life among Cambodian adults living with HIV/AIDS and using antiretroviral therapy: a short report. *AIDS Care*. 2016;28(12):1546–50.
- Earnshaw VA, Lang SM, Lippitt M, et al. HIV stigma and physical health symptoms: do social support, adaptive coping, and/or identity centrality act as resilience resources? *AIDS Behav*. 2015;19(1):41–9.
- China NHFPC. 2015 China AIDS Response Progress Report.2015.
- Liping M, Peng X, Haijiang L, et al. Quality of life of people living with HIV/AIDS: a cross-sectional study in Zhejiang Province, China. *PLoS One*. 2015; 10(8):e135705.
- Liu H, He X, Levy JA, et al. Psychological impacts among older and younger people living with HIV/AIDS in Nanning, China. *J Aging Res*. 2014;2014: 576592.
- Lan GL, Yuan ZK, Clements-Nolle KD, et al. Social capital and quality of life among people living with HIV/AIDS in Southeast China. *Asia Pac J Public Health*. 2016;28(4):325–35.
- Chen JP, Han MM, Liao ZJ, et al. HIV-related behaviors, social support and health-related quality of life among men who have sex with men and women (MSMW): a cross-sectional study in Chongqing, China. *PLoS One*. 2015;10(2):e118651.
- Xiao Z, Li X, Qiao S, et al. Social support, depression, and quality of life among people living with HIV in Guangxi, China. *AIDS Care*. 2017;29(3): 319–25.
- Li XM, Yuan XQ, Wang JJ, et al. Evaluation of impact of social support and care on HIV-positive and AIDS individuals' quality of life: a nonrandomised community trial. *J Clin Nurs*. 2017;26(3–4):369–78.
- Lan G, Yuan Z, Cook A, et al. The relationships among social support and quality of life in persons living with HIV/AIDS in Jiangxi and Zhejiang provinces, China. *AIDS Care*. 2015;27(8):946–53.
- Wu X, Chen J, Huang H, et al. Perceived stigma, medical social support and quality of life among people living with HIV/AIDS in Hunan, China. *Appl Nurs Res*. 2015;28(2):169–74.
- Song B, Yan C, Lin Y, et al. Health-related quality of life in HIV-infected men who have sex with men in China: a cross-sectional study. *Med Sci Monit*. 2016;22:2859–70.
- Liu H, Xu Y, Lin X, et al. Associations between perceived HIV stigma and quality of life at the dyadic level: the actor-partner interdependence model. *PLoS One*. 2013;8(2):e55680.
- Zhu Y, Liu J, Qu B, et al. Relationship between quality of life and unprotected anal intercourse among Chinese men who have sex with men: a cross-sectional study. *BMC Public Health*. 2016;16:382.
- Hidru TH, Wang F, Lolokote S, et al. Associated factors of self-reported psychopathology and health related quality of life among men who have sex with men (MSM) with HIV/AIDS in Dalian, China: a pilot study. *Infect Dis Poverty*. 2016;5(1):108.
- Xu Y, Lin X, Chen S, et al. Ageism, resilience, coping, family support, and quality of life among older people living with HIV/AIDS in Nanning, China. *Glob Public Health*. 2016;13:1–14.
- Tang X, Zhang Y, Bo Y, et al. Quality of life and its related Factors in patients with HIV/AIDS in rural China. *Iran J Public Health*. 2015;44(3):412–4.
- Xu JF, Ming ZQ, Zhang YQ, et al. Family support, discrimination, and quality of life among ART-treated HIV-infected patients: a two-year study in China. *Infect Dis Poverty*. 2017;6(1):152.
- Ji G, Detels R, Wu Z, et al. Correlates of HIV infection among former blood/plasma donors in rural China. *AIDS*. 2006;20(4):585–91.
- Wu Z, Rou K, Detels R. Prevalence of HIV infection among former commercial plasma donors in rural eastern China. *Health Policy Plan*. 2001; 16(1):41–6.
- He N, Detels R. The HIV epidemic in China: history, response, and challenge. *Cell Res*. 2005;15(11–12):825–32.
- Teng T, Shao Y. Scientific approaches to AIDS prevention and control in China. *Adv Dent Res*. 2011;23(1):10–2.
- Li L, Ji G, Lin C, et al. Antiretroviral therapy initiation following policy changes: observations from China. *Asia Pac J Public Health*. 2016;28(5):416–22.
- Lin C, Li L, Ji G, et al. Emotional social support and access to care among older people living with HIV in rural China. *Int J Geriatr Psychiatry*. 2015; 30(10):1041–7.
- Wu AW. MOS HIV health survey users manual. Baltimore: John Hopkins University; 1999.
- Wu AW, Revicki DA, Jacobson D, et al. Evidence for reliability, validity and usefulness of the medical outcomes study HIV health survey (MOS-HIV). *Qual Life Res*. 1997;6(6):481–93.
- Wu AW, Hays RD, Kelly S, et al. Applications of the medical outcomes study health-related quality of life measures in HIV/AIDS. *Qual Life Res*. 1997;6(6): 531–54.
- Revicki DA, Sorensen S, Reliability WAW. Validity of physical and mental health summary scores from the medical outcomes study HIV health survey. *Med Care*. 1998;36(2):126–37.

41. Huang ZJ, Tian M, Dai SY, et al. Feasibility, reliability and validity of the Chinese simplified version of the MOS-HIV health survey among AIDS patients in China. *Qual Life Res.* 2013;22(2):403–7.
42. Pereira M, Canavarro MC. Gender and age differences in quality of life and the impact of psychopathological symptoms among HIV-infected patients. *AIDS Behav.* 2011;15(8):1857–69.
43. Li L, Lin C, Ji G. Gendered aspects of perceived and internalized HIV-related stigma in China. *Women Health.* 2016;57:1–13.
44. Chen Z, Ge Y, Lai H, et al. Globalization and gender wage inequality in China. *World Dev.* 2013;44:256–66.
45. Balderson BH, Grothaus L, Harrison RG, et al. Chronic illness burden and quality of life in an aging HIV population. *AIDS Care.* 2013;25(4):451–8.
46. Botros D, Somarriba G, Neri D, et al. Interventions to address chronic disease and HIV: strategies to promote exercise and nutrition among HIV-infected individuals. *Curr HIV/AIDS Rep.* 2012;9(4):351–63.
47. Krause DD, Butler KR, May WL. Associations between factors affecting access to care and health-related quality of life: results of a statewide HIV/AIDS cross-sectional study. *AIDS Care.* 2013;25(1):77–84.
48. Hasanah CI, Zaliha AR, Mahiran M. Factors influencing the quality of life in patients with HIV in Malaysia. *Qual Life Res.* 2011;20(1):91–100.
49. Li J, Wang J. Individuals are inadequate: recognizing the family-centeredness of Chinese bioethics and Chinese health system. *J Med Philos.* 2012;37(6):568–82.
50. Li L, Lin C, Ji G, et al. Parents living with HIV in China: family functioning and quality of life. *J Child Fam Stud.* 2009;18(1):93–101.
51. Chandran V, Madi D, Chowta N, et al. Caregiver burden among adults caring for people living with HIV/AIDS (PLWHA) in southern India. *J Clin Diagn Res.* 2016;10(5):C41–3.
52. Yu YJ, Li X, Qiao S, et al. Family relations in the context of HIV/AIDS in Southwest China. *AIDS Care.* 2016;28(10):1261–8.
53. Li L, Liang LJ, Lin C, et al. Gender differences in depressive symptoms among HIV-positive concordant and discordant heterosexual couples in China. *Psychol Women Q.* 2017;41(1):89–99.
54. Talley AE, Ann BB. A relationship-oriented model of HIV-related stigma derived from a review of the HIV-affected couples literature. *AIDS Behav.* 2010;14(1):72–86.
55. Mbada CE, Onayemi O, Ogunmoyole Y, et al. Health-related quality of life and physical functioning in people living with HIV/AIDS: a case-control design. *Health Qual Life Outcomes.* 2013;11:106.
56. Tran BX. Quality of life outcomes of antiretroviral treatment for HIV/AIDS patients in Vietnam. *PLoS One.* 2012;7(7):e41062.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

