

RESEARCH ARTICLE

Open Access



The influence of the municipal human development index and maternal education on infant mortality: an investigation in a retrospective cohort study in the extreme south of Brazil

Carolina Ribeiro Anele^{1*} , Vânia Naomi Hirakata², Marcelo Zubaran Goldani^{1,3,4} and Clécio Homrich da Silva^{1,3,4}

Abstract

Background: Infant mortality is considered an important and sensitive health indicator in several countries, especially in underdeveloped and developing countries. Most of the factors influencing infant mortality are interrelated and are the result of social issues. Therefore, this study performed an investigation of the influence of the MHD and maternal education on infant mortality in a capital in the extreme south of Brazil.

Methods: It is a retrospective cohort study with data on births and deaths in the first year of life for the period of 2000–2017. The association between the independent variables and the outcome was done by bivariate analysis through simple Poisson regression. The variables that can potentially be considered confounding factors were used in a multiple Poisson regression for robust variances - adjusted model.

Results: The study included 317,545 children, of whom 3107 died. The medium MHD showed associated with infant death in the first year of life. Maternal education, individually and jointly analyzed with the MHD, showed association with the outcome of infant death in the first year of life, particularly for children of mothers with lower maternal education ($p < 0.001$). In relation to other related factors, maternal age; number of Prenatal Care Consultations; gestational age, weight, gender and Apgar Index (5th minute) of the newborn showed association with IM ($p < 0.001$).

Conclusions: The HDI is considered a good predictor of infant mortality by some authors and the analyzes of the present study also confirm an association of the medium MHD and its low MHDIE component with infant mortality. In addition, it was maternal education with less than 8 years of study that that demonstrated a higher risk of death, revealing itself to be a social determinant with a relevant impact on infant mortality. Thus, it is possible to conclude that maternal education is available information, and it is superior to the MHD to assess the infant mortality outcome.

Keywords: Infant mortality, Live birth, Vital statistics, Human development, Education

* Correspondence: carolina.anele@hotmail.com

¹Programa de Pós Graduação em Saúde da Criança e do Adolescente - Faculdade de Medicina, Universidade Federal do Rio Grande do Sul (UFRGS), Rua Ramiro Barcelos, 2400, Porto Alegre, RS 90035003, Brazil
Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Infant mortality (IM) is considered, internationally, as one of the most sensitive health indicators, since it is related to social conditions and can also be influenced by prenatal and perinatal factors, as well as environmental factors, especially nutritional and infectious ones [1–3]. Worldwide, countries have dedicated themselves to achieving Millennium Development Goal 4 (MDG 4), which was to reduce by two-thirds the mortality of children under five by the year of 2015. Currently, efforts are focused on the new agenda in which Millennium Development Goal 3 (MDG 3) keeps the reduction of under-five mortality to at least 25 per 1000 live births as a priority [4]. In Brazil there was a significant reduction in IM in the period from 1990 to 2012, when MDG 4 was achieved 3 years before the deadline [5, 6]. However, as in other countries, neonatal deaths still remained high and corresponded to approximately 70% of mortality in the first year of life [1, 7]. Porto Alegre, the state capital located in the extreme south of Brazil and where the present study was carried out, presented a variation in the infant mortality rate from 18.22 in 1994 to 9.13 deaths per thousand live births in 2018, which represented a decrease of 49.89% [8]. For the epidemiological surveillance of such vital events, in Brazil, the Live Births Information System (SINASC) and the Mortality Information System (SIM) are used [9].

Most of the factors influencing IM are interrelated and are the result of social issues, including the educational level of the mother, which is an important social determinant for better child health conditions because it leads to a better maternal perception of health needs and problems. Maternal education seems to be directly related to the healthy growth and development of the child [10, 11]. The region of residence is also related, albeit indirectly, to IM, being an important social factor in child health outcomes [12], as less developed regions may have worse physical structure and sanitation conditions and limited access to health services [3, 13, 14].

There are important differences in these factors between countries, especially in developing countries like Brazil, where there are differences between regions according to socio-economic characteristics, with significant social inequities. As an alternative to assess these development disparities in different countries, the Human Development Index (HDI) evaluates human development through three indicators: schooling (expected years of schooling for children of school age and average schooling in years for adults over 25), longevity (life expectancy at birth) and income (per capita income). In this context, the Municipal Human Development Index (MHDI) was launched in Brazil in 2013. It was a strategy to adapt the HDI to the local context, allowing the calculation of HDI at the municipal level of the

metropolitan regions of Brazil. MHDI considers the same three dimensions of Global HDI (longevity, education and income), including its three components, MHDI Income (MHDII), MHDI Longevity (MHDIL), MHDI Education (MHDIE), but it is adapted to the national context and uses local indicators available for the calculation. In the MHDI, the geometric mean is used: the dimensions (longevity, income and education) are multiplied and the product is extracted by the cube root, for this calculation the three components have the same weight [15]. The general MHDI of Porto Alegre was 0.805 in 2010, ranking as the 7th best Brazilian capital [15].

Because it is a sensitive indicator for socioeconomic and environmental changes, IM should be constantly monitored, and a better understanding of the association of HDI and maternal education with IM assists in the planning of actions and policies aimed at maternal and child health, particularly in regions of greater social vulnerability [1, 3]. Therefore, this study performed an investigation of the influence of the MHDI and maternal education on infant mortality in a capital in the extreme south of Brazil.

Methods

Study design and participants

This is a retrospective cohort study that used the annual SIM and SINASC databases provided by the Non transmissible Vital Events, Diseases and Aggravates Surveillance Team of the General Health Surveillance Coordination of the Porto Alegre Municipal Health Secretariat.

The city of Porto Alegre has an estimated population of 1,479,101 people and has eight Planning Management Regions that are subdivided into seventeen Participatory Budget Regions (macro-regions), each of which is made up of a set of neighborhoods with affinities to each other [8], it has an area of 471.85km². In relation to education level, in 2017, the illiteracy rate was 2.27%, the Basic Education Development Index (IDEB) was 4.9 (target 5.5) for the initial years and 3.9 (target 4.7) for the final years. Regarding mothers heads of families from 2000 to 2010, the percentage of women without complete primary education showed a decrease of 10%, from 39.87 to 29.15% [8].

The study included all live births in the city of Porto Alegre from January 2000 to December 2016 obtained at SINASC and all deaths of children under 1 year of age from January 2000 to the month December 2017, obtained from SIM. Thus, all children were included in the study until they completed their first year of life. A linkage was made between the two databases through the number of the Declaration of Live Birth (DLB), that refers to document filled in at the maternity hospital when

the child is born, the name of the mother and the date of birth, in order to establish a unique database. After a first automatic attempt, there was no agreement on the number of DLB of 269 children who died in the first year of life (7.2% of total deaths) between the two databases. From then on, a manual search was carried out in an attempt to correct and locate the number of the DLB, through the name of the mother, which was confirmed by the date of birth and weight of the newborn and, in a few cases, the address of the maternal residence. From this process, a further 165 deaths were located (4.4% of the total deaths of children in the first year) and the DLB numbers were corrected according to the SINASC database, totaling a final loss of 104 children (2.8% of the total deaths).

After performing the linkage between the SINASC and SIM databases, the newborns twins or higher (2.5% of all live births in the period studied) were excluded to avoid double maternal data and to avoid confusion bias as they are a population that represents greater use of assisted conception, a method used in the private sector. Newborns with birth weight of less than 500 g (0.09% of all live births) also were excluded following publications from other studies investigating infant mortality. Also excluded who were not born in the municipality of Porto Alegre (0.65% of live births) and who had their maternal address ignored (0.36% of live births).

Although there may have been double entry of mothers due to the effect of the long period analyzed, it was considered that, eventually, they may have changed homes and also improved their level of education. However, it is important to note that, throughout the study period, the percentage of mothers who were under 18 years of age and, therefore, with greater possibility of changing their level of education, was only 7.8%. In addition, the study population is based on the newborn that will become a new case even if it is a second child of a given mother.

From SINASC, the following variables were used: number of Declaration of Live Birth; maternal age (10–17, 18–34, ≥ 35 years); maternal education in completed years of study (< 8 ; 8–11; ≥ 12 years); maternal home district (see Additional file 1) [16]; Planning Management Region of Maternal Residence; Participatory budgeting Regions of Maternal Residence, which were later called macro-regions (see Additional file 2 and Additional file 3) [16]; maternal marital status (married/stable union, single/separated/widowed); number of previous living children; number of previous dead children (none; ≥ 1); number of Prenatal Care Consultations (None; 1–3; 4–6; ≥ 7); gestational age in weeks (< 27 , 28–31, 32–36, ≥ 37); type of hospital (public, private, mixed); type of delivery (vaginal, cesarean); weight of the newborn (grams); Apgar rate at the 5th minute of life (≥ 7 , < 7); gender of

the newborn (female, male). The study considered the maternal data at birth contained in the database of SINASC, since this system presents a greater completeness of the data, mainly in the city of the study that presents itself as one of the capitals with the best coverage and filling of this system [9, 11]. From SIM, the variables used were: the number of the Declaration of Live Birth and date of death. The following variables were also introduced in the bank: year; parity, created from the number of live children and number of dead children (primipara, multipara) and the value of the MHDI of the macro-regions of maternal domicile and its three components: MHDI Income (MHDII), MHDI Longevity (MHDIL) and MHDI Education (MHDIE). The study used the following cutoff points previously calculated and established by the United Nations Development Program Brazil (UNDP Brazil), the Institute for Applied Economic Research (IPEA) and the João Pinheiro Foundation, classified as very low (0–0.499), low (0.5–0.599), medium (0.6–0.699), high (0.7–0.799) and very high (0.8–1.0) and available through the Atlas Human Development online platform at Brazil [15]. The values of each component and the MHDI were calculated based on the 2010 Demographic Census of the Brazilian Institute of Geography and Statistics (IBGE). In the municipality studied, no macro-region has an MHDI lower than 0.6. Thus, according to the cutoff points already established, the MHDI of the macro-regions was classified only as medium, high and very high. The exception was only for the MHDIE component, which had a low rating in some macro-regions of the study.

Statistical analyses

Preliminarily, a descriptive analysis of variables with absolute and relative frequencies was presented. The association of variables of the study, according to the MHDI classification, with the death outcome was performed by Chi-square test. Later, the association between the determinants involved (independent variables) with the outcome of death was verified by a bivariate analysis through simple Poisson regression. Finally, the variables that can potentially be considered confounding factors, as well as maternal education, were used in four models adjusted to estimate the influence of MHDI, MHDII, MHDIL and MHDIE on infant mortality in a multiple Poisson regression for robust variances.

Ethical aspects

The study was approved by the Research Ethics Committees of Hospital de Clínicas of Porto Alegre and the Research Ethics Committees of Municipal Health Department of Porto Alegre, respectively, under the respective protocol numbers 2,940,235 and 3,153,671.

Results

Three hundred seventeen thousand five hundred forty-five children were included in the study. Of these 3107 died in the first year of life (Fig. 1).

The proportion of mothers aged 35 years or older increases according to the MHDI classification, being higher (25.2%) in those residing in macro-regions with very high MHDI. Independent of the MHDI classification, a maternal age under 18 years showed association with child death in the first year of life ($p < 0.05$). In macro-regions with high and very high MHDI, single mothers, judicially separated or widowed, multiparas and with one or more previous dead children also showed association with death ($p < 0.05$). Regardless of the MHDI classification, the lower the number of Prenatal Care Consultations the higher the percentage of deaths. The percentage of cesarean sections was higher in macro-regions with high and very high MHDI (43 and 60.4%, respectively). In the medium and high MHDI classifications, cesarean section showed a significant association with death, while in very high MHDI vaginal delivery showed a significant association with death. In the three classifications of MHDI (medium, high and very high), the proportion of deaths was higher for those newborns with a lower gestational age, with low weight or with an Apgar Index of less than 7 at the fifth minute

of life. Independent of the MHDI classification have congenital anomaly showed an association with child death in the first year of life ($p < 0.05$). The public hospital showed a significant association with death, regardless of the MHDI classification (Table 1). In the three classifications of the MHDI, there is a decrease in MI in the city of Porto Alegre during the study period (see Additional file 4).

The majority of mothers were between 18 and 34 years old, 64.2% ($n = 203,417$) of them were single, legally separated or widowed and the minority, 44.8% ($n = 142,143$) were primiparas which proved to be a protective factor for infant mortality. Regarding prenatal care, 32% ($n = 101,322$) had less than seven consultations, which proved to be a risk factor for infant death in the first year of life. Vaginal delivery occurred in 53.6% ($n = 170,220$) of births and most of them in mixed or private hospitals (53.4%). Newborns with a gestational age of less than 37 weeks, an Apgar Index of less than 7 at the fifth minute of life, low weight and congenital anomaly were also considered risk factors for death ($p < 0.001$). With the exception of the age of 35 or more and type of delivery, all other variables presented significant association with death in the preliminary crude analysis (Table 2), but in the consecutive multivariate analysis, maternal age and previous number of dead children were not associated with the infant mortality outcome.




Fig. 1 Description of the study population: live births and infant deaths in the first year of life in the municipality of Porto Alegre, Brazil (2000–2017). figure made by the author. Source: figure made by the author

Table 1 Distribution of the absolute frequency of sociodemographic, perinatal and neonatal characteristics with the respective mortality percentages according to the MHDl classification in Porto Alegre (Rio Grande do Sul, Brazil) from 2000 to 2017

	MHDl					
	MEDIUM n(%)		HIGH n(%)		VERY HIGH n(%)	
	TOTAL	DEATH	TOTAL	DEATH	TOTAL	DEATH
Maternal Age						
10 to 17 years	5540 (10.6)	86 (1.6) ^a	15,708 (8.6)	218 (1.4) ^a	3628 (4.4)	66 (1.8) ^a
18 to 34 years	40,411 (77.5)	446 (1.1)	140,685 (77.2)	1358 (1.1)	58,530 (70.5)	436 (0.7)
≥ 35 years	6170 (11.9)	80 (1.3)	25,859 (14.2)	280 (1.1)	20,907 (25.2)	132 (0.6)
	52,121 (100)	612 (1.2)	182,252 (100)	1856 (1.1)	83,065 (100)	634 (0.8)
Maternal Education						
< 8 years	20,301 (39.2)	301 (1.5) ^a	59,084 (32.5)	867 (1.5) ^a	13,486 (16.3)	205 (1.5) ^a
≥ 8 to 11 years	24,981 (48.2)	251 (1.0)	83,658 (46.1)	752 (0.9)	25,528 (30.8)	216 (0.8) ^a
≥ 12 years	6541 (12.6)	50 (0.8)	38,806 (21.4)	224 (0.6)	43,780 (52.9)	205 (0.5)
	51,823 (100)	602 (1.2)	181,548 (100)	1843 (1.0)	82,794 (100)	626 (0.8)
Marital Status						
Single/Separated/Widowed	37,022 (71.3)	452 (1.2)	123,195 (67.8)	1428 (1.2) ^a	43,144 (52.1)	434 (1.0) ^a
Married/stable union	14,937 (28.7)	156 (1.0)	58,639 (32.2)	416 (0.7)	39,739 (47.9)	192 (0.5)
	51,959 (100)	608 (1.2)	181,834 (100)	1844 (1.0)	82,883 (100)	626 (0.8)
Parity						
Primipara	20,293 (39)	221 (1.1)	78,595 (43.2)	685 (0.9)	43,217 (52.1)	273 (0.6)
Multipara	31,684 (61)	388 (1.2)	103,341 (56.8)	1162 (1.1) ^a	39,707 (47.9)	355 (0.9) ^a
	51,977 (100)	609 (1.2)	181,936 (100)	1847 (1.0)	82,924 (100)	628 (0.8)
Number of Dead Children						
≥ 1	5038 (9.7)	67 (1.3)	17,376 (9.5)	225 (1.3) ^a	8038 (9.7)	83 (1.0) ^a
None	47,067 (90.3)	545 (1.2)	164,833 (90.5)	1630 (1.0)	75,008 (90.3)	550 (0.7)
	52,105 (100)	612 (1.2)	182,209 (100)	1855 (1.0)	83,046 (100)	633 (0.8)
Number Prenatal Care Consultations						
None	1844 (3.6)	78 (4.2) ^a	6048 (3.3)	286 (4.7) ^a	1808 (2.2)	92 (5.1) ^a
from 1 to 3	5530 (10.7)	137 (2.5) ^a	15,889 (8.7)	427 (2.7) ^a	4285 (5.2)	103 (2.4) ^a
from 4 to 6	13,463 (26.0)	192 (1.4) ^a	40,524 (22.3)	518 (1.3) ^a	11,887 (14.3)	164 (1.4) ^a
7 and more	30,993 (59.8)	199 (0.6)	119,233 (65.6)	605 (0.5)	64,885 (78.3)	266 (0.4)
	51,830 (100)	606 (1.2)	181,694 (100)	1836 (1.0)	82,865 (100)	625 (0.8)
Birth						
Vaginal	33,379 (64.0)	361 (1.1)	103,908 (57.0)	1007 (1.0)	32,869 (39.6)	300 (0.9) ^a
Cesarean	18,744 (36.0)	252 (1.3) ^a	78,340 (43.0)	848 (1.1) ^a	50,196 (60.4)	334 (0.7)
	52,123 (100)	613 (1.2)	182,248 (100)	1855 (1.1)	83,065 (100)	634 (0.8)
Gestational Age						
< 27 weeks	232 (0.4)	133 (57.3) ^a	790 (0.4)	465 (58.9) ^a	295 (0.4)	151 (51.2) ^a
28 to 31 weeks	473 (0.9)	85 (18.0) ^a	1523 (0.8)	234 (15.4) ^a	634 (0.8)	78 (12.3) ^a
32 to 36 weeks	4331 (8.3)	123 (2.8) ^a	14,926 (8.2)	349 (2.3) ^a	6979 (8.4)	115 (1.6) ^a
37 or more	46,967 (90.3)	261 (0.6)	164,641 (90.7)	764 (0.5)	75,032 (90.5)	276 (0.4)
	52,003 (100)	602 (1.2)	181,880 (100)	1812 (1.0)	82,940 (100)	620 (0.7)
Apgar At 5 Minutes						
< 7	686 (1.3)	168 (24.5) ^a	2205 (1.2)	592 (26.8) ^a	694 (0.8)	179 (25.8) ^a
≥ 7	50,961 (98.7)	410 (0.8)	178,989 (98.8)	1158 (0.6)	82,091 (99.2)	416 (0.5)

Table 1 Distribution of the absolute frequency of sociodemographic, perinatal and neonatal characteristics with the respective mortality percentages according to the MHDl classification in Porto Alegre (Rio Grande do Sul, Brazil) from 2000 to 2017 (Continued)

	MHDl					
	MEDIUM n(%)		HIGH n(%)		VERY HIGH n(%)	
	TOTAL	DEATH	TOTAL	DEATH	TOTAL	DEATH
	51,647 (100)	578 (1.1)	181,194 (100)	1750 (1.0)	82,785 (100)	595 (0.7)
Low Weight						
No (≥ 2500 g)	47,425 (91.0)	262 (0.6)	166,558 (91.4)	762 (0.5)	76,581 (92.2)	268 (0.3)
Yes (< 2500 g)	4698 (9.0)	351 (7.5) ^a	15,694 (8.6)	1094 (7.0) ^a	6485 (7.8)	366 (5.6) ^a
	52,123 (100)	613 (1.2)	182,252 (100)	1856 (1.0)	83,066 (100)	634 (0.8)
Congenital Anomaly						
Yes	790 (1.5)	153 (19.4) ^a	2857 (1.6)	469 (16.4) ^a	1062 (1.3)	168 (15.8) ^a
No	50,958 (98.5)	450 (0.9)	178,634 (98.4)	1362 (0.8)	81,668 (98.7)	456 (0.6)
	51,748 (100)	603 (1.2)	181,491 (100)	1831 (1.0)	82,730 (100)	624 (0.8)
Type of Hospital						
Public	25,051 (48.3)	317 (1.3) ^a	93,969 (51.7)	1135 (1.2) ^a	28,492 (34.4)	320 (1.1) ^a
Mixed	20,679 (39.9)	247 (1.2)	51,443 (28.3)	544 (1.1)	13,217 (16.0)	154 (1.2) ^a
Private	6106 (11.8)	34 (0.6)	36,204 (19.9)	125 (0.3)	41,108 (49.6)	126 (0.3)
	51,836 (100)	598 (1.2)	181,616 (100)	1804 (1.0)	82,817 (100)	600 (0.7)

^aStatistically significant association by the residue test adjusted to 5% significance. MHDl Municipal Human Development Index. n Number of participants. %: percentage

In the bivariate analysis, through Poisson's simple regression, the maternal education equal or superior to 12 years of study was a protective factor for death. In relation to the general MHDl in its three components, the lower its classification, the higher was the risk of child death in the first year of life. Children of mothers residing in the medium MHDl score died 1.54 times more when compared to those of mothers with very high MHDl scores. On the other hand, children of mothers residing in macro-regions with low MHDl died 1.66 times more often when compared to those of mothers with very high MHDl (Table 3).

The MHDl and its components were analyzed separately with the other variables, so an adjusted model with MHDl and schooling, a model with MHDII and schooling, a model with MHDIL and schooling and finally a model with MHDIE and schooling through multiple Poisson regression adjusting for statistically significant variables in the bivariate analysis to assess the influence of MHDl and maternal education on infant mortality. The first column of Table 4 describes the classifications of each MHDl, as well as maternal education. The MHDl with a medium classification and its MHDIE component with a low classification remained associated with infant death in the first year of life ($p = 0.036$ and $p = 0.037$ respectively), both presenting a 16% higher risk of death. The analysis of maternal education showed that a number less than 8 years of study maintained an association with infant death ($p < 0.001$) both in the model

with the general MHDl and in its 3 components, presenting a 37 to 40% higher risk of death (Table 4).

Discussion

In recent years, different authors have investigated factors related to infant mortality and maternal-infant health conditions. Regarding the influence of socioeconomic factors, the literature has shown that there are differences according to the place studied [17–19]. In the present study, besides macro-regions with medium MHDl being associated with death ($p = 0.036$), maternal education, individually and jointly analyzed with the MHDl, showed association with the outcome of infant death in the first year of life, particularly for children of mothers with lower maternal education ($p < 0.001$). In relation to other related factors, low maternal age; number of Prenatal Care Consultations; gestational age, weight, gender, congenital anomaly, and Apgar Index (5th minute) of the newborn showed association with IM ($p < 0.001$) in the municipality of Porto Alegre from 2000 to 2017. These results corroborate previous studies that identified biological and prenatal care factors related to infant death [1, 20–22].

Although most mothers were between 18 and 34 years of age, the highest proportion of child deaths in the first year of life (1.5%) occurred among the children of those under 18. This age group presents a higher risk for complications, since younger women are still in the phase of physical and psychological development and, part of

Table 2 Distribution of absolute frequency and association of socio-demographic, perinatal and neonatal characteristics on Infant Mortality in Porto Alegre (Rio Grande do Sul, Brazil) from 2000 to 2017

	TOTAL n(%)	DEATH n(%)	RR [CI95%]	Value of <i>p</i>
Maternal Age				
10 to 17 years	24,882 (7.8)	370 (1.5)	1.59 [1.42–1.77]	< 0.001
≥ 35 years	52,952 (16.7)	493 (0.9)	0.99 [0.90–1.09]	0.926
18 to 34 years	239,706 (75.5)	2242 (0.9)	1	
Marital Status				
Single/Separated/Widowed	203,417 (64.2)	2317 (1.1)	1.69 [1.56–1.83]	< 0.001
Married/stable union	113,361 (35.8)	764 (0.7)	1	
Parity				
Primipara	142,143 (44.8)	1180 (0.8)	0.76 [0.71–0.82]	< 0.001
Multipara	174,796 (55.2)	1907 (1.1)	1	
Number of dead children				
≥ 1	30,460 (9.6)	377 (1.2)	1.30 [1.17–1.45]	< 0.001
None	287,004 (90.4)	2727 (1.0)	1	
Number of Prenatal Care Consultations				
None	9713 (3.1)	457 (4.7)	9.45 [8.48–10.53]	< 0.001
From 1 to 3	25,711 (8.1)	668 (2.6)	5.22 [4.74–5.74]	< 0.001
From 4 to 6	65,898 (20.8)	874 (1.3)	2.66 [2.44–2.91]	< 0.001
7 and more	215,170 (68.0)	1071 (0.5)	1	
Birth				
Vaginal	170,220 (53.6)	1671 (1.0)	1.0 [0.94–1.08]	0.828
Cesarean	147,320 (46.4)	1435 (1.0)	1	
Gestational Age				
Less than 27 weeks	1317 (0.4)	749 (56.9)	124.96 [116.31–134.25]	< 0.001
28 to 31 weeks	2631 (0.8)	397 (15.1)	33.15 [29.83–36.85]	< 0.001
32 to 36 weeks	26,244 (8.3)	587 (2.2)	4.91 [4.46–5.41]	< 0.001
37 weeks or more	286,735 (90.5)	1305 (0.5)	1	
Apgar at 5 min				
< 7	3587 (1.1)	940 (26.2)	41.19 [38.39–44.19]	< 0.001
≥ 7	312,140 (98.9)	1986 (0.6)	1	
Low Weight				
No (≥2500 g)	290,659 (91.5)	1294 (0.4)	0.07 [0.06–0.07]	< 0.001
Yes (< 2500 g)	26,886 (8.5)	1813 (6.7)	1	
Congenital Anomaly				
Yes	4711 (1.5)	791 (16.8)	23.02 [21.34–24.83]	< 0.001
No	311,361 (98.5)	2271 (0.7)	1	
Newborn Sex				
Female	154,886 (48.8)	1411 (0.9)	0.88 [0.82–0.95]	< 0.001
Male	162,630 (51.2)	1680 (1.0)	1	
Type of Hospital				
Public	147,558 (46.6)	1774 (1.2)	3.50 [3.09–3.97]	< 0.001
Mixed	85,357 (27.0)	945 (1.1)	3.23 [2.83–3.69]	< 0.001
Private	83,456 (26.4)	286 (0.3)	1	

Dependent variable: death. RR Relative risk, CI95% 95% confidence interval. n Number of participants. %: percentage. <: less

Table 3 Association of the MHDl (and its components) and maternal education on Infant Mortality (unadjusted model) in Porto Alegre (Rio Grande do Sul, Brazil) in the period 2000 to 2017

	Total (n)	Death n(%)	RR [CI95%]	Value of p
MHDl				
Medium	52,123	613 (1.2)	1.54 [1.38–1.72]	< 0.001
High	182,252	1856 (1.0)	1.33 [1.22–1.46]	< 0.001
Very high	83,066	634 (0.8)	1	
MHDII				
Medium	31,958	381 (1.2)	1.40 [1.25–1.57]	< 0.001
High	137,472	1466 (1.1)	1.26 [1.17–1.35]	< 0.001
Very high	148,011	1256 (0.8)	1	
MHDIL				
High	11,554	139 (1.2)	1.24 [1.05–1.47]	0.012
Very high	305,887	2964 (1.0)	1	
MHDIE				
Low	59,957	704 (1.2)	1.66 [1.47–1.87]	< 0.001
Medium	136,948	1451 (1.1)	1.49 [1.34–1.66]	< 0.001
High	59,988	519 (0.9)	1.22 [1.07–1.39]	0.002
Very high	60,548	429 (0.7)	1	
Maternal Education				
< 8 years	92,914	1375 (1.5)	2.75 [2.48–3.05]	< 0.001
≥ 8 to 11 years	134,210	1219 (0.9)	1.69 [1.52–1.87]	< 0.001
	89,143	480 (0.5)	1	

Dependent variable: death. RR Relative risk, CI95% 95% confidence interval, MHDl Municipal Human Development Index, MHDII Income component of the Municipal Human Development Index, MHDIL Longevity component of the Municipal Human Development Index, MHDIE Education component of the Municipal Human Development Index, n Number of participants. %: percentage. <: less

them, performing their studies in primary, secondary or even higher education [23, 24]. Single, legally separated or widowed women presented a higher risk of death than those married or in a stable union (RR = 1.69; CI95% 1.56–1.83). In literature, other studies have described that the condition of women without a fixed partner negatively affects the health of the mother and child, because it generates some degree of suffering for the woman, who may develop, among other conditions, depression, thus influencing the mother-baby interaction. Moreover, it represented a decrease in emotional and economic support for the family [25–27].

As for the type of birth, most women have had vaginal birth. However, the present study did not evaluate the temporal evolution of this variable, since it is known that this percentage has varied according to the year. Although the evidence does not show an association between cesarean section and a decrease in IM, the number of cesarean sections has been steadily increasing in both developing and developed countries [28, 29]. In 2015, for the first time in Brazil, the number of operative deliveries stabilized [6]. The proportion of cesarean sections was higher in macroregions with better MHDl scores. This ratio, according to different authors, could be explained by the preference for cesarean sections with higher socioeconomic level, white ethnicity and higher maternal education [30, 31]. Moreover, the most visible discrepancies are observed when comparing cesarean rates in public and private hospitals in several regions of Brazil [31, 32]. These rates in the private sector are significantly higher than in the public health system, exceeding 90% in some hospitals [33, 34].

Table 4 Association of MHDl, MHDII, MHDIL, MHDIE and its classification (in four different robust Poisson regression models) and maternal education on Infant Mortality (adjusted models)^a in Porto Alegre (Rio Grande do Sul, Brazil) in the period 2000 to 2017

	MHDl		MHDII		MHDIL		MHDIE	
	RR [CI95%]	p						
Classification of MHDl								
Low	–	–	–	–	–	–	1.16 [1.01–1.34]	0.037
Medium	1.16 [1.01–1.33]	0.036	1.09 [0.96–1.24]	0.170	–	–	1.07 [0.94–1.22]	0.305
High	1.03 [0.91–1.15]	0.637	1.04 [0.95–1.14]	0.360	0.99 [0.83–1.20]	0.978	0.92 [0.78–1.07]	0.284
Very high	1		1		1		1	
Maternal Education								
< 8 years	1.38 [1.20–1.59]	< 0.001	1.39 [1.20–1.60]	< 0.001	1.40 [1.22–1.61]	< 0.001	1.37 [1.19–1.58]	< 0.001
≥ 8 to 11 years	1.04 [0.91–1.19]	0.532	1.05 [0.92–1.19]	0.482	1.06 [0.93–1.20]	0.394	1.04 [0.91–1.18]	0.565
≥ 12 years	1		1		1		1	

Dependent variable: death. RR Relative risk, CI95% 95% confidence interval, MHDl Municipal Human Development Index, MHDII Income component of the Municipal Human Development Index, MHDIL Longevity component of the Municipal Human Development Index, MHDIE Education component of the Municipal Human Development Index, n Number of participants. %: percentage. <: less

^aThe model was adjusted for maternal age, parity, number of prenatal care consultations, gestational age, number of dead children, maternal marital status, type of delivery, Apgar, weight (appropriate; low), congenital anomaly, sex of the infant, type of hospital

The MHDI has proven to be an indicator that has a relationship with the variables studied. The lower the classification of the MHDI of the macro-region of maternal residence, the worse were the socioeconomic and demographic conditions. Previous studies that evaluated infant mortality in different regions or ethnic groups found similar results [18, 19]. Cruz and collaborators [17] observed that, in Brazil, women living in more developed regions have lower chances of having an early pregnancy. Approximately, an analysis conducted in Nigeria with population data found that lower maternal age, as well as lower education, especially of mothers who had deceased children showed association with child death [18]. In this context, this same association is found with the worst maternal conditions observed in less developed or developing countries, as well as in regions with lower income and limited access to health services [3, 35].

In the same perspective, although studies show that prenatal care coverage in Brazil has increased by up to 98.7%, the percentage of women with less than 7 prenatal visits is frequent and may vary according to the region analyzed [36, 37]. Data regarding the city in this study showed an improvement of this indicator by 41.71% in the period from 2001 to 2017. However, it is known that this percentage varied according to interurban and health care differences, and there may be regions with adequate Prenatal Care Consultations above 80% and others below 65% [38]. Although the present study did not specifically investigate the annual temporal evolution of prenatal care coverage, these percentages are similar to the results found for this variable.

Research that assessed the influence of the HDI of the maternal home region, when at birth, on health outcomes, showed that this indicator when low is associated with worse outcomes with respect to child neuropsychomotor development [39, 40] and also with higher infant mortality rates [19, 41]. In a study with data from 188 countries, Ruiz and collaborators [41] found that IM was correlated with low HDI and low inequality-adjusted HDI, with the latter being more strongly correlated than the former ($Z = 2.524$, $p = 0.012$).

Although higher maternal education does not guarantee protection for infant death in the first year of life, as shown in the analysis, schooling less than 8 years of study represents a risk factor increasing the chance of death by 37 to 40%, both in the MHDI and its three components. That is, even though the higher maternal education level did not show statistical significance in reducing infant mortality in the present study, the low educational level showed an association with infant death. It is assumed that these results are attributed to the worst socioeconomic conditions of that mother and the social and community environment in which she

lives with limited access to various resources and health services as well as worse conditions of basic sanitation [22, 42, 43]. Viellas and collaborators [36] found that in Brazil, women with lower education had less prenatal care coverage, fewer prenatal visits and higher utilization of public services. In contrast, they found that those with higher education used more specialized prenatal services, mostly in private services, with monitoring by the same professional throughout pregnancy. In this sense, prenatal care is a process of health education for women and their families, collaborating in the care of pregnancy and children, including, often, the beginning of health care for some individuals, especially adolescents [44].

However, other studies have shown that, in addition to socioeconomic conditions, higher maternal education has improved the child's health conditions and prevented infant mortality. The higher educational level of mothers, together with their better intellectual capacities, help in health choices, increase the understanding and use of information and improve the perception of health problems [45]. They also invest in more health-beneficial behaviors, both at the individual and child levels, showing better adherence to health recommendations in terms of nutrition, exercise and reduction of harmful habits [46–49].

Some particular features of the study are worth highlighting. The study was a pioneer in the country in analyzing the relationship of the MHDI of the different regions of a municipality with infant mortality. It was also unprecedented in performing a parallel analysis of the influence of maternal schooling and MHDI on IM. Furthermore, it used two well consolidated health information systems in the municipality over an extended period of time (2000–2016), which provided a robust database with higher quality in its processing and analysis. This study was important because it revealed that maternal education was shown to be superior to the MHDI as a predictor of infant mortality, reinforcing that when schooling is very low, it can contribute to risky behaviors, and perhaps in addition to being a social indicator, it should be considered a health indicator for the child, as it helps in the health choices of the child's mother and adherence to recommendations. With a health indicator, the mother's educational level could guide the elaboration of public policies, assisting in the screening of risk groups not only by the region of residence, but also by the individual maternal conditions and thus enabling more agile and targeted actions for these individuals of greater risk.

On the other hand, the study presented some limitations. Among them, the non-use of the race/skin color variable, as it was incomplete in some of the years analyzed, and the use of secondary data with pre-established variables and lack of information on serious diseases and

maternal characteristics (income, maternal weight and smoking) and more specific neonatal characteristics (breastfeeding). One hypothesis to explain that MHDl did not present a higher risk of death, as well as what has already been evidenced in the literature is that just like the HDI, the MHDl when analyzed in macro-regions can hide areas of low development and social vulnerability. This could have influenced the analyses since the MHDl values are distributed to the macro-region, not including in the analytical process the micro-regions classified with low or very low MHDl. Porto Alegre currently has 335 micro-regions [15], of which many have very low MHDl values, especially when analyzing the education component. However, the database used did not have a list with the address codes for each micro-region.

Conclusions

The HDI is considered a good predictor of infant mortality by some authors and the analyzes of the present study also confirm an association of the medium MHDl and its low MHDIE component with infant mortality. In addition, it was maternal education with less than 8 years of study that that demonstrated a higher risk of death, revealing itself to be a social determinant with a relevant impact on infant mortality. Thus, it is possible to conclude that maternal education is available information, and it is superior to the MHDl to assess the infant mortality outcome.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-021-10226-9>.

Additional file 1: Figure 1. Map of the studied city by districts (Porto Alegre, Rio Grande do Sul, Brazil). Map of the studied city by districts (Porto Alegre, Rio Grande do Sul, Brazil). OBSERVAPOA and PROCENPA, 2016. Public domain. http://observapoa.com.br/default.php?reg=259&p_secao=46

Additional file 2: Figure 2. Map of the studied city by Participatory budgeting Regions - macro-regions (Porto Alegre, Rio Grande do Sul, Brazil). Map of the studied city by Participatory budgeting Regions - macro-regions (Porto Alegre, Rio Grande do Sul, Brazil). OBSERVAPOA and PROCENPA, 2016. Public domain. http://proweb.procempa.com.br/pmpa/prefpoa/observatorio/usu_doc/sem_grids.pdf

Additional file 3: Table 1. Planning Management Regions and Participatory budgeting Regions (macro-regions) and their corresponding MHDl values and their 3 components of Porto Alegre (Rio Grande do Sul, Brazil). Planning Management Regions and Participatory budgeting Regions (macro-regions) and their corresponding MHDl values and their 3 components of Porto Alegre (Rio Grande do Sul, Brazil). OBSERVAPOA and PROCENPA, 2016. http://observapoa.com.br/default.php?reg=259&p_secao=46

Additional file 4: Graphic 1. Infant Mortality Rate per thousand live births according to the Municipal Human Development Index (MHDl) classification in Porto Alegre (Rio Grande do Sul, Brazil) from 2000 to 2017. Infant Mortality Rate per thousand live births according to the Municipal Human Development Index (MHDl) classification in Porto

Alegre (Rio Grande do Sul, Brazil) from 2000 to 2017. Graphic made by the author.

Additional file 5. Title of data: Study dataset. Dataset generated by the similarity of the two databases related to the current study.

Abbreviations

IM: Infant mortality; MDG 4: Millennium Development Goal 4; MDG 3: Millennium Development Goal 3; HDI: Human Development Index; MHDl: Municipal Human Development Index; MHDIE: Income Component of the Municipal Human Development Index; MHDIL: Longevity Component of the Municipal Human Development Index; MHDIE: Education Component of the Municipal Human Development Index; SINASC: Live Births Information System; SIM: Mortality Information System; IBGE: Brazilian Institute of Geography and Statistics

Acknowledgements

I would like to thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES for the scholarship received during my master's degree and the Non transmissible Vital Events, Diseases and Aggravates Surveillance Team of the General Health Surveillance Coordination of the Porto Alegre Municipal Health Secretariat for making the database available.

Authors' contributions

CRA Preparation of the original project; data collection; data processing and analysis; writing of the preliminary and final version of the manuscript; review and editing. VNH Database organization; data processing and analysis. MZG Preparation of the original project; writing of the final version of the manuscript. CHS Preparation of the original project; data processing and analysis; writing of the final version of the manuscript; project administration. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The dataset generated by the linkage of the two databases related to the current study are available in the "Additional file 5". For the purpose of maintaining the confidentiality of individuals, the variables number of the Declaration of Live Birth, mother's name and home address number were excluded.

Ethics approval and consent to participate

The study was approved by the Research Ethics Committees of Hospital de Clínicas de Porto Alegre and the Research Ethics Committees of Municipal Health Department of Porto Alegre, respectively, under the respective protocol numbers 2,940,235 and 3,153,671.

The Term of Commitment for the Use of Secondary Data was signed for the ethical approval and availability of the two databases (Live Births Information System - SINASC) and the Mortality Information System - SIM)) by the Non transmissible Vital Events, Diseases and Aggravates Surveillance Team of the General Health Surveillance Coordination of the Porto Alegre Municipal Health Secretariat.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Programa de Pós Graduação em Saúde da Criança e do Adolescente - Faculdade de Medicina, Universidade Federal do Rio Grande do Sul (UFRGS), Rua Ramiro Barcelos, 2400, Porto Alegre, RS 90035003, Brazil. ²Grupo de Pesquisa e Pós-Graduação (GPPG) do Hospital de Clínicas de Porto Alegre (HCPA), Rua Ramiro Barcelos, 2400, Porto Alegre, RS 90035003, Brazil. ³Serviço de Pediatria e de Atenção Primária em Saúde, Hospital de Clínicas de Porto Alegre (HCPA), Rua Ramiro Barcelos, 2350, Porto Alegre, RS 90035007, Brazil. ⁴Departamento de Pediatria - Faculdade de Medicina, Universidade Federal do Rio Grande do Sul (UFRGS), Rua Ramiro Barcelos, 2400, Porto Alegre 90035003, RS, Brazil.

Received: 11 May 2020 Accepted: 12 January 2021

Published online: 22 January 2021

References

- Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, et al. Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the sustainable development goals. *Lancet*. 2016;388:3027–35. [https://doi.org/10.1016/S0140-6736\(16\)31593-8](https://doi.org/10.1016/S0140-6736(16)31593-8).
- Denno DM, Paul SL. Child health and survival in a changing world. *Pediatr Clin North Am*. 2017;64(4):735–54. <https://doi.org/10.1016/j.pcl.2017.03.013>.
- WHO. MCEE-WHO methods and data sources for child causes of death 2000–2017. Geneva: Department Of Evidence, Information and Research and Maternal Child Epidemiology Estimation; 2018. https://www.who.int/healthinfo/global_burden_disease/childcod_methods_2000_2017.pdf. Accessed 10 Dec 2019
- UNITED NATIONS. The millennium development goals report 2015: UN Department of Public Information; 2018. [http://www.un.org/millenniumgoals/2015_MDG_Report/pdf/MDG%202015%20rev%20\(July%2015\).pdf](http://www.un.org/millenniumgoals/2015_MDG_Report/pdf/MDG%202015%20rev%20(July%2015).pdf). Accessed 10 Dec 2019
- Barufi AM, Haddad E, Paez A. Infant mortality in Brazil, 1980–2000: a spatial panel data analysis. *BMC Public Health*. 2012;12(1):181. <https://doi.org/10.1186/1471-2458-12-181>.
- Brasil, Saúde Md, em SdV, Saúde, e DdVdDeAnT, Saúde Pd. Saúde Brasil 2014: uma análise da situação de saúde e das causas externas. Brasília: Editora do Ministério da Saúde; 2015. http://bvsm.sau.gov.br/bvsm/publicacoes/saude_brasil_2014_analise_situacao.pdf. Accessed 10 Dec 2019
- Brasil, Saúde Md, Saúde SdAa, Ações Dd, Estratégicas. Pe: Cuidados com O Recém-Nascido Pré-Termo. Brasília: Editora do Ministério da Saúde; 2011. p. 4. http://bvsm.sau.gov.br/bvsm/publicacoes/atencao_recem_nascido%20guia_profissionais_saude_v1.pdf. Accessed 10 Dec 2019
- OBSERVAPOA. Indicadores: Porto Alegre em análise; 2019. http://www.observapoa.com.br/default.php?p_secao=4. Accessed 10 Dec 2019
- Brasil, Saúde Md. A experiência Brasileira em sistemas de informação em saúde. Brasília: Editora do Ministério da Saúde; 2009. p. 2. http://bvsm.sau.gov.br/bvsm/publicacoes/experiencia_brasileira_sistemas_saude_volume1.pdf. Accessed 10 Dec 2019
- Basu AM, Stephenson R. Low levels of maternal education and the proximate determinants of childhood mortality: a little learning is not a dangerous thing. *Soc Sci Med*. 2005;60(9):2011–23. <https://doi.org/10.1016/j.socscimed.2004.08.057>.
- Silvestrin S, Silva CH, Hirakata VN, Goldani AA, Silveira PP, Goldani MZ. Maternal education level and low birth weight: a meta-analysis. *J Pediatr*. 2013;89(4):339–45. <https://doi.org/10.1016/j.jpeds.2013.01.003>.
- Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382:427–51. [https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X).
- Kimani-Murage EW, Fotso JC, Egondi T, Abuya B, Elungata P, Ziraba AK, et al. Trends in childhood mortality in Kenya: the urban advantage has seemingly been wiped out. *Health Place*. 2014;29:95–103. <https://doi.org/10.1016/j.healthplace.2014.06.003>.
- Gruebner O, Khan MMH, Burkart K, Lautenbach S, Lakes T, Krämer A, et al. Spatial variations and determinants of infant and under-five mortality in Bangladesh. *Health Place*. 2017;47:156–64. <https://doi.org/10.1016/j.healthplace.2017.08.012>.
- IPEA, PNUD, Pinheiro FJ. Atlas do Desenvolvimento Humano nas Regiões Metropolitanas Brasileiras. Brasília: IPEA, Fundação João Pinheiro, PNUD; 2014. http://www.atlasbrasil.org.br/2013/data/rawData/publicacao_atlas_rm_pt.pdf. Accessed 10 Dec 2019
- OBSERVAPOA. Geoprocessamento, Arquivos e shapes: Bairros Oficiais de Porto Alegre - 2016; 2016. http://observapoa.com.br/default.php?reg=259&p_secao=46 Accessed 10 Dec 2019
- Cruz MS, Carvalho FJV, Irffi G. Perfil socioeconômico, demográfico, cultural, regional e comportamental da gravidez na adolescência no Brasil; 2016. <http://www.ipea.gov.br/ppp/index.php/PPP/article/view/567>. Accessed 10 Dec 2019
- Koffi AK, Maina A, Yaroh AG, Habi O, Bensaïd K, Kalter HD. Social determinants of child mortality in Niger: results from the 2012 National Verbal and social autopsy study. *J Glob Health*. 2016;6(1):010603.
- Siahianidou T, Dessypris N, Analitis A, Mihis C, Evangelou E, Chrousos G, et al. Disparities of infant and neonatal mortality trends in Greece during the years of economic crisis by ethnicity, place of residence and human development index: a nationwide population study. *BMJ Open*. 2019;9(8):e025287.
- Lee AC, Cousens S, Darmstadt GL, Blencowe H, Pattinson R, Moran NF, et al. Care during labor and birth for the prevention of intrapartum-related neonatal deaths: a systematic review and Delphi estimation of mortality effect. *BMC Public Health*. 2011;11 Suppl 3:S10. <https://doi.org/10.1186/1471-2458-11-S3-S10>.
- Santos HG, Andrade SM, Silva AMR, Mathias TAF, Ferrari LL, Mesas AE. Mortes infantis evitáveis por intervenções do Sistema Único de Saúde: comparação de duas coortes de nascimentos. *Cien Saude Colet*. 2014;19:907–16. <https://doi.org/10.1590/1413-81232014193.01182013>.
- Maia LTS, Souza WW, Mendes ACG. Diferenciais nos fatores de Risco Para a mortalidade infantil em cinco cidades brasileiras: um estudo de Caso-controlado com base no SIM e no SINASC. *Cad Saude Publica*. 2012;28:2163–76. <https://doi.org/10.1590/S0102-311X2012001100016>.
- Torvie AJ, Callegari LS, Schiff MA, Debiec KE. Labor and delivery outcomes among young adolescents. *Am J Obstet Gynecol*. 2015;213(1):95.e91–8.
- Socolov DG, Iorga M, Carauleanu A, Ilea C, Bliaru I, Boiculescu L, et al. Pregnancy during adolescence and associated risks: an 8-year hospital-based cohort study (2007–2014) in Romania, the country with the highest rate of teenage pregnancy in Europe. *Biomed Res Int*. 2017;2017:9205016.
- Rousou E, Kouta C, Middleton N, Karanikola M. Single mothers' self-assessment of health: a systematic exploration of the literature. *Int Nurs Rev*. 2013;60(4):425–34. <https://doi.org/10.1111/inr.12044>.
- Gomes AG, Marin AH, Piccinini CA, Lopes RdCS. Expectativas e sentimentos de gestantes solteiras em relação aos seus bebês. *Temas em Psicologia*. 2015;23:399–411. <https://doi.org/10.9788/TP2015.2-12>.
- Cabeza de Baca T, Wojcicki JM, Epel ES, Adler NE. Lack of partner impacts newborn health through maternal depression: a pilot study of low-income immigrant Latina women. *Midwifery*. 2018;64:63–8. <https://doi.org/10.1016/j.midw.2018.05.014>.
- Gibbons L, Belizán JM, Lauer JA, Betrán AP, Meriandi M, Althabe F. The global numbers and costs of additionally needed and unnecessary caesarean sections performed per year: overuse as a barrier to universal coverage. *World Health Rep*. 2010;30:1–31 <https://www.who.int/healthsystems/topics/financing/healthreport/30C-sectioncosts.pdf>. Accessed 10 Dec 2019.
- Betrán AP, Ye J, Moller AB, Zhang J, Gülmezoglu AM, Torloni MR. The increasing trend in caesarean section rates: global, regional and National Estimates: 1990–2014. *PLoS One*. 2016;11(2):e0148343.
- Leal MC, Gama SGN, Cunha CB. Desigualdades raciais, sociodemográficas e na assistência ao pré-natal e ao parto, 1999–2001. *Rev Saude Publica*. 2005;39:100–7. <https://doi.org/10.1590/S0034-89102005000100013>.
- Freitas PF, Fernandes TM, Buxhspsa, pid SX, nrm=iso: Associação entre fatores institucionais, perfil da assistência ao parto e as taxas de cesariana em Santa Catarina. *Rev Bras Epidemiol* 2016, 19:525–538. <https://doi.org/10.1590/1980-5497201600030005>.
- Raspantini PR, Miranda MJ, ZPd S, Alencar GP, Diniz SG, Almeida MF. O impacto do tipo de hospital e tipo de parto sobre a idade gestacional ao nascer no Município de São Paulo, 2013–2014. *Rev Bras Epidemiol*. 2016;19:878–82. <https://doi.org/10.1590/1980-5497201600040016>.
- Victora CG, Aquino EM, do Carmo Leal M, Monteiro CA, Barros FC, Szwarzwald CL. Maternal and child health in Brazil: progress and challenges. *Lancet*. 2011;377(9780):1863–76. [https://doi.org/10.1016/S0140-6736\(11\)60138-4](https://doi.org/10.1016/S0140-6736(11)60138-4).
- Domingues RMSM, Dias MAB, Nakamura-Pereira M, Torres JA, d'Orsi E, Pereira APE, et al. Processo de decisão pelo tipo de parto no Brasil: da preferência inicial das mulheres à via de parto final. *Cad Saude Publica*. 2014;30:S101–16. <https://doi.org/10.1590/0102-311X00105113>.
- Cubbin C, Kim Y, Vohra-Gupta S, Margerison C. Longitudinal measures of neighborhood poverty and income inequality are associated with adverse birth outcomes in Texas. *Soc Sci Med*. 2020;245:112665.
- Viellas EF, Domingues RMSM, Dias MAB, Gama SGN, Theme Filha MM, Costa JV, et al. Assistência pré-natal no Brasil. *Cad Saude Publica*. 2014;30:S85–S100. <https://doi.org/10.1590/0102-311X00126013>.
- Magalhães ML, Cortez-Escalante J, Cannon LRC, Coimbra TS, Padilla H. Relatório 30 anos de SUS, que SUS Para 2030? In: Saúde OP-Ad. Brasília, editor. *Desafios da mortalidade infantil e na infância*; 2018.
- OBSERVAPOA. A Cidade de Porto Alegre; 2017. http://www.observapoa.com.br/default.php?p_secao=4. Accessed 10 Dec 2019

39. McCoy DC, Peet ED, Ezzati M, Danaei G, Black MM, Sudfeld CR, et al. Early childhood developmental status in low- and middle-income countries: national, regional, and global prevalence estimates using predictive modeling. *PLoS Med.* 2016;13(6):e1002034.
40. Tran TD, Luchters S, Fisher J. Early childhood development: impact of national human development, family poverty, parenting practices and access to early childhood education. *Child Care Health Dev.* 2017;43(3):415–26. <https://doi.org/10.1111/cch.12395>.
41. Ruiz JI, Nuhu K, McDaniel JT, Popoff F, Izcovich A, Criniti JM. Inequality as a powerful predictor of infant and maternal mortality around the world. *PLoS One.* 2015;10(10):e0140796.
42. Hernandez AR, Silva CH, Agranonik M, Quadros FM, Goldani MZ. Análise de tendências das taxas de mortalidade infantil e de seus fatores de Risco na cidade de Porto Alegre, Rio Grande do Sul, Brasil, no período de 1996 a 2008. *Cad Saude Publica.* 2011;27:2188–96. <https://doi.org/10.1590/S0102-311X2011001100012>.
43. Fonseca SC, Flores PVG, Camargo KR Jr, Pinheiro RS, Coeli CM. Maternal education and age: inequalities in neonatal death. *Rev Saude Publica.* 2017; 51. <https://doi.org/10.11606/s1518-8787.2017051007013>.
44. Liu X, Behrman JR, Stein AD, Adair LS, Bhargava SK, Borja JB, et al. Prenatal care and child growth and schooling in four low- and medium-income countries. *PLoS One.* 2017;12(2):e0171299.
45. WHO. World health statistics overview 2019: monitoring health for the SDGs, sustainable development goals. Geneva: World Health Organization; 2019. <https://apps.who.int/iris/bitstream/handle/10665/324835/9789241565707-eng.pdf?ua=1>. Accessed 10 Dec 2019
46. Ickes SB, Hurst TE, Flax VL. Maternal literacy, facility birth, and education are positively associated with better infant and young child feeding practices and nutritional status among Ugandan children. *J Nutr.* 2015;145(11):2578–86. <https://doi.org/10.3945/jn.115.214346>.
47. Prickett KC, Augustine JM. Maternal education and Investments in Children's health. *J Marriage Fam.* 2016;78(1):7–25. <https://doi.org/10.1111/jomf.12253>.
48. Andriano L, Monden CWS. The causal effect of maternal education on child mortality: evidence from a quasi-experiment in Malawi and Uganda. *Demography.* 2019;56(5):1765–90. <https://doi.org/10.1007/s13524-019-00812-3>.
49. Mensch BS, Chuang EK, Melnikas AJ, Psaki SR. Evidence for causal links between education and maternal and child health: systematic review. *Trop Med Int Health.* 2019;24(5):504–22. <https://doi.org/10.1111/tmi.13218>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

