

CORRECTION

Open Access



# Correction to: Modeling mortality risk effects of cigarettes and smokeless tobacco: results from the national health interview survey linked mortality file data

Esther Salazar<sup>\*</sup>, Chunfeng Ren, Brian L. Rostron and Ghideon Solomon

## Correction to: *BMC Public Health* 21, 1773 (2021)

<https://doi.org/10.1186/s12889-021-11,801-w>

It was highlighted that the in the original article [1] in Table 1 and Table S3, the labels for the "Poverty level" variable (Below 100% threshold, At or above 100% threshold) were presented in the incorrect order. That is, label "At or above 100% threshold" should have appeared first. Also, in the Results section, the following sentence should have appeared as: "Exclusive current SLT users, along with exclusive current and former smokers, were more likely to be non-Hispanic white and have less education (less than high school or high school diploma) and **higher poverty level (at or above 100% poverty threshold).**" The original article has been updated.

Published online: 04 January 2022

## Reference

1. Salazar E, et al. Modeling mortality risk effects of cigarettes and smokeless tobacco: results from the National Health Interview Survey Linked Mortality File Data. *BMC Public Health*. 2021;21:1773. <https://doi.org/10.1186/s12889-021-11801-w>.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-021-12242-1>.

**Additional file 1: Table S3.** Demographic and socioeconomic characteristics (weighted estimates and 95% CIs) for NHIS participants aged 35+ at the time of interview.<sup>a</sup>

The original article can be found online at <https://doi.org/10.1186/s12889-021-11801-w>.

\*Correspondence: [Esther.Salazar@fda.hhs.gov](mailto:Esther.Salazar@fda.hhs.gov)  
Center for Tobacco Products, U.S. Food and Drug Administration, 11785  
Beltsville Dr, Calverton, MD 20705, USA



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Table 1** Demographic and socioeconomic characteristics (weighted estimates)<sup>a</sup> for NHIS participants<sup>b</sup> aged 35+ at the time of interview

Variables	Current Smokers		Former Smokers		Never Smokers		All
	Current SLT Users	Former SLT Users	Current SLT Users	Former SLT Users	Current SLT Users	Former SLT Users	
<b>Mean age in years</b>	48.75	48.99	51.41	57.53	53.68	50.51	55.23
<b>Sex %</b>							
Male	79.11	75.25	37.56	79.68	75.81	77.10	36.25
female	20.89	24.75	62.44	20.32	24.19	22.90	63.75
<b>Race/ethnicity %</b>							
Hispanic	3.16	5.18	9.24	5.62	2.14	4.84	11.85
Non-Hispanic white	85.38	80.95	73.51	85.54	77.35	82.23	67.47
Non-Hispanic black	9.80	11.33	13.5	7.58	18.00	11.10	12.00
Non-Hispanic other	1.66	2.54	3.74	1.27	2.52	1.83	6.54
<b>Education %</b>							
Less than high school	31.85	23.05	22.31	23.66	36.76	14.18	17.70
High school diploma	35.88	40.96	38.64	32.75	29.93	23.06	29.80
Some college and higher	32.28	35.75	38.5	43.48	33.04	62.44	51.91
Missing	0.00	0.24	0.56	0.11	0.28	0.32	0.58
<b>Poverty level %</b>							
At or above 100% threshold	76.94	76.96	73.05	83.89	72.32	82.39	77.39
Below 100% threshold	13.03	16.11	14.08	6.39	14.63	7.38	9.39
Missing	10.02	6.93	12.87	9.73	13.05	10.23	13.22
<b>Body mass index (kg/m<sup>2</sup>) %</b>							
Underweight (< 18.5)	1.60	2.58	2.96	0.46	1.04	0.50	1.71
Normal weight (18.5–24.9)	30.58	31.06	40.94	21.56	21.61	19.66	35.19
Overweight (25.0–29.9)	38.56	41.31	32.14	42.77	39.64	43.31	34.35
Obese (30+)	27.27	24.00	21.25	33.26	35.49	35.63	25.49
Missing	1.99	1.05	2.72	1.95	2.22	0.89	3.27
Never Tobacco Users							54.97

<sup>a</sup> Additional file 1: Table S3, shows the 95% CIs for these weighted estimates

<sup>b</sup> Excluding participants with missing tobacco-use status, poly-users, and users of other tobacco products including pipe, hookah, e-cigarettes, bidi, and cigars. Survey years: 1987, 1991, 1992, 1994, 1998, 2000, 2005, 2010, and 2012–2014